



(12) UK Patent (19) GB (11) 2 364 059 (13) B

(45) Date of publication: 14.01.2004

(54) Title of the invention: **Beta-Secretase enzyme compositions and methods**

(51) Int Cl⁷: C07K 14/47, A01K 67/027, A61K 38/08, A61P 25/28, C07K 7/06, C12N 1/19 1/21 5/16 9/64 15/57, C12Q 1/68, G01N 33/50

(21) Application No: 0121314.9

(22) Date of Filing: 10.02.2000

(30) Priority Data:

(31) 60119571 (32) 10.02.1999 (33) US
(31) 60139172 (32) 15.06.1999 (33) US

(60) Parent of Application No(s)
0318113.8, 0318107.0, 0318112.0 under
Section 15(4) of the Patents Act 1977

(86) International Application Data:
PCT/US2000/003819 En 10.02.2000

(87) International Publication Data:
WO2000/047618 En 17.08.2000

(43) Date A Publication: 16.01.2002

(52) UK CL (Edition W):
C3H HA5 HB7E HB7M HB7T HB7V HC1
H339 H650 H655 H656 H685 H734
A5B BE B316
G1B BAX
U1S S2418 S4001

(56) Documents Cited:
EP 0855444 A WO 1998/026059 A
WO 1997/047314 A WO 1996/020725 A
Brain Res; Vol 750, pp 11-19 (1997).
Chevallier et al.
J Neurochem; Vol 66, pp 2436-2445 (1996).
Brown et al.
Proc Natl Acad Sci USA; Vol 90, pp
6796-6800 (1993). Baldwin et al.
Mol Brain Res; Vol 48, pp 206-214 (1997).
Thompson et al.
Nucl Acid Res; Vol 18, pp 7184 (1990).
Diedrich et al.

(72) Inventor(s):

John P Anderson
Guriqbal Basi
Minh Tam Doane
Normand Frigon
Varghese John
Michael Power
Sukanto Sinha
Gwen Tatsuno
Jay Tung
Shuwen Wang
Lisa McConlogue

(73) Proprietor(s):

Elan Pharmaceuticals Inc.
(Incorporated in USA - Delaware)
800 Gateway Boulevard,
South San Francisco, CA 94080,
United States of America

(74) Agent and/or Address for Service:

Kilburn & Strode
20 Red Lion Street, LONDON, WC1R 4PJ,
United Kingdom

(continued on next page)

GB 2 364 059 B - continuation

- (56) J Biol Chem; Vol 271, pp 27241-27244
(1996). Saftig et al
Protein Sci; Vol 6, pp 1458-1466 (1997).
Pavel et al
Biochem Biophys Res Commun; Vol 27, pp
157-162. Schechter et al
J Med Chem; Vol 35, pp 1702-1709. Young
et al
Nature; Vol 402, pp 537-540 (1999). Sinha
et al.
Nature; Vol 402, pp 533-537 (1999). Yan et
al
Mol Cell Neurosci; Vol 14, pp 419-427
(1999). Hussain et al
Science; Vol 286, pp 735-741 (1999).
Vassar et al.

- (58) Field of Search:
As for published application 2364059 A viz:
Other: **NOT YET ADVISED**
updated as appropriate

Additional Fields
INT CL⁷ A01K, A61K, C07K, C12N, C12Q,
G01N
Other: EPO-Internal, BIOSIS, WPI Data,
PAJ, MEDLINE, SCISEARCH, CHEM ABS
Data,
BIOTECHNOLOGY ABS, EMBASE

ATGGCCCAAGCCCTGCCCTGGCTCCTGCTGTGGATGGGCGCGGGAGT
GCTGCCTGCCCACGGCACCCAGCACGGCATCCGGCTGCCCCCTGCGCA
GCGGCCTGGGGGGCGCCCCCTGGGGCTGCGGCTGCCCCGGGAGAC
CGACGAAGAGCCCGAGGAGCCCGGCCGGAGGGGCAGCTTTGTGGAGA
TGGTGGACAACCTGAGGGGCAAGTCGGGGCAGGGGCTACTACGTGGAG
ATGACCGTGGGCAGCCCCCGCAGACGCTCAACATCCTGGTGGATACA
GGCAGCAGTAACTTTGCAGTGGGTGCTGCCCCCACCCTTCCCTGCAT
CGCTACTACCAGAGGCAGCTGTCCAGCACATACCGGGACCTCCGGAAG
GGTGTGTATGTGCCCTACACCCAGGGCAAGTGGGAAGGGGAGCTGGG
CACCGACCTGGTAAGCATCCCCCATGGCCCCAACGTCACTGTGCGTGC
CAACATTGCTGCCATCACTGAATCAGACAAGTTCTTCATCAACGGCTCC
AACTGGGAAGGCATCCTGGGGCTGGCCTATGCTGAGATTGCCAGGCCT
GACGACTCCCTGGAGCCTTTCTTTGACTCTCTGGTAAAGCAGACCCACG
TTCCCAACCTCTTCTCCCTGCAGCTTTGTGGTGCTGGCTTCCCCCTCAA
CCAGTCTGAAGTGCTGGCCTCTGTCTGGAGGGAGCATGATCATTGGAGG
TATCGACCACTCGCTGTACACAGGCAGTCTCTGGTATACACCCATCCGG
CGGGAGTGGTATTATGAGGTGATCATTGTGCGGGTGGAGATCAATGGA
CAGGATCTGAAAATGGACTGCAAGGAGTACAACCTATGACAAGAGCATTG
TGGACAGTGGCACCACCAACCTTCGTTTGCCCAAGAAAGTGTTTGAAGC
TGCAGTCAAATCCATCAAGGCAGCCTCCTCCACGGAGAAGTTCCCTGAT
GGTTTCTGGCTAGGAGAGCAGCTGGTGTGCTGGCAAGCAGGCACCACC
CCTTGGAACATTTTCCCAGTCATCTCACTCTACCTAATGGGTGAGGTTAC
CAACCAGTCCTTCCGCATCACCATCCTTCCGCAGCAATACCTGCGGCCA
GTGGAAGATGTGGCCACGTCCCAAGACGACTGTTACAAGTTTGCCATCT
CACAGTCATCCACGGGCACTGTTATGGGAGCTGTTATCATGGAGGGCTT
CTACGTTGTCTTTGATCGGGCCCCGAAAACGAATTGGCTTTGCTGTCAGC
GCTTGCCATGTGCACGATGAGTTCAGGACGGCAGCGGTGGAAGGCCCT
TTTGTACCTTGGACATGGAAGACTGTGGCTACAACATTCCACAGACAG
ATGAGTCAACCCTCATGACCATAGCCTATGTCATGGCTGCCATCTGCGC
CCTCTTCATGCTGCCACTCTGCCTCATGGTGTGTCAGTGGCGCTGCCTC
CGCTGCCTGCGCCAGCAGCATGATGACTTTGCTGATGACATCTCCCTGC
TGAAG

FIG. 1A

CCATGCCGGCCCCCTCACAGCCCCGCCGGGAGCCCCGAGCCCCGCTGCCCAGGCTC
CGCCGCSGTGCCGATGTAGCGGGCTCCGGATCCCAGCCTCTCCCCTGCTCCCG
TCTGCGGATCTCCCCTGACCGCTCTCCACAGCCCCGACCCGGGGGCTGGCCCA
GCCCTGCAGGCCCTGGCGTCCTGATGCCCCCAAGCTCCCTCTCCTGAGAAGCCA
AGCACCAACCAGACTTGGGGGCAGGCGCCAGGGACGGACGTGGGCCAGTGCG/
CCAGAGGGCCCCGAAGGCCGGGGCCCCACCATGGCCCCAAGCCCTGCCCTGGCTCC
CTGTGGATGGGCGCGGGAGTGCTGCCTGCCACGGCACCCAGCACGGCATCCG
TGCCCTGCGCAGCGGCCTGGGGGGCGCCCCCTGGGGCTGCGGCTGCCCCG
AGACCGACGAAGAGCCCCGAGGAGCCCCGCCGGAGGGGGCAGCTTTGTGGAGATC
GGACAACCTGAGGGGCAAGTCGGGGCAGGGCTACTACGTGGAGATGACCGTGG
AGCCCCCGCAGACGCTCAACATCCTGGTGGATACAGGCAGCAGTAACCTTTGCA
GGGTGCTGCCCCCACCCTTCTGTCATCGCTACTACCAGAGGCAGCTGTCCAG
CATACCGGGACCTCCGGAAGGGTGTGTATGTGCCCTACACCCAGGGCAAGTGGC
GGGGAGCTGGGCACCGACCTGGTAAGCATCCCCCATGGCCCCAAGCTCACTGT
TGCCAACATTGCTGCCATCACTGAATCAGACAAGTTCTTCATCAACGGCTCCAAC
GAAGGCATCCTGGGGCTGGCCTATGCTGAGATTGCCAGGCCTGACGACTCCCTG
GCCTTTCTTTGACTCTCTGGTAAAGCAGACCCACGTTCCCAACCTCTTCTCCCTG
CTTTGTGGTGCTGGCTTCCCCCTCAACCAGTCTGAAGTGCTGGCCTCTGTCCGA
GAGCATGATCATTGGAGGTATCGACCACTCGCTGTACACAGGCAGTCTCTGGTA
ACCCATCCGGCGGGAGTGGTATTATGAGGTGATCATTGTGCGGGTGAGATCAA
GACAGGATCTGAAAATGGACTGCAAGGAGTACAACCTATGACAAGAGCATTGTGG
GTGGCACCAACCAACCTTCGTTTGCCCAAGAAAGTGTGTTGAAGCTGCAGTCAAATC
TCAAGGCAGCCTCCTCCACGGAGAAGTTCCTGATGGTTTCTGGCTAGGAGAGC
CTGGTGTGCTGGCAAGCAGGCACCAACCCTTGGAAACATTTTCCAGTCATCTCAC
TACCTAATGGGTGAGGTTACCAACCAGTCTTCCGCATCACCATCCTTCCGCAGC
TACCTGCGGCGAGTGGAAGATGTGGCCACGTCCCAAGACGACTGTTACAAGTTT
ATCTCACAGTCATCCACGGGCACTGTTATGGGAGCTGTTATCATGGAGGGCTTCT
GTTGTCTTTGATCGGGCCCCGAAAACGAATTGGCTTTGCTGTCAGCGCTTGCCATG
CACGATGAGTTCAGGACGGCAGCGGTGGAAGGCCCTTTTGTACCTTGGACATG
AGACTGTGGCTACAACATTCCACAGACAGATGAGTCAACCCTCATGACCATAGCC
TGTCATGGCTGCCATCTGCGCCCTCTTCATGCTGCCACTCTGCCTCATGGTGTGT
GTGGCGCTGCCTCCGCTGCCTGCGCCAGCAGCATGATGACTTTGCTGATGACAT
CCCTGCTGAAGTGAGGAGGCCCATGGGCAGAAGATAGAGATTCCCCTGGACCAC
CTCCGTGGTTCACTTTGGTCACAAGTAGGAGACACAGATGGCACCTGTGGCCAG
CACCTCAGGACCCTCCCCACCCACCAATGCCTCTGCCTTGATGGAGAAGGAAA
GCTGGCAAGGTGGGTTCCAGGGACTGTACCTGTAGGAAACAGAAAAGAGAAGAA
AAGCACTCTGCTGGCGGGAATACTCTTGGTCACCTCAAATTTAAGTCGGGAAATT
GCTGCTTGAACTTCAGCCCTGAACCTTTGTCCACCATTCTTTAAATTCTCCAAC
AAAGTATTCTTCTTTTCTTAGTTTCAGAAGTACTGGCATCACACGCAGGTTACCTT
CGTGTGTCCCTGTGGTACCCTGGCAGAGAAGAGACCAAGCTTGTTTCCCTGCTG
CAAAGTCAGTAGGAGAGGATGCACAGTTTGCTATTTGCTTTAGAGACAGGGACTC
TAAACAAGCCTAACATTGGTGCAAAGATTGCCTCTTGAATT

FIG 1R

MAQALPWLLWWMGAGVLP AHGTQH GIRLPLRSGLG G APLGLRLP
RETDEEPEEPGRRGSFVEMVDNLRGKSGQGYVEMTVGSP PQT
LNILVDTGSSNFAVGAAPHPFLHRYYQRQLSSTYRDLRKG VYVPY
TQGWEGELGTDLV SIPHGPNVTVRANIAAITESDKFFINGSNWE
GILGLAYAEIARPDDSLEPFFDSL VKQTHVPNLFSLQLCGAGFPLN
QSEVLASVGGSMIIGGIDHSLYTGSLWYTPIRREWYYEVIIVRVEIN
GQDLKMDCKEYNYDKSIVDSGTTNLR LPPKKVFEAAVKS IKAASST
EKFPDGFWLGEQLVCWQAGTTPWNIFPVISLYLMGEVTNQSFRT
ILPQQYL RPVEDVATSQDDCYKFAISQSSTGTVMGAVIMEGFYV
FDRARKRIGFAVSACHVHDEFRTAAVEGPFVTLDMEDCGYNIPQ
TDESTLMTIAYVMAAICALFMLPLCLMVCQWRCLRCLRQQHDDF
ADDISLLK

FIG. 2A

ETDEEPEEPGRRGSFVEMVDNLRGKSGQGYYVEMTVGSPPQT
LNILVDTGSSNFAVGAAPHPFLHRYYQRQLSSTYRDLRKGVYVP
TQGWEGELGTDLVSIHPGPNVTVRANIAAITESDKFFINGSNWE
GILGLAYAEIARPDDSLEPFFDSL VKQTHVPNLFSLQLCGAGFPLN
QSEVLASVGGSMIIGGIDHSLYTGSLWYTPIRREWYYEVIIVRVEIN
GQDLKMDCKEYNYDKSIVDSGTTNLRLPKKVFEEAAVKSIIKAASST
EKFPDGFWLGEQLVCWQAGTTPWNIFPVISLYLMGEVTNQSFRIT
ILPQQYL RPVEDVATSQDDCYKFAISQSSTGTVMGAVIMEGFYVW
FDRARKRIGFAVSACHVHDEFRTAAVEGPFVTLDMEDCGYNIPQ
TDESTLMTIAYVMAAICALFMLPLCLMVCQWRCLRCLRQQHDDF
ADDISLLK

FIG. 2B

FIG. 3A

MAQALPWLLLWMGAGVLP AHGTQH GIRLPLRSGLG GAPLGLRLPRETDEEPE
EPGRRGSFVEMVDNLRGKSGQGYVEMT
VGSPQTLNILVD TGSSNFAVGAAPHPFLHRYYQRQLSSTYRDLRKGVVYPYT
QGKWE GELGTDLV SIPHGPNVTVRANI
AAITESDKFFINGSNWEGILGLAYAEIARPDDSLEPFFDSL VKQTHV PNLFSLQL
CGAGFPLNQSEVLASVGGSMIIGGI
DHSLYTGSLWYTPIRREWYVEIIVRVEINGQDLKMDCKEYNYDKSIVDSGTTNL
RLPKKVFEAAVKSIIKAASSTEKFPD
GFWLGEQLVCWQAGTTPWNIFPVISLYLMGEVTNQSFRITILPQQYL RPVEDVA
TSQDDCYKFAISQSSTGTVMGAVIME
GFYVVFDRARKRIGFAVSACHVHDEFRTAAVEGPFVTLDMEDCGYNIPQTDED
YKDDDDK

FIG. 3B

ETDEEPEEPGRRGSFVEMVDNLRGKSGQGYVEMT
VGSPQTLNILVD TGSSNFAVGAAPHPFLHRYYQRQLSSTYRDLRKGVVYPYT
QGKWE GELGTDLV SIPHGPNVTVRANI
AAITESDKFFINGSNWEGILGLAYAEIARPDDSLEPFFDSL VKQTHV PNLFSLQL
CGAGFPLNQSEVLASVGGSMIIGGI
DHSLYTGSLWYTPIRREWYVEIIVRVEINGQDLKMDCKEYNYDKSIVDSGTTNL
RLPKKVFEAAVKSIIKAASSTEKFPD
GFWLGEQLVCWQAGTTPWNIFPVISLYLMGEVTNQSFRITILPQQYL RPVEDVA
TSQDDCYKFAISQSSTGTVMGAVIME
GFYVVFDRARKRIGFAVSACHVHDEFRTAAVEGPFVTLDMEDCGYNIPQTDED
YKDDDDK


GEC of recombinant β -secretase

FIG. 4




FIG. 5

8/26




FIG. 6A

FIG. 6B

9/26




FIG. 7




FIG. 8

C




FIG. 9

501
501

FIG. 10

CTGTTGGGCTCGCGGTTGAGGACAAACTCTTCGCGGTCTTTCCAGTACTCT
 TGGATCGGAAACCCGTCCGGCCTCCGAACGGTACTCCGCCACCGAGGGACCT
 GAGCGAGTCCGCATCGACCGGATCGGAAAACCTCTCGACTGTTGGGGTGA
 TACTCCCTCTCAAAAGCGGGCATGACTTCTGCGCTAAGATTGTCAGTTTC
 AAAAACGAGGAGGATTTGATATTCACCTGGCCCCGCGGTGATGCCTTTGAG
 GTGGCCGCGTCCATCTGGTCAGAAAAGACAATCTTTTGTGTCAAGCTT
 AGGTGTGGCAGGCTTGAGATCTGGCCATACACTTGAGTGACAATGACATC
 ACTTTGCCTTTCTCTCCACAGGTGTCCACTCCAGGTCCAACCTGCAGGTC
 ACTCTAGACCC

FIG. 11A




FIG. 11B




FIG. 12

1 TTCTCATGTTTCACAGCTTATCATGTCAGATCCCGCAAGCTGTTGTCATTTCTTCCAGGCGCAGAACTGGTAGGTATGGAAGATCCGATGTACCGGCCAGATATAT.

107 CGGTGACATTGATTATTCAGTACTGATTATTAATAGTAATCAUATTAAGGGTCATTAGTTCATAGCCCATATATGAGTTCCGGGTACATAAATTAAGGTAATAAT

213 CCCCCTGCGTACGCGCCCAAGCAGCCCGCCGCCCATTTGAGCTCAATAATGACGTATGTTCCCATAGTAAGCCCAATAGGACTTTCCATTGACGTCAATGGTGGAA

319 ATTTAAGGTAAACTGCCCCTTGGCAGTACATCAAGTGTATCATATGCCAAGTAAGCCGCCCTATTGAGCTCAATGAGGGTAAATGGCCCGCCCTGCCATTATGCC

425 GTACATGACCTTATGGGACTTTCTCTACTTGGCAGTACATCTAGTATTAGTCATGCTATTACCATGGTCATGGGTTTTCGGCAGTACATCAATGGCGGTGGAT

531 CCGTTTGACTCAGCGGGATTTCGAGTCTCCAGCCCATTTGACGTCAATGGGAGTTTGTGTTTGGCAACAAAATCAAGCGGACTTTCCAAAATGTGGTAACAACTC

637 CCCCATTGACCCAAATGGCCCGTGAAGCGTGTAGCGTGGAGGCTCTATATAACGAGAGCTCTCTGGCTAACTAGAGAACCCACTGCTTACTGCTTATGGAAATTT

743 TAGGACTCACTATAGGAGACCCAACTCTGTTGGGCTGGGGTTGAGGACAACTCTTGGGGTCTTTCCAGTACTCTTGATAGGAAACCGGTGGGCTGGG

849 CCGTACTCGGCAACCGGAGGAGCTGAGCGAGTCCCATCGAGCCGATCGGAAACCTCTCGACTGTTGGGGTGAGTACTCCCTCTCAAAAGCGGCGATGACTTC

955 CCTAAGATTGTCACTTTCCAAAAGGAGGAGATTGATATTCACTGGCCCGCGGGTATGCTTTGAGGGTGGCCCGGCTCCATCTGGTCAGAAAGCAATC

1061 TTGTGTCTCAAGCTTGAAGTGTGGCAGGCTGAGATCTGGCCATACACTTGAGTCAAAATGACATCCACTTTGCCCTTCTCTCCAGGCTGTCACCTCCCAAGT

1167 AACTCCAGGTGAGCTCTAGACCCCGGAATTCGACATATCCAATCACTGCGGCACTCGTCCCAAGCCCGCCCGGAGCTGGGAGCCCGCACTGGATTAAT

1273 TGGCTGAGCGAGCAACCGAGCCCGAGGAGCCCGAGGAGCCCTTGGGCTGCGCCGCTGCGGCGCGCGCGCGGAGCAGGGAAAGCCCGCAAGCGCCCGCATGCT

1379 CCCCCTCCAGCCCGCCCGGAGCCCGCCCGCCCGCTGGCCAGGCTGGCCCGCCCGCTGGCGATGTAGCGGCTCCCGATCCCAAGCTCTCCCTGCTCCCGTCT

1485 CGGATCTCCCTGAGCCCTCTCCAGCGCCCGGAGCCCGGCGGCTGGCCAGGCGCTCGAGGCTCTGATGCGCCCAAGCTCCCTCTCTCTGAGAAAG

1591 CAGCAACCAACAGACTTGGGGGCAAGCGCCAGGAGCGAGCTGGGCGAGTGGGAACCCAGAGCGCCCGGAAGCCCGGGGCCACC ATG GGC CAA GGC CT

1690 CCC TGG CTC CTG CTG TGG ATG GGC GCG GGA GTG CTG CCT GGC CAC GGC ACC CAG CAC GGC ATC GGC CTG CCC CTG GGC

6P Pro Trp Leu Leu Leu Trp Met Gly Ala Gly Val Leu Pro Ala His Gly Thr Gln His Gly Ile Arg Leu Pro Leu Arg

1768 AOC GGC CTG GGC GGC GGC GGC CTG GGC CTG GGC CTG GGC GGC GAG AOC GAC GAA GAG GGC GAG GAG GGC GGC GGC AOC

32P Ser Gly Leu Gly Gly Ala Pro Leu Gly Leu Arg Leu Pro Arg Gu Thr Asp Glu Glu Pro Glu Glu Pro Gly Arg Arg

1846 GGC AOC TTT GTG GAG ATG GTG GAC AAC CTG AOG GGC AAG TCG GGC CAG GGC TAC TAC GTG GAG ATG AOC GTG GGC AOC

58P Gly Ser Phe Val Glu Met Val Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu Met Thr Val Gly Ser

1924 GGC GGC CAG AOG CTC AAC ATC CTG GTG GAT ACA GGC AOC AGT AAC TTT GCA GTG GGT OCT GGC GGC CAC GGC TTC CTG

84P Pro Pro Gln Thr Leu Asn Ile Leu Val Asp Thr Gly Ser Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu

2002 CAT GGC TAC TAC CAG AOG CAG CTG TCC AOC ACA TAC GGC GAG CTC GGC AAG GGT GTG TAT GTG CCC TAC ACC CAG GGC

110P His Arg Tyr Tyr Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val Tyr Val Pro Tyr Thr Gln Glu

2080 AAG TGG GAA GGC GAG CTG GGC ACC GAC CTG GTA AOC ATC GGC CAT GGC GGC AOC GTC ACT GTG GGT GGC AAC ATT GGC

136P Lys Trp Glu Gly Glu Leu Gly Thr Asp Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg Ala Asn Ile Ala

2158 GGC ATC ACT GAA TCA GAC AAG TTC TTC ATC AAC GGC TCC AAC TGG GAA GGC ATC CTG GGC CTG GGC TAT OCT GAG AT

162P Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser Asn Trp Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile

2236 GGC AOG CCT GAC GAC TCC CTG GAG OCT TTC TTT GAC TCT CTG GTA AAG CAG AOC CAC GTT GGC AAC CTC TTC TCC CT

188P Ala Arg Pro Asp Asp Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr His Val Pro Asn Leu Phe Ser Leu

2314 CAG CTT TGT GGT OCT GGC TTC CCC CTC AAC CAG Ser TCT GAA GTG CTG GGC TCT GTC GGA GGC AOC ATC ATC ATT GCA G

214P Gln Leu Cys Gly Ala Gly Phe Pro Leu Asn Gln Ser Glu Val Leu Ala Gly Thr Val Gly Tyr Ser Met Ile Ile Gly Glu

2392 ATC GAC CAC TGG CTG TAC ACA GGC AGT CTC TGG TAT ACA GGC ATC GGC GGC GAG TGG TAT TAT GAG GTC ATC ATT G

240P Ile Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro Ile Arg Arg Glu Trp Tyr Tyr Glu Val Ile Ile Val

2470 GGC GTG GAG ATC AAT GCA CAG GAT CTG AAA ATG GAC TCC AAG CAG TAC AAC TAT GAC AAG AOC ATT GTG GAC AGT G

266P Arg Val Glu Ile Asn Gly Gln Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys Ser Ile Val Asp Ser G

2548 AOC AOC AAC CTT GGT TTG GGC AAG AAA GTG TTT GAA GGT GCA GTC AAA TCC ATC AAG GCA GGC TCC TCC AOC GAG A

292P Thr Thr Asn Leu Arg Leu Pro Lys Lys Val Phe Glu Ala Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu L

2626 TTC CCT CAT GGT TTC TGG CTA GGA GAG CAG CTG GTG TCC TGG CTA GCA GGC AOC ACC OCT TGG AAC ATT TTT CCA G

318P Phe Pro Asp Gly Thr Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala Ala Gly Thr Pro Trp Asn Ile Thr Pro Val

[illegible]

[illegible]

FIG. 13D

[illegible]

15414 CTGCCAACCTGCTCATGAGCAGCAACGGTCAGTCGGCTCCGTCGCCCAACCTCTCGATGAGCTGATGCTTTGGGCGGAGCACTGGCCCGAAGTCCGCCACCTCGTGC

15147 GCGGATTTCGGCTCCACAAATGCTCTGACCGGAGCATGCCCGCATACAGCGGTCAATTGACTCGAGCGGACGGATGTTTGGGCGATTCCCAATACGAGGTCGCCAA

15253 TCTTCTCTCGAGCGGGTGCTTTGGGCGGTATCGAGCCAGCAGCGGGCTACTTTCAGCGCGAGCCATCGCGAGCTTCGAGGATCCCGCGCGCTCGCGGGGTATATG

15359 GCGCATTCGCTCTGACCACTCTATCAGAGCTTGGTTGACCGCAATTTCGATCATCCACTTTGCCCGCAGGTCGATGCCACCGCAATGCTCGGATCCCGAGCGCG

15465 ACTGTGCGCGGTACACAAATCCCGCGCGAGAGCGGGCGGCTCTCGAGCGCATTCGCTGTGAGAGTACTGCCCGTAGTGCAAAACCGGAGATGCCCGAGGCTAACT

15571 AAACACCGAGCAGAGCAATACCGGAGCGAGCGGGCGCTATGAGCGCATATAAAGACAGCAATAAAGCGCAAGGGTGTTCGGTGGTTTGTGTCATAAAGCGGGGGTT

15677 GGTGCCACGGCTTCGCTCTCTGTCGATATGCCCGAGAGCGCCATTCGGGCGCAATAGCGGGGGGTTTCTTCTGCTTTTGGCGACCGCAACCGCGCAAGTTCCGGTGAG

15783 CCGAGCGCTCCGACCGCAACGTTGGGGCGGGCGCGCGCTGCGATAGCGACTGGCGGGTGGGTTAGCGACCGGGTGGCGCATCGGCAATGCTTTATGTTGGTGGTGGG

15889 TTAATATTTTGGGGTTGGTGGGGTCTGGTCCAGGACTGACTGAGCGACAGAGCCCATGCTTTTGGATGGCGCTGGCGCATCGACCGCATGTACTGGGGGACA

15995 GACACCGGGGGTCTGTGCTGCCAAACCGGGCGAGCGCGCAAAACCAACCGGGCGGATTTCTGGGGTCCAGCTAGTCGACCA

SalI

CTGTTGGGCTCGCGGTTGAGGACAACTCTTCGCGGTCTTTCCAGTACTCTTGGATCGGAAAC
 CCGTCGGCCTCCGAACGGTACTCCGCCACCGAGGGACCTGAGCGAGTCCGCATCGACCGGAT
 CGGAAAACCTCTCGACTGTTGGGGTGAGTACTCCCTCTCAAAGCGGGCATGACTTCTGCGCT
 AAGATTGTCAGTTTCCAAAAACGAGGAGGATTTGATATTACCTGGCCCGCGGTGATGCCTTT
 GAGGGTGGCCGCGTCCATCTGGTCAGAAAAAGACAATCTTTTGTGTCAAGCTTGAGGTGTGG
 CAGGCTTGAGATCTGGCCATACACTTGAGTGACAATGACATCCACTTGCCTTTCTCTCCACAG
 GTGTCCACTCCCAGGTCCAACCTGCAGGTCGACTCTAGACCC

FIG. 14A




FIG. 14B




FIG. 15A




FIG. 15B




FIG. 16A

FIG. 16B





FIG. 17A

FIG. 17B

**FIG. 18**




FIG. 19A

Wild-Type SequenceVal-Lys-Met-Asp...
Swedish SequenceVal-Asn-Leu-Asp...

FIG. 19B




FIG. 20




FIG. 21

β-SECRETASE ENZYME COMPOSITIONS AND METHODS

5 Field of the Invention

The invention relates to the discovery of various active forms of β-secretase, an enzyme that cleaves β-amyloid precursor protein (APP) at one of the two cleavage sites necessary to produce β-amyloid peptide (Aβ). The invention also relates to inhibitors of this
10 enzyme, which are considered candidates for therapeutics in the treatment of amyloidogenic diseases such as Alzheimer's disease. Further aspects of the present invention include screening methods, assays, and kits for discovering such therapeutic inhibitors, as well as diagnostic methods for determining whether an individual carries a mutant form of the enzyme.

15

Background of the Invention

Alzheimer's disease is characterized by the presence of numerous amyloid plaques and neurofibrillary tangles present in the brain, particularly in those regions of the brain involved in memory and cognition. β-amyloid peptide (Aβ) is a 39-43 amino acid peptide
20 that is major component of amyloid plaques and is produced by cleavage of a large protein known as the amyloid precursor protein (APP) at a specific site(s) within the N-terminal region of the protein. Normal processing of APP involves cleavage of the protein at point 16-17 amino acids C-terminal to the N-terminus of the β-AP region, releasing a secreted ectodomain, α-sAPP, thus precluding production of β-AP. Cleavage by β-secretase enzyme
25 of APP between Met⁶⁷¹ and Asp⁶⁷² and subsequent processing at the C-terminal end of APP produces Aβ peptide, which is highly implicated in the etiology of Alzheimer's pathology (Seubert, *et al.*, in *Pharmacological Treatment of Alzheimer's disease*, Wiley-Liss, Inc., pp. 345-366, 1997; Zhao, J., *et al.* *J. Biol. Chem.* 271: 31407-31411, 1996).

It is not clear whether β-secretase enzyme levels and/or activity is inherently higher
30 than normal in Alzheimer's patients; however, it is clear that its cleavage product, Aβ peptide, is abnormally concentrated in amyloid plaques present in their brains. Therefore, it would be desirable to isolate, purify and characterize the enzyme responsible for the

pathogenic cleavage of APP in order to help answer this and other questions surrounding the etiology of the disease. In particular, it is also desirable to utilize the isolated enzyme, or active fragments thereof, in methods for screening candidate drugs for ability to inhibit the activity of β -secretase. Drugs exhibiting inhibitory effects on β -secretase activity are expected to be useful therapeutics in the treatment of Alzheimer's disease and other amyloidogenic disorders characterized by deposition of A β peptide containing fibrils.

U.S. Patent 5,744,346 (Chrysler, *et al.*) describes the initial isolation and partial purification of β -secretase enzyme characterized by its size (apparent molecular weight in the range of 260 to 300 kilodaltons when measured by gel exclusion chromatography) and enzymatic activity (ability to cleave the 695-amino acid isotype of β -amyloid precursor protein between amino acids 596 and 597). The present invention provides a significant improvement in the purity of β -secretase enzyme, by providing a purified β -secretase enzyme that is at least 200 fold purer than that previously described. Such a purified protein has utility in a number of applications, including crystallization for structure determination. The invention also provides methods for producing recombinant forms of β -secretase enzymes that have the same size and enzymatic profiles as the naturally occurring forms. It is a further discovery of the present invention that human β -secretase is a so-called "aspartyl" (or "aspartic") protease.

Summary of the Invention

This invention is directed to a protein purified to apparent homogeneity comprising a segment of a β -secretase enzyme protein lacking the signal sequence (amino acid residues 1-21 with respect to SEQ ID NO:2) and lacking the putative pro region (amino acid residues 22-45 with respect to SEQ ID NO:2).

The protein may be characterized by a specific activity of at least about 0.2×10^5 and preferably at least 1.0×10^5 nM/h/ μ g protein in a representative β -secretase assay, the MBP-C125sw substrate assay. The protein, which has a characteristic activity in cleaving the 695-amino acid isotype of β -amyloid precursor protein (β -APP) between amino acids 596 and 597 thereof, may be at least 10,000-fold, preferably at least 20,000-fold and, more preferably in excess of 200,000-fold higher specific activity than an activity exhibited by a solubilized but unenriched membrane fraction from human 293 cells, such as have been earlier characterized.

In one embodiment, the purified enzyme is fewer than 450 amino acids in length, comprising a polypeptide having the amino acid sequence SEQ ID NO: 70 [63-452]. The purified protein exists in a variety of "truncated forms" relative

to the proenzyme referred to herein as SEQ ID NO: 2 [1-501], such as forms having amino acid sequences SEQ ID NO: 70 [63-452], SEQ ID NO: 69 [63-501], SEQ ID NO: 67 [58-501], SEQ ID NO: 68 [58-452], SEQ ID NO: 58 [46-452]. More generally, it has been found that particularly useful forms of the enzyme, particularly with regard to the crystallization studies described herein, are characterized by an N-terminus at position 46 with respect to SEQ ID NO: 2 and a C-terminus between positions 452 and 470 with respect to SEQ ID NO: 2. These forms are considered to be cleaved in the transmembrane "anchor" domain. Another particularly useful purified form of the enzyme is SEQ ID NO: 43 [46-501]. More generally, it is appreciated that useful forms of the enzyme have an N-terminal residue corresponding to a residue selected from the group consisting of residues 46, 58 and 63 with respect to SEQ ID NO: 2 and a C-terminus selected from a residue between positions 452 and 501 with respect to SEQ ID NO: 2 or a C-terminus between residue positions 452 and 470 with respect to SEQ ID NO: 2.

This invention is further directed to a crystalline protein composition formed from a purified β -secretase protein, as described above. According to one embodiment, the purified protein is characterized by an ability to bind to the β -secretase inhibitor substrate P10-P4'sta D->V which is at least equal to an ability exhibited by a protein having the amino acid sequence SEQ ID NO: 71 [46-419], when the proteins are tested for binding to said substrate under the same conditions. According to another embodiment, the purified protein forming the crystallization composition is characterized by a binding affinity for the β -secretase inhibitor substrate SEQ ID NO: 72 (P10-P4'sta D->V) which is at least 1/100 of an affinity exhibited by a protein having the amino acid sequence SEQ ID NO: 43 [46-501], when said proteins are tested for binding to said substrate under the same conditions. Proteins forming the crystalline composition may be glycosylated or deglycosylated.

The crystalline protein composition of the invention may further comprise a β -secretase substrate selected from the group consisting of MBP-C125wt, MBP-C125sw, APP, APPsw, and β -secretase-cleavable fragments thereof or a β -secretase inhibitor molecule, examples of which are provided herein, particularly exemplified by peptide-derived inhibitors such as SEQ ID NO: 78, SEQ ID NO: 72, SEQ ID NO: 81, and derivatives thereof. Generally useful inhibitors in this regard will have a K_i of no more than about 50 μ M to 0.5 mM.

Another aspect of the invention is directed to an isolated protein, comprising a

polypeptide that (i) is fewer than about 450 amino acid residues in length, (ii) includes an amino acid sequence that is at least 90% identical to SEQ ID NO: 75 [63-423] including conservative substitutions thereof, and (iii) exhibits β -secretase activity, as evidenced by an ability to cleave a substrate selected from the group consisting of the 695 amino acid isotype of beta amyloid precursor protein (β APP) between amino acids 596 and 597 thereof, MBP-C125wt and MBP-C125sw. Peptides which fit these criteria include, but are not limited to a polypeptide which includes or has the sequence SEQ ID NO: 75 [63-423]. The amino acid sequence may be SEQ ID NO: 58 [46-452] or at least 95% thereto, and may also include conservative substitutions within such sequences.

10 According to a further embodiment, the invention includes isolated protein compositions, such as those described above, in combination with a β -secretase substrate, such as MBP-C125wt, MBP-C125sw, APP, APPsw, and β -secretase-cleavable fragments thereof or a β -secretase inhibitor molecule. Additional β -secretase-cleavable fragments useful in this regard are described in the specification hereof. Particularly useful
15 inhibitors include peptides derived from or including SEQ ID NO: 78, SEQ ID NO: 81 and SEQ ID NO: 72. Generally, such inhibitors will have K_i s of less than about 1 μ M. Such inhibitors may be labeled with a detectable reporter molecule. Such labeled molecules are particularly useful, for example, in ligand binding assays.

In accordance with a further aspect, the invention includes protein compositions,
20 such as those described above, expressed by a heterologous cell. In accordance with a further embodiment, such cells may also co-express a β -secretase substrate or inhibitor protein or peptide. One or both of the expressed molecules may be heterologous to the cell.

In a related embodiment, the invention includes antibodies that bind specifically
25 to a β -secretase protein comprising a polypeptide that includes an amino acid sequence that is at least 90% identical to SEQ ID NO: 75 [63-423] including conservative substitutions thereof, but which lacks significant immunoreactivity with a protein having a sequence selected from the group consisting of SEQ ID NO: 2 [1-501] and SEQ ID NO: 43 [46-501]. The antibody may be reactive with a protein selected from the group
30 consisting of SEQ ID NO: 67 [58-501], SEQ ID NO: 69 [63-501], SEQ ID NO: 58 [46-452], SEQ ID NO: 68 [58-452] and SEQ ID NO: 70 [63-452].

In a further aspect, the invention provides isolated nucleic acids comprising a

sequence of nucleotides that encodes a β -secretase protein that is at least 95% identical to a protein selected from the group consisting of SEQ ID NO: 43 [46-501], SEQ ID NO: 58 [46-452], SEQ ID NO: 67 [58-501], SEQ ID NO: 68 [58-452], SEQ ID NO: 69 [63-501], SEQ ID NO: 70 [63-452], SEQ ID NO: 75 [63-423], and SEQ ID NO: 71 [46-419],
5 or a complementary sequence of any of such nucleotides, and specifically excluding a nucleic acid encoding a protein having the sequence SEQ ID NO: 2 [1-501].

Additionally, the invention includes an expression vector comprising such isolated nucleic acids operably linked to the nucleic acid with regulatory sequences effective for expression of the nucleic acid in a selected host cell, for heterologous
10 expression. The invention also includes a heterologous cell transfected with such an expression vector which expresses a biologically-active β -secretase. The cell can be a bacterial cell or a eukaryotic cell, such as an insect cell or a yeast cell. The invention also provides a method of producing a recombinant β -secretase enzyme, comprising
15 culturing a cell of the invention under conditions to promote growth of said cell and subjecting an extract or cultured medium from said cell to an affinity matrix containing a β -secretase inhibitor molecule. The matrix may contain an antibody characterised by an ability to bind β -secretase, such as an antibody of the invention.

Also provided is a heterologous cell, comprising (i) a nucleic acid molecule encoding an active β -secretase protein of the present invention or a nucleic acid molecule
20 encoding the active β -secretase protein of SEQ ID NO:2 [1-501];
(ii) a nucleic acid molecule encoding a β -secretase substrate molecule selected from the group consisting of MBP-C125wt, MBP-C125sw, APPwt, APPsw, and β -secretase cleavable fragments thereof; and (iii) operatively linked to (i) and (ii), a regulatory sequence effective for expression of said nucleic acid molecules in said cell. The nucleic
25 acid encoding the β -secretase protein may be heterologous to the cell, and the nucleic acid encoding the β -secretase may be heterologous to the cell.

The invention is also directed to a method of screening for compounds that inhibit $A\beta$ production, comprising contacting a β -secretase polypeptide of the invention with (i)
a test compound and (ii) a β -secretase substrate, and selecting the test compound as
30 capable of inhibiting $A\beta$ production if the β -secretase polypeptide exhibits less β -secretase activity in the presence of the test compound than in the absence of the test compound. Such an assay may be cell-based, with one or both of the enzyme and the

substrate produced by the cell, such as the co-expression cell referred to above. Kits embodying such screening methods also form a part of the invention.

5 β -secretase inhibitor compounds selected according to the methods described above may be selected, for example, from a phage display selection system ("library"), such as are known in the art. Such libraries may be "biased" for the sequence peptide SEQ ID NO: 97 [P10-P4'D \rightarrow V]. Other inhibitors include, or may be derived from peptide inhibitors herein identified, such as inhibitors SEQ ID NO: 78, SEQ ID NO: 72, SEQ ID NO: 78 and SEQ ID NO: 81.

10 The β -secretase polypeptide may have a sequence selected from the group consisting of SEQ ID NO: 43 [46-501] and SEQ ID NO: 58 [46-452]. The β -secretase polypeptide and the substrate may be produced by a cell of the invention, and the β -secretase substrate may be selected from the group consisting of MBP-C125wt, MBP-C125sw, APP, APPsw, and β -secretase-cleavable fragments thereof, such as those fragments discussed herein.

15 The invention also provides a method of screening for compounds that inhibit A β production, comprising measuring binding of a purified β -secretase polypeptide of the invention with a β -secretase inhibitor compound in the presence of a test compound, and selecting the test compound as β -secretase active-site binding compound, if binding of the inhibitor in the presence of said test compound is less than binding of the inhibitor in the absence of said test compound. The inhibitor compound may be labelled with a
20 detectable marker, may be an inhibitor as discussed herein, and may have a K_i with respect to β -secretase of less than about 50 μ M.

The invention further provides a screening kit, comprising an isolated β -secretase protein of the present invention, a cleavable β -secretase substrate selected from the group
25 consisting of MBP-C125wt, MBP-C125sw, APPwt, APPsw, and β -secretase cleavable fragments thereof, and means for detecting cleavage of said substrate by β -secretase. The β -secretase protein may be present in a heterologous cell and the β -secretase-cleavable fragment may be as discussed herein.

30 In a yet further aspect, the invention provides the use of a compound effective to inhibit a β -secretase enzyme of the invention in the manufacture of a medicament for treating a patient afflicted with or having a predilection for Alzheimer's disease or other cerebrovascular amyloidosis wherein the compound has a K_i of less than

50 μ M in a MBP-C125sw assay.

According to this aspect, the enzymatic hydrolysis of APP to A β is blocked by administering to the patient a pharmaceutically effective dose of a compound effective to inhibit one or more of the various forms of the enzyme described herein. The therapeutic compound may be derived from a peptide selected from the group consisting of SEQ ID NO: 72, SEQ ID NO: 78, SEQ ID NO: 81 and SEQ ID NO: 97. Such derivation may be effected by the various phage selection systems described herein, in conjunction with the screening methods of the invention, or other such methods. Alternatively, or in addition, derivation may be achieved via rational chemistry approaches, including molecular modelling, known in the medicinal chemistry art. Also provided is an *in vitro* method of inhibiting enzymatic proteolysis of APP to A β in a tissue, comprising contacting said tissue with a compound effective to inhibit the enzymatic activity of a β -secretase protein of the present invention, wherein the compound has a K_i of less than 1-50 μ M in a MBP-C125sw assay, and a therapeutic drug composition for the treatment of Alzheimer's disease or other cerebrovascular amyloidosis characterised by deposition of A β peptide, wherein the active compound in said drug is selected for its ability to inhibit the enzymatic activity of a β -secretase protein of the present invention. The therapeutic drug may be derived from a peptide selected from the group consisting of SEQ ID NO: 72, SEQ ID NO: 78, SEQ ID NO: 81 and SEQ ID NO: 97, and may also include a pharmaceutically effective excipient.

According to yet another related aspect, the invention includes a method of diagnosing the presence of or a predilection for Alzheimer's disease in a patient, comprising detecting the expression level of a gene comprising a nucleic acid encoding a segment of a β -secretase enzyme lacking the signal sequence (amino acid residues 1-21 with respect to SEQ ID NO:2) and lacking the putative pro region (amino acid residues 22-45 with respect to SEQ ID NO:2) in a cell sample from said patient, and diagnosing the patient as having or having a predilection for Alzheimer's disease, if said expression level is significantly greater than a pre-determined control expression level. Detectable nucleic acids, and primers useful in such detection, are described in detail herein. The diagnostic method may be carried out in a whole cell assay and/or on a nucleic acid derived from a cell sample of said patient.

The invention will be described in more detail in the following detailed description of the invention read in conjunction with the accompanying drawings.

Brief Description of the Figures

5

FIG. 1A shows the sequence of a polynucleotide (SEQ ID NO: 1) which encodes human β -secretase translation product shown in FIG. 2A.

FIG. 1B shows the polynucleotide of FIG. 1A, including putative 5'- and 3'- untranslated regions (SEQ ID NO: 44).

FIG. 2A shows the amino acid sequence (SEQ ID NO: 2) of the predicted translation product of the open reading frame of the polynucleotide sequence shown in FIGS. 1A and 1B.

FIG. 2B shows the amino acid sequence of an active fragment of human β -secretase (SEQ ID NO: 43) [46-501].

FIG. 3A shows the translation product that encodes an active fragment of human β -secretase, 452stop, (amino acids 1-452 with reference to SEQ ID NO: 2; SEQ ID NO: 59) including a FLAG-epitope tag (underlined; SEQ ID NO: 45) at the C-terminus.

FIG. 3B shows the amino acid sequence of a fragment of human β -secretase (amino acids 46-452 (SEQ ID NO: 58) with reference to SEQ ID NO: 2; including a FLAG-epitope tag (underlined; SEQ ID NO: 45) at the C-terminus.

FIG. 4 shows an elution profile of recombinant β -secretase eluted from a gel filtration column.

FIG. 5 shows the full length amino acid sequence of β -secretase 1-501 (SEQ ID NO: 2), including the ORF which encodes it (SEQ ID NO: 1), with certain features indicated, such as "active-D" sites indicating the aspartic acid active catalytic sites, a transmembrane region commencing at position 453, as well as leader ("Signal") sequence (residues 1-21; SEQ ID NO: 46) and putative pro region (residues 22-45; SEQ ID NO: 47) and where the polynucleotide region corresponding the proenzyme region corresponding to amino acids 46-501 (SEQ ID NO: 43) (nt 135-1503) is shown as SEQ ID NO: 44 and contains an internal peptide region (SEQ ID NO: 56) and a transmembrane region (SEQ ID NO: 62).

FIGS. 6A and 6B show images of silver-stained SDS-PAGE gels on which purified β -secretase-containing fractions were run under reducing (6A) and non-reducing (6B) conditions.

FIG. 7 shows a silver-stained SDS-PAGE of β -secretase purified from heterologous 293T cells expressing the recombinant enzyme.

FIG. 8 shows a silver-stained SDS-PAGE of β -secretase purified from heterologous Cos A2 cells expressing the recombinant enzyme.

FIG. 9 shows a scheme in which primers derived from the polynucleotide (SEQ ID NO. 76 encoding N-terminus of purified naturally occurring β -secretase (SEQ ID NO. 77) were used to PCR-clone additional portions of the molecule, such as fragment SEQ ID NO. 79 encoding by nucleic acid SEQ ID NO. 98, as illustrated.

FIG. 10 shows an alignment of the amino acid sequence of human β -secretase ("Human Imapain.seq," 1-501, SEQ ID NO: 2) compared to ("pBS/mImpain 11#3 cons") consensus mouse sequence: SEQ ID NO: 65.

FIG. 11A shows the nucleotide sequence (SEQ ID NO: 80) of an insert used in preparing vector pCF.

FIG. 11B shows a linear schematic of pCEK.

FIG. 12 shows a schematic of pCEK.clone 27 used to transfect mammalian cells with β -secretase.

FIG. 13(A-E) shows the nucleotide sequence of pCEK clone 27 (SEQ ID NO: 48), with the OFR indicated by the amino acid sequence SEQ ID NO: 2.

FIG. 14A shows a nucleotide sequence inserted into parent vector pCDNA3 (SEQ ID NO: 10

FIG. 14B shows a plot of β -secretase activity in cell lysates from COS cells transfected with vectors derived from clones encoding β -secretase.

FIGS. 15A shows an image of an SDS PAGE gel loaded with triplicate samples of the lysates made from heterologous cells transfected with mutant APP (751 wt) and β -galactosidase as control (lanes d) and from cells transfected with mutant APP (751 wt) and β -secretase (lanes f) where lanes a, b, and c show lysates from untreated cells, cells transfected with β -galactosidase alone and cells transfected with β -secretase alone, respectively, and lane e indicates markers.

FIG. 15B shows an image of an SDS PAGE gel loaded with triplicate samples of the lysates made from heterologous cells transfected with mutant APP (Swedish mutation) and β -galactosidase as control (lanes c) and from cells transfected with mutant APP (Swedish mutation) and β -secretase (lanes e) where lanes a and b show lysates from cells transfected with β -galactosidase alone and cells transfected with β -secretase alone, and lane d indicates markers.

FIGS. 16A and 16B show Western blots of cell supernatants tested for presence or increase in soluble APP (sAPP).

FIGS. 17A and 17B show Western blots of α -cleaved APP substrate in co-expression cells.

FIG. 18 shows AB (x-40) production in 293T cells cotransfected with APP and β -secretase.

FIG. 19A shows a schematic of an APP substrate fragment, and its use in conjunction with antibodies SW192 and 8E-192 in the assay.

FIG. 19B shows the β -secretase cleavage sites in the wild-type APP sequence (SEQ ID NO: 103) and Swedish APP sequence (SEQ ID NO: 104).

FIG. 20 shows a schematic of a second APP substrate fragment derived from APP 638, and its use in conjunction with antibodies SW192 and 8E-192 in the assay.

FIG. 21 shows a schematic of pohCK751 vector.

Brief Description of the Sequences

10 This section briefly identifies the sequence identification numbers referred to herein. Number ranges shown in brackets here and throughout the specification are referenced to the amino acid sequence SEQ ID NO: 2, using conventional N \rightarrow C-terminus order.

SEQ ID NO: 1 is a nucleic acid sequence that encodes human β -secretase, including an active fragment, as exemplified herein.

15 SEQ ID NO: 2 is the predicted translation product of SEQ ID NO: 1 [1-501].

SEQ ID NOS: 3-21 are degenerate oligonucleotide primers described in Example 1 (Table 4), designed from regions of SEQ ID NO: 2.

SEQ ID NOS: 22-41 are additional oligonucleotide primers used in PCR cloning methods described herein, shown in Table 5.

20 SEQ ID NO: 42 is a polynucleotide sequence that encodes the active enzyme β -secretase shown as SEQ ID NO: 43.

SEQ ID NO: 43 is the sequence of an active enzyme portion of human β -secretase, the N-terminus of which corresponds to the N-terminus of the predominant form of the protein isolated from natural sources [46-501].

25 SEQ ID NO: 44 is a polynucleotide which encodes SEQ ID NO: 2, including 5' and 3' untranslated regions.

SEQ ID NO: 45 is the FLAG sequence used in conjunction with certain polynucleotides.

SEQ ID NO: 46 is the putative leader region of β -secretase [1-22].

30 SEQ ID NO: 47 is the putative pre-pro region of β -secretase [23-45].

SEQ ID NO: 48 is the sequence of the clone pCEK Cl.27 (FIG. 13A-E).

SEQ ID NO: 49 is a nucleotide sequence of a fragment of the gene which encodes human β -secretase.

SEQ ID NO: 50 is the predicted translation product of SEQ ID NO: 49.

SEQ ID NO: 51 is the predicted internal amino acid sequence of a portion of human β -secretase.

SEQ ID NOS: 52 and 53 are peptide substrates suitable for use in β -secretase assays used in the present invention.

SEQ ID NO: 54 is a peptide sequence cleavage site of APP (wild-type) recognized by a human β -

SEQ ID NO: 55 is amino acids 46-69 of SEQ ID NO: 2.

SEQ ID NO: 56 is an internal peptide just N-terminal to the transmembrane domain of β -secretase.

SEQ ID NO: 57 is β -secretase [1-419].

SEQ ID NO: 58 is β -secretase [46-452].

SEQ ID NO: 59 is β -secretase [1-452].

SEQ ID NO: 60 is β -secretase [1-420].

SEQ ID NO: 61 is EVM[hydroxyethylene]AEF.

SEQ ID NO: 62 is the amino acid sequence of the transmembrane domain of β -secretase shown in (FIG. 5).

SEQ ID NO: 63 is P26-P4' of APPwt.

SEQ ID NO: 64 is P26-P1' of APPwt.

SEQ ID NO: 65 is mouse β -secretase (FIG. 10, lower sequence).

SEQ ID NO: 66 is β -secretase [22-501].

SEQ ID NO: 67 is β -secretase [58-501].

SEQ ID NO: 68 is β -secretase [58-452].

SEQ ID NO: 69 is β -secretase [63-501].

SEQ ID NO: 70 is β -secretase [63-452].

SEQ ID NO: 71 is β -secretase [46-419].

SEQ ID NO: 72 is P10-P4'staD \rightarrow V.

SEQ ID NO: 73 is P4-P4'staD \rightarrow V.

SEQ ID NO: 74 is β -secretase [22-452].

SEQ ID NO: 75 is β -secretase [63-423].

SEQ ID NO: 76 is nucleic acid encoding the N-terminus of naturally occurring β -secretase.

SEQ ID NO: 77 is a peptide fragment at the N-terminus of naturally occurring β -secretase.

5 SEQ ID NO: 78 is a P3-P4'XD \rightarrow V (VMXVAEF, where X is hydroxyethylene or statine).

SEQ ID NO: 79 is a peptide fragment of naturally occurring β -secretase.

SEQ ID NO: 80 is a nucleotide insert in vector pCF used herein.

10 SEQ ID NO: 81 is P4-P4'XD \rightarrow V (EVMXVAEF, where X is hydroxyethylene or statine).

SEQ ID NO: 82 is APP fragment SEVKMDAEF (P5-P4'wt).

SEQ ID NO: 83 is APP fragment SEVNLDAEF (P5-P4'sw).

SEQ ID NO: 84 is APP fragment SEVKLDAEF.

SEQ ID NO: 85 is APP fragment SEVKFDAEF.

15 SEQ ID NO: 86 is APP fragment SEVNFDAEF.

SEQ ID NO: 87 is APP fragment SEVKMAAEF.

SEQ ID NO: 88 is APP fragment SEVNLAEEF.

SEQ ID NO: 89 is APP fragment SEVKLAAEF.

SEQ ID NO: 90 is APP fragment SEVKMLAEF.

20 SEQ ID NO: 91 is APP fragment SEVNLLAEF.

SEQ ID NO: 92 is APP fragment SEVKLLAEF.

SEQ ID NO: 93 is APP fragment SEVKFAAEF.

SEQ ID NO: 94 is APP fragment SEVNFAAEF.

SEQ ID NO: 95 is APP fragment SEVKFLAEF.

25 SEQ ID NO: 96 is APP fragment SEVNFLAEF.

SEQ ID NO: 97 is APP-derived fragment P10-P4'(D \rightarrow V): KTEEISEVNLVAEF.

SEQ ID NO: 98 is a nucleic acid fragment (FIG.9).

SEQ ID NO: 99 is the N terminal peptide sequence of β -secretase isolated from human brain, recombinant 293T cells and recombinant Cos A2 cells (Table 3).

SEQ ID NO: 100 is the N terminal peptide sequence of a form of β -secretase isolated from recombinant 293T cells.

SEQ ID NO: 101 is the N terminal peptide sequence of a form of β -secretase isolated from recombinant 293T cells.

SEQ ID NO: 102 is the N terminal peptide sequence of a form of β -secretase isolated from recombinant Cos A2 cells.

SEQ ID NO: 103 is the β -secretase cleavage sites in the wild-type APP sequence.

SEQ ID NO: 104 is the β -secretase cleavage sites in the Swedish APP sequence.

Detailed Description of the Invention

I. Definitions

Unless otherwise indicated, all terms used herein have the same meaning as they would to one skilled in the art of the present invention. Practitioners are particularly directed to Sambrook, *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (Second Edition), Cold Spring Harbor Press, Plainview, N.Y., and Ausubel, F.M., *et al.* (1998) *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, NY, for definitions, terms of art and standard methods known in the art of molecular biology, particularly as it relates to the cloning protocols described herein. It is understood that this invention is not limited to the particular methodology, protocols, and reagents described, as these may be varied to produce the same result.

The terms "polynucleotide" and "nucleic acid" are used interchangeably herein and refer to a polymeric molecule having a backbone that supports bases capable of hydrogen bonding to typical polynucleotides, where the polymer backbone presents the bases in a manner to permit such hydrogen bonding in a sequence specific fashion between the polymeric molecule and a typical polynucleotide (e.g., single-stranded DNA). Such bases are typically inosine, adenosine, guanosine, cytosine, uracil and thymidine. Polymeric molecules include double and single stranded RNA and DNA, and backbone modifications thereof, for example, methylphosphonate linkages.

The term "vector" refers to a polynucleotide having a nucleotide sequence that can assimilate new nucleic acids, and propagate those new sequences in an appropriate host. Vectors include, but are not limited to recombinant plasmids and viruses. The vector (e.g., plasmid or recombinant virus) comprising the nucleic acid of the invention can be in a carrier, for example, a plasmid complexed to protein, a plasmid complexed with lipid-based nucleic acid transduction systems, or other non-viral carrier systems.

The term "polypeptide" as used herein refers to a compound made up of a single chain of amino acid residues linked by peptide bonds. The term "protein" may be synonymous with the term "polypeptide" or may refer to a complex of two or more polypeptides.

The term "modified", when referring to a polypeptide of the invention, means a polypeptide which is modified either by natural processes, such as processing or other post-translational modifications, or by chemical modification techniques which are well known in the art. Among the numerous known modifications which may be present include, but are not limited to, acetylation, acylation, amidation, ADP-ribosylation, glycosylation, GPI anchor

formation, covalent attachment of a lipid or lipid derivative, methylation, myristylation, pegylation, prenylation, phosphorylation, ubiquitination, or any similar process.

The term "β-secretase" is defined in Section III, herein.

5 The term "biologically active" used in conjunction with the term β-secretase refers to possession of a β-secretase enzyme activity, such as the ability to cleave β-amyloid precursor protein (APP) to produce β-amyloid peptide (Aβ).

The term "fragment," when referring to β-secretase of the invention, means a polypeptide which has an amino acid sequence which is the same as part of but not all of the amino acid sequence of full-length β-secretase polypeptide. In the context of the present
10 invention, the full length β-secretase is generally identified as SEQ ID NO: 2, the ORF of the full-length nucleotide; however, according to a discovery of the invention, the naturally occurring active form is probably one or more N-terminal truncated versions, such as amino acids 46-501 (SEQ ID NO: 43), 22-501 (SEQ ID NO: 66), 58-501 (SEQ ID NO: 67) or 63-501 (SEQ ID NO: 69); other active forms are C-terminal truncated forms ending between about amino acids 450 and
15 452. The numbering system used throughout is based on the numbering of the sequence SEQ ID NO: 2.

An "active fragment" is a β-secretase fragment that retains at least one of the functions or activities of β-secretase, including but not limited to the β-secretase enzyme activity discussed above and/or ability to bind to the inhibitor substrate described herein as P10-P4'staD->V (SEQ ID NO: 72). Fragments contemplated include, but are not limited to, a β-secretase
20 fragment which retains the ability to cleave β-amyloid precursor protein to produce β-amyloid peptide. Such a fragment preferably includes at least 350, and more preferably at least 400, contiguous amino acids or conservative substitutions thereof of β-secretase, as described herein. More preferably, the fragment includes active aspartyl acid residues in the structural proximities identified and defined by the primary polypeptide structure shown as
25 SEQ ID NO: 2 and also denoted as "Active-D" sites herein.

A "conservative substitution" refers to the substitution of an amino acid in one class by an amino acid in the same class, where a class is defined by common physicochemical amino acid sidechain properties and high substitution frequencies in homologous proteins found in nature (as determined, e.g., by a standard Dayhoff frequency exchange matrix or
30 BLOSUM matrix). Six general classes of amino acid sidechains, categorized as described above, include: Class I (Cys); Class II (Ser, Thr, Pro, Ala, Gly); Class III (Asn, Asp, Gln, Glu); Class IV (His, Arg, Lys); Class V (Ile, Leu, Val, Met); and Class VI (Phe, Tyr, Trp).

For example, substitution of an Asp for another class III residue such as Asn, Gln, or Glu, is considered to be a conservative substitution.

"Optimal alignment" is defined as an alignment giving the highest percent identity score. Such alignment can be performed using a variety of commercially available sequence analysis programs, such as the local alignment program LALIGN using a ktp of 1, default parameters and the default PAM. A preferred alignment is the pairwise alignment using the CLUSTAL-W program in MacVector, operated with default parameters, including an open gap penalty of 10.0, an extended gap penalty of 0.1, and a BLOSUM30 similarity matrix.

"Percent sequence identity," with respect to two amino acid or polynucleotide sequences, refers to the percentage of residues that are identical in the two sequences when the sequences are optimally aligned. Thus, 80% amino acid sequence identity means that 80% of the amino acids in two or more optimally aligned polypeptide sequences are identical. If a gap needs to be inserted into a first sequence to optimally align it with a second sequence, the percent identity is calculated using only the residues that are paired with a corresponding amino acid residue (i.e., the calculation does not consider residues in the second sequences that are in the "gap" of the first sequence.

A first polypeptide region is said to "correspond" to a second polypeptide region when the regions are essentially co-extensive when the sequences containing the regions are aligned using a sequence alignment program, as above. Corresponding polypeptide regions typically contain a similar, if not identical, number of residues. It will be understood, however, that corresponding regions may contain insertions or deletions of residues with respect to one another, as well as some differences in their sequences.

A first polynucleotide region is said to "correspond" to a second polynucleotide region when the regions are essentially co-extensive when the sequences containing the regions are aligned using a sequence alignment program, as above. Corresponding polynucleotide regions typically contain a similar, if not identical, number of residues. It will be understood, however, that corresponding regions may contain insertions or deletions of bases with respect to one another, as well as some differences in their sequences.

The term "sequence identity" means nucleic acid or amino acid sequence identity in two or more aligned sequences, aligned as defined above.

"Sequence similarity" between two polypeptides is determined by comparing the amino acid sequence and its conserved amino acid substitutes of one polypeptide to the sequence of a second polypeptide. Thus, 80% protein sequence similarity means that 80%

the amino acid residues in two or more aligned protein sequences are conserved amino acid residues, *i.e.* are conservative substitutions.

"Hybridization" includes any process by which a strand of a nucleic acid joins with a complementary nucleic acid strand through base pairing. Thus, strictly speaking, the term
5 refers to the ability of the complement of the target sequence to bind to the test sequence, or vice-versa.

"Hybridization conditions" are based in part on the melting temperature (T_m) of the nucleic acid binding complex or probe and are typically classified by degree of "stringency" of the conditions under which hybridization is measured. The specific conditions that define
10 various degrees of stringency (*i.e.*, high, medium, low) depend on the nature of the polynucleotide to which hybridization is desired, particularly its percent GC content, and can be determined empirically according to methods known in the art. Functionally, maximum stringency conditions may be used to identify nucleic acid sequences having strict identity or near-strict identity with the hybridization probe; while high stringency conditions are used to
15 identify nucleic acid sequences having about 80% or more sequence identity with the probe.

The term "gene" as used herein means the segment of DNA involved in producing a polypeptide chain; it may include regions preceding and following the coding region, *e.g.* 5' untranslated (5' UTR) or "leader" sequences and 3' UTR or "trailer" sequences, as well as intervening sequences (introns) between individual coding segments (exons).

20 The term "isolated" means that the material is removed from its original environment (*e.g.*, the natural environment if it is naturally occurring). For example, a naturally occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such isolated polynucleotides may be part of a vector and/or such
25 polynucleotides or polypeptides may be part of a composition, such as a recombinantly produced cell (heterologous cell) expressing the polypeptide, and still be isolated in that such vector or composition is not part of its natural environment.

An "isolated polynucleotide having a sequence which encodes β -secretase" is a polynucleotide that contains the coding sequence of β -secretase, or an active fragment
30 thereof, (i) alone, (ii) in combination with additional coding sequences, such as fusion protein or signal peptide, in which the β -secretase coding sequence is the dominant coding sequence, (iii) in combination with non-coding sequences, such as introns and control elements, such as promoter and terminator elements or 5' and/or 3' untranslated regions,

effective for expression of the coding sequence in a suitable host, and/or (iv) in a vector or host environment in which the β -secretase coding sequence is a heterologous gene.

The terms "heterologous DNA," "heterologous RNA," "heterologous nucleic acid," "heterologous gene," and "heterologous polynucleotide" refer to nucleotides that are not endogenous to the cell or part of the genome in which they are present; generally such nucleotides have been added to the cell, by transfection, microinjection, electroporation, or the like. Such nucleotides generally include at least one coding sequence, but this coding sequence need not be expressed.

The term "heterologous cell" refers to a recombinantly produced cell that contains at least one heterologous DNA molecule.

A "recombinant protein" is a protein isolated, purified, or identified by virtue of expression in a heterologous cell, said cell having been transduced or transfected, either transiently or stably, with a recombinant expression vector engineered to drive expression of the protein in the host cell.

The term "expression" means that a protein is produced by a cell, usually as a result of transfection of the cell with a heterologous nucleic acid.

"Co-expression" is a process by which two or more proteins or RNA species of interest are expressed in a single cell. Co-expression of the two or more proteins is typically achieved by transfection of the cell with one or more recombinant expression vectors(s) that carry coding sequences for the proteins. In the context of the present invention, for example, a cell can be said to "co-express" two proteins, if one or both of the proteins is heterologous to the cell.

The term "expression vector" refers to vectors that have the ability to incorporate and express heterologous DNA fragments in a foreign cell. Many prokaryotic and eukaryotic expression vectors are commercially available. Selection of appropriate expression vectors within the knowledge of those having skill in the art.

The terms "purified" or "substantially purified" refer to molecules, either polynucleotides or polypeptides, that are removed from their natural environment, isolated or separated, and are at least 90% and more preferably at least 95-99% free from other components with which they are naturally associated. The foregoing notwithstanding, such descriptor does not preclude the presence in the same sample of splice- or other protein variants (glycosylation variants) in the same, otherwise homogeneous, sample.

A protein or polypeptide is generally considered to be "purified to apparent homogeneity" if a sample containing it shows a single protein band on a silver-stained polyacrylamide electrophoretic gel.

5 The term "crystallized protein" means a protein that has co-precipitated out of solution in pure crystals consisting only of the crystal, but possibly including other components that are tightly bound to the protein.

A "variant" polynucleotide sequence may encode a "variant" amino acid sequence that is altered by one or more amino acids from the reference polypeptide sequence. The variant polynucleotide sequence may encode a variant amino acid sequence, which contains
10 "conservative" substitutions, wherein the substituted amino acid has structural or chemical properties similar to the amino acid which it replaces. In addition, or alternatively, the variant polynucleotide sequence may encode a variant amino acid sequence, which contains "non-conservative" substitutions, wherein the substituted amino acid has dissimilar structural or chemical properties to the amino acid which it replaces. Variant polynucleotides may also
15 encode variant amino acid sequences, which contain amino acid insertions or deletions, or both. Furthermore, a variant polynucleotide may encode the same polypeptide as the reference polynucleotide sequence but, due to the degeneracy of the genetic code, has a polynucleotide sequence that is altered by one or more bases from the reference polynucleotide sequence.

20 An "allelic variant" is an alternate form of a polynucleotide sequence, which may have a substitution, deletion or addition of one or more nucleotides that does not substantially alter the function of the encoded polypeptide.

"Alternative splicing" is a process whereby multiple polypeptide isoforms are generated from a single gene, and involves the splicing together of nonconsecutive exons
25 during the processing of some, but not all, transcripts of the gene. Thus, a particular exon may be connected to any one of several alternative exons to form messenger RNAs. The alternatively-spliced mRNAs produce polypeptides ("splice variants") in which some parts are common while other parts are different.

"Splice variants" of β -secretase, when referred to in the context of an mRNA
30 transcript, are mRNAs produced by alternative splicing of coding regions, i.e., exons, from the β -secretase gene.

"Splice variants" of β -secretase, when referred to in the context of the protein itself, are β -secretase translation products that are encoded by alternatively-spliced β -secretase mRNA transcripts.

5 A "mutant" amino acid or polynucleotide sequence is a variant amino acid sequence, or a variant polynucleotide sequence, which encodes a variant amino acid sequence that has significantly altered biological activity or function from that of the naturally occurring protein.

A "substitution" results from the replacement of one or more nucleotides or amino acids by different nucleotides or amino acids, respectively.

10 The term "modulate" as used herein refers to the change in activity of the polypeptide of the invention. Modulation may relate to an increase or a decrease in biological activity, binding characteristics, or any other biological, functional, or immunological property of the molecule.

15 The terms "antagonist" and "inhibitor" are used interchangeably herein and refer to a molecule which, when bound to the polypeptide of the present invention, modulates the activity of enzyme by blocking, decreasing, or shortening the duration of the biological activity. An antagonist as used herein may also be referred to as a " β -secretase inhibitor" or " β -secretase blocker." Antagonists may themselves be polypeptides, nucleic acids, carbohydrates, lipids, small molecules (usually less than 1000 kD), or derivatives thereof, or
20 any other ligand which binds to and modulates the activity of the enzyme.

β -Secretase Compositions

The present invention provides isolated, active truncated forms of human β -secretase enzyme, which is an aspartyl (aspartic) protease or proteinase. As defined more fully in the sections that follow, β -secretase exhibits a proteolytic activity that is
25 involved in the generation of β -amyloid peptide from β -amyloid precursor protein (APP), such as is described in U.S. Patent 5,744,346, incorporated herein by reference. Alternatively, or in addition, the β -secretase is characterized by its ability to bind, with moderately high affinity, to an inhibitor substrate described herein as P10-P4' staD \rightarrow V (SEQ ID NO: 72). According to an important feature of the present invention,
30 truncated forms of human β -secretase have been isolated, and its naturally occurring form has been characterized, purified and sequenced.

According to another aspect of the invention, nucleotide sequences encoding the truncated enzyme have been identified. The truncated forms have similar protease activity to the naturally occurring or full-length recombinant enzyme. Using the information provided herein, practitioners can isolate DNA encoding various active forms of the protein from available sources and can express the protein recombinantly in a convenient expression system. Alternatively and in addition, practitioners can purify the enzyme from natural or recombinant sources and use it in purified form to further characterize its structure and function. According to a further feature of the invention, polynucleotides and proteins of the invention are particularly useful in a variety of screening assay formats, including cell-based screening for drugs that inhibit the enzyme. Examples of uses of such assays, as well as additional utilities for the compositions are provided in Section IV, below.

β -secretase is of particular interest due to its activity and involvement in generating fibril peptide components that are the major components of amyloid plaques in the central nervous system (CNS), such as are seen in Alzheimer's disease, Down's syndrome and other CNS disorders. Accordingly, a useful feature of the present invention includes an isolated truncated form of the enzyme that can be used, for example, to screen for inhibitory substances which are candidates for therapeutics for such disorders.

A. Isolation of Polynucleotides encoding Human β -secretase

Polynucleotides encoding human β -secretase were obtained by PCR cloning and hybridization techniques as detailed in Examples 1-3 and described below. FIG. 1A shows the sequence of a polynucleotide (SEQ ID NO: 1) which encodes a form of human β -secretase (SEQ ID NO: 2 [1-501]). Polynucleotides encoding human β -secretase are conveniently isolated from any of a number of human tissues, preferably tissues of neuronal origin, including but not limited to neuronal cell lines such as the commercially available human neuroblastoma cell line IMR-32 available from the American Type Culture Collection (Manassas, VA; ATTC CCL 127) and human fetal brain, such as a human fetal brain cDNA library available from OriGene Technologies, Inc. (Rockville, MD).

Briefly, human β -secretase coding regions were isolated by methods well known in the art, using hybridization probes derived from the coding sequence provided as SEQ ID NO: 1. Such probes can be designed and made by methods well known in the art. Exemplary probes, including degenerate probes, are described in Example 1. Alternatively, a

cDNA library is screened by PCR, using, for example, the primers and conditions described in Example 2 herein. Such methods are discussed in more detail in Part B, below.

cDNA libraries were also screened using a 3'-RACE (Rapid Amplification of cDNA Ends) protocol according to methods well known in the art (White, B.A., ed., PCR Cloning Protocols; Humana Press, Totowa, NJ, 1997; shown schematically in FIG. 9). Here primers derived from the 5' portion of SEQ ID NO: 1 are added to partial cDNA substrate clone found by screening a fetal brain cDNA library as described above. A representative 3'RACI reaction used in determining the longer sequence is detailed in Example 3 and is described in more detail in Part B, below.

- 10 Human β -secretase, as well as additional members of the neuronal aspartyl protease family described herein may be identified by the use of random degenerate primers designed in accordance with any portion of the polypeptide sequence shown as SEQ ID NO: 2. For example, in experiments carried out in support of the present invention, and detailed in Example 1 herein, eight degenerate primer pools, each 8-fold degenerate, were designed
15 based on a unique 22 amino acid peptide region selected from SEQ ID: 2. Such techniques can be used to identify further similar sequences from other species and/or representing other members of this protease family.

Preparation of polynucleotides

- 20 The polynucleotides described herein may be obtained by screening cDNA libraries using oligonucleotide probes, which can hybridize to and/or PCR-amplify polynucleotides that encode human β -secretase, as disclosed above. cDNA libraries prepared from a variety of tissues are commercially available, and procedures for screening and isolating cDNA clones are well known to those of skill in the art. Genomic libraries can likewise be screened
25 to obtain genomic sequences including regulatory regions and introns. Such techniques are described in, for example, Sambrook *et al.* (1989) Molecular Cloning: A Laboratory Manual (2nd Edition), Cold Spring Harbor Press, Plainview, N.Y. and Ausubel, FM *et al.* (1998) Current Protocols in Molecular Biology, John Wiley & Sons, New York, N.Y.

- The polynucleotides may be extended to obtain upstream and downstream sequences
30 such as promoters, regulatory elements, and 5' and 3' untranslated regions (UTRs). Extension of the available transcript sequence may be performed by numerous methods known to those of skill in the art, such as PCR or primer extension (Sambrook *et al.*, *supra*), or by the RAC

method using, for example, the MARATHON RACE kit (Cat. # K1802-1; Clontech, Palo Alto, CA).

Alternatively, the technique of "restriction-site" PCR (Gobinda *et al.* (1993) PCR Methods Applic. 2:318-22), which uses universal primers to retrieve flanking sequence adjacent a known locus, may be employed to generate additional coding regions. First, genomic DNA is amplified in the presence of primer to a linker sequence and a primer specific to the known region. The amplified sequences are subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

Inverse PCR can be used to amplify or extend sequences using divergent primers based on a known region (Triglia T *et al.* (1988) Nucleic Acids Res 16:8186). The primers may be designed using OLIGO(R) 4.06 Primer Analysis Software (1992; National Biosciences Inc, Plymouth, Minn.), or another appropriate program, to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68-72°C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

Capture PCR (Lagerstrom M *et al.* (1991) PCR Methods Applic 1:111-19) is a method for PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA. Capture PCR also requires multiple restriction enzyme digestions and ligations to place an engineered double-stranded sequence into a flanking part of the DNA molecule before PCR.

Another method which may be used to retrieve flanking sequences is that of Parker, JD *et al.* (1991; Nucleic Acids Res 19:3055-60). Additionally, one can use PCR, nested primers and PromoterFinder(TM) libraries to "walk in" genomic DNA (Clontech, Palo Alto, CA). This process avoids the need to screen libraries and is useful in finding intron/exon junctions. Preferred libraries for screening for full length cDNAs are ones that have been size-selected to include larger cDNAs. Also, random primed libraries are preferred in that they will contain more sequences which contain the 5' and upstream regions of genes. A randomly primed library may be particularly useful if an oligo d(T) library does not yield a full-length cDNA. Genomic libraries are useful for extension into the 5' nontranslated regulatory region.

The polynucleotides and oligonucleotides of the invention can also be prepared by solid-phase methods, according to known synthetic methods. Typically, fragments of up to about 100 bases are individually synthesized, then joined to form continuous sequences up to several hundred bases.

B. Isolation of β -Secretase

The amino acid sequence for a full-length human β -secretase translation product is shown as SEQ ID NO: 2 in FIG. 2A. According to the discovery of the present invention, this sequence represents a "pre pro" form of the enzyme that was deduced from the nucleotide sequence information described in the previous section in conjunction with the methods described below. Comparison of this sequence with sequences determined from the biologically active form of the enzyme purified from natural sources, as described in Part 4, below, indicate that it is likely that an active and predominant form of the enzyme is represented by sequence shown in FIG. 2B (SEQ ID NO: 43), in which the first 45 amino acids of the open-reading frame deduced sequence have been removed. This suggests that the enzyme may be post-translationally modified by proteolytic activity, which may be autocatalytic in nature. Further analysis, illustrated by the schematics shown in FIG. 5 herein, indicates that the enzyme contains a hydrophobic, putative transmembrane region near its C-terminus. As described below, a further discovery of the present invention is that the enzyme can be truncated prior to this transmembrane region and still retain β -secretase activity.

1. Purification of β -secretase from Natural and Recombinant Sources

β -secretase has now been purified from natural and recombinant sources. U.S. Patent 5,744,346, incorporated herein by reference, describes isolation of β -secretase in a single peak having an apparent molecular weight of 260-300,000 (Daltons) by gel exclusion chromatography. The native enzyme can be purified to apparent homogeneity by affinity column chromatography. The methods revealed herein have been used on preparations from brain tissue as well as on preparations from 293T and recombinant cells; accordingly, these methods are believed to be generally applicable over a variety of tissue sources. The practitioner will realize that certain of the preparation steps, particularly the initial steps, may require modification to accommodate a particular tissue source and will adapt such procedures according to methods known in the art. Methods for purifying β -secretase from

human brain as well as from cells are detailed in Example 5. Briefly, cell membranes or brain tissue are homogenized, fractionated, and subjected to various types of column chromatographic matrices, including wheat germ agglutinin-agarose (WGA), anion exchange chromatography and size exclusion. Activity of fractions can be measured using any appropriate assay for β -secretase activity, such as the MBP-C125 cleavage assay detailed in Example 4. Fractions containing β -secretase activity elute from this column in a peak elution volume corresponding to a size of about 260-300 kilodaltons.

The foregoing purification scheme, which yields approximately 1,500-fold purification, is similar to that described in detail in U.S. Patent 5,744,346, incorporated herein by reference. Further purification can be achieved by applying the cation exchange flow-through material to an affinity column that employs as its affinity matrix a specific inhibitor of β -secretase, termed "P10—P4'staD->V" (NH_2 -KTEEISEVN[sta]VAEF-CO₂H; SEQ ID NO.: 72). This inhibitor, and methods for making a Sepharose affinity column which incorporates it, are described in Example 7. After washing the column, β -secretase and a limited number of contaminating proteins were eluted with pH 9.5 borate buffer. The eluate was then fractionated by anion exchange HPLC, using a Mini-Q column. Fractions containing the activity peak were pooled to give the final β -secretase preparation. Results of an exemplary run using this purification scheme are summarized in Table 1. FIG. 6A shows a picture of a silver-stained SDS PAGE gel run under reducing conditions, in which β -secretase runs as a 70 kilodalton band. The same fractions run under non-reducing conditions (FIG. 6B) provide evidence for disulfide cross-linked oligomers. When the anion exchange pool fractions 18-21 (see FIG. 6B) were treated with dithiothreitol (DTT) and re-chromatographed on a Mini Q column, then subjected to SDS-PAGE under non-reducing conditions, a single band running at about 70 kilodaltons was observed. Surprisingly, the purity of this preparation is at least about 200 fold higher than the previously purified material, described in U.S. Patent 5,744,346. By way of comparison, the most pure fraction described therein exhibited a specific activity of about 253 nM/h/ μ g protein, taking into consideration the MW of substrate MBP-C26sw (45 kilodaltons). The method therefore provides a preparation that is at least about 1000-fold higher purity (affinity eluate) and as high as about 6000-fold higher purity than that preparation, which represented at least 5 to 100-fold higher purity than the enzyme present in a solubilized but unenriched membrane fraction from human 293 cells.

Table 1

Preparation of β -secretase from Human Brain

	Total Activity ^a nM/h	Specific Activity ^b nM/h/ μ g prot.	% Yield	Purification (fold)
Brain Extract	19,311,150	4.7	100	
WGA Eluate	21,189,600	81.4	110	
Affinity Eluate	11,175,000	257,500	53	54.8
Anion Exchange Pool	3,267,685	1,485,312	17	316.3

^aActivity in MBP-C125sw assay^bSpecific Activity = $\frac{(\text{Product conc. nM})(\text{Dilution factor})}{(\text{Enzyme sol. vol})(\text{Incub. time h})(\text{Enzyme conc. } \mu\text{g/vol})}$

Example 5 also describes purification schemes used for purifying recombinant materials from heterologous cells transfected with the β -secretase coding sequence. Results from these purifications are illustrated in FIGS. 7 and 8. Further experiments carried out in support of the present invention, showed that the recombinant material has an apparent molecular weight in the range from 260,000 to 300,000 Daltons when measured by gel exclusion chromatography. FIG. 4 shows an activity profile of this preparation run on a gel exclusion chromatography column, such as a Superdex 200 (26/60) column, according to the methods described in U.S. Patent 5,744,346, incorporated herein by reference.

1. Sequencing of β -secretase Protein

A schematic overview summarizing methods and results for determining the cDNA sequence encoding the N-terminal peptide sequence determined from purified β -secretase is shown in FIG. 9. N-terminal sequencing of purified β -secretase protein isolated from natural sources yielded a 21-residue peptide sequence, SEQ ID NO. 77, as described above. This peptide sequence, and its reverse translated fully degenerate nucleotide sequence, SEQ ID NO. 76, is shown in the top portion of FIG. 4. Two partially degenerate primer sets used for RT-PCR amplification of a cDNA fragment encoding this peptide are also summarized in FIG. 4. Primer set 1 consisted of DNA nucleotide primers #3427-3434 (SEQ ID NOS: 22-29), shown in Table 3 (Example 3). Matrix RT-PCR using combinations of primers from this set with cDNA reverse transcribed from primary human neuronal cultures as template yielded the predicted 54 bp cDNA product with primers #3428-3433 (SEQ ID NOS: 23-28), also described in Table 3.

In further experiments carried out in support of the present invention, it was found that oligonucleotides from primer sets 1 and 2 could also be used to amplify cDNA fragments of the predicted size from mouse brain mRNA. DNA sequence demonstrated that such primers could also be used to clone the murine homolog(s) and other species homologs of

human β -secretase and/or additional members of the aspartyl protease family described herein by standard RACE-PCR technology. The sequence of a murine homolog is presented in FIG. 10 (lower sequence; "pBS/MulmPain H#3 cons"); SEQ ID NO. 65. The murine polypeptide sequence is about 95% identical to the human polypeptide sequence.

2. 5' and 3' RACE-PCR for Additional Sequence, Cloning, and mRNA Analysis

The unambiguous internal nucleotide sequence from the amplified fragment provided information which facilitated the design of internal primers matching the upper (coding) strand for 3' RACE, and lower (non-coding) strand for 5' RACE (Frohman, M. A., M. K. Dush and G. R. Martin (1988). "Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene specific oligo-nucleotide primer." Proc. Natl. Acad. Sci. U. S. A. 85 (23): 8998-9002.) The DNA primers used for this experiment (#3459 & #3460) (SEQ ID NOS: 38 & 39) are illustrated schematically in FIG. 9, and the exact sequence of these primers is presented in Table 4 of Example 3.

Primers #3459 and #3476 (Table 5, SEQ ID NOS: 38 & 41) were used for initial 3' RACE amplification of downstream sequences from the IMR-32 cDNA library in the vector pLPCXlox. The library had previously been sub-divided into 100 pools of 5,000 clones per pool, and plasmid DNA was isolated from each pool. A survey of the 100 pools with the primers described in Part 2, above, identified individual pools containing β -secretase clones from the library. Such clones can be used for RACE-PCR analysis.

An approximately 1.8 Kb PCR fragment was observed by agarose gel fractionation of the reaction products. The PCR product was purified from the gel and subjected to DNA sequence analysis using primer #3459 (Table 5, SEQ ID NO: 38). The resulting clone sequence, designated 23A, was determined. Six of the first seven deduced amino-acids from one of the reading frames of 23A were an exact match with the last 7 amino-acids of the N-terminal sequence (SEQ ID NO. 77) determined from the purified protein isolated from natural sources in other experiments carried out in support of this invention. This observation provided internal validation of the sequences, and defined the proper reading frame downstream. Furthermore, this DNA sequence facilitated design of additional primers for extending the sequence further downstream, verifying the sequence by sequencing the opposite strand in the upstream direction, and further facilitated isolating the cDNA clone.

A DNA sequence of human β -secretase is illustrated as SEQ ID NO: 42 corresponding to SEQ ID NO: 1 including 5'- and 3'-untranslated regions. This sequence was determined from a partial cDNA clone (9C7c.35) isolated from a commercially available

human fetal brain cDNA library purchased from OriGene™, the 3' RACE product 23A, and additional clones – a total of 12 independent cDNA clones were used to determine the composite sequence. The composite sequence was assembled by sequencing overlapping stretches of DNA from both strands of the clone or PCR fragment. The predicted full length translation product is shown as SEQ ID NO: 2 in Fig. 1B.

4. Tissue Distribution of β -secretase and Related Transcripts

Oligonucleotide primer #3460 (SEQ ID NO. 39, Table 5) was employed as an end-labeled probe on Northern blots to determine the size of the transcript encoding β -secretase and to examine its expression in IMR-32 cells. Additional primers were used to isolate the mouse cDNA and to characterize mouse tissues, using Marathon RACE ready cDNA preparations (Clontech, Palo Alto, CA). TABLE 2 summarizes the results of experiments in which various human and murine tissues were tested for the presence of β -secretase-encoding transcripts by PCR or Northern blotting.

For example, the oligo-nucleotide probe 3460 (SEQ ID NO: 39) hybridized to a 2 Kb transcript in IMR-32 cells, indicating that the mRNA encoding the β -secretase enzyme is 2 Kb in size in this tissue. Northern blot analysis of total RNA isolated from the human T-cell line Jurkat, and human myelomonocyte line Thp1 with the 3460 oligo-nucleotide probe 3460 also revealed the presence of a 2 kb transcript in these cells.

The oligonucleotide probe #3460 (SEQ ID NO: 39) also hybridizes to a ~2 kb transcript in Northern blots containing RNA from all human organs examined to date, from both adult and fetal tissue. The organs surveyed include heart, brain, liver, pancreas, placenta, lung, muscle, uterus, bladder, kidney, spleen, skin, and small intestine. In addition, certain tissues, e. g. pancreas, liver, brain, muscle, uterus, bladder, kidney, spleen and lung, show expression of larger transcripts of ~4.5 kb, 5 kb, and 6.5 kb which hybridize with oligonucleotide probe #3460 (SEQ ID NO: 39).

In further experiments carried out in support of the present invention, Northern blot results were obtained with oligonucleotide probe #3460 (SEQ ID NO: 39) by employing a riboprobe derived from SEQ ID NO: 1, encompassing nucleotides #155-1014. This clone provides an 860 bp riboprobe, encompassing the catalytic domain-encoding portion of β -secretase, for high stringency hybridization. This probe hybridized with high specificity to the exact match mRNA expressed in the samples being examined. Northern blots of mRNA isolated from IMR-32 and 1°HNC probed

with this riboprobe revealed the presence of the 2 kb transcript previously detected with oligonucleotide #3460, as well as a novel, higher MW transcript of ~5 kb. Hybridization of RNA from adult and fetal human tissues with this 860 nt riboprobe also confirmed the result obtained with the oligonucleotide probe #3460 (SEQ ID NO: 39). The mRNA
5 encoding β -secretase is expressed in all tissues examined, predominantly as an ~5 kb transcript. In adult, its expression appeared lowest in brain, placenta, and lung, intermediate in uterus, and bladder, and highest in heart, liver, pancreas, muscle, kidney, spleen, and lung. In fetal tissue, the message is expressed uniformly in all tissues examined.

Table 2 Tissue distribution of human and murine β -secretase transcripts

Tissue/Organ	Size Messages Found (Kb):		Clontech Human Brain region	
	Human	Mouse	Tissue/Organ	Human
Heart	2 ^a	3.5, 3.8, 5 & 7	Cerebellum	2Kb, 4Kb, 6Kb
Brain	2, 3, 4, and 7	3.5, 3.8, 5 & 7	Cerebral Cx	2Kb, 4Kb, 6Kb
Liver	2, 3, 4, and 7	3.5, 3.8, 5 & 7	Medulla	2Kb, 4Kb, 6Kb
Pancreas	2, 3, 4, and 7	nd ^d	Spinal Cord	2Kb, 4Kb, 6Kb
Placenta	2 ^a , 4 and 7 ^b	nd	Occipital Pole	2Kb, 4Kb, 6Kb
Lung	2 ^a , 4 and 7 ^b	3.5, 3.8, 5 & 7	Frontal Lobe	2Kb, 4Kb, 6Kb
Muscle	2 ^a and 7 ^b	3.5, 3.8, 5 & 7	Amygdala	2Kb, 4Kb, 6Kb
Uterus	2 ^a , 4, and 7	nd	Caudate N.	2Kb, 4Kb, 6Kb
Bladder	2 ^a , 3, 4, and 7	nd	Corpus Callosum	2Kb, 4Kb, 6Kb
Kidney	2 ^a , 3, 4, and 7	3.5, 3.8, 5 & 7	Hippocampus	2Kb, 4Kb, 6Kb
Spleen	2 ^a , 3, 4, and 7	nd		
Testis	nd	4.5Kb, 2Kb	Substantia Nigra	2Kb, 4Kb, 6Kb
Stomach	nd	5 ^a	Thalamus	2Kb, 4Kb, 6Kb
Sm. Intestine	nd	3.5, 3.8, 5 & 7		
f Brain ^c	2 ^a , 3, 4, and 7	nd		
f Liver	2 ^a , 3, 4, and 7	nd		
f Lung	2 ^a , 3, 4, and 7	nd		
f Muscle	2 ^a , 3, 4, and 7	nd		
f Heart	2 ^a , 3, 4, and 7	nd		
f Kidney	2 ^a , 3, 4, and 7	nd		
f Skin	2 ^a , 3, 4, and 7	nd		
f Sm. Intestine	2 ^a , 3, 4, and 7	nd		
Cell Line	Human	Mouse		
IMR32	2 ^a , 5 & 7			
U937	2 ^a			
THP1	2 ^a			
Jurkat	2 ^a			
HL60	none			
A293	5 & 7			
NALM6	5 & 7			
A549	5 & 7			
Hela	2, 4, 5, & 7			
PC12		2 & 5		
J774		5Kb, 2Kb		
P388D1 ccl46		5Kb (very little), 2Kb		
P19		5Kb, 2Kb		
RBL		5Kb, 2Kb		
EL4		5Kb, 2Kb		

^aby oligo 3460 probe only

^cf = fetal

^bfaint

^dnd=not determined

5. Active Forms of β -secretase

a. N-terminus

5 The full-length open reading frame (ORF) of human β -secretase is described above, and its sequence is shown in FIG. 2A as SEQ ID NO: 2. However, as mentioned above, a further discovery of the present invention indicates that the predominant form of the active, naturally occurring molecule is truncated at the N-terminus by about 45 amino acids. That is, the protein purified from natural sources was N-terminal sequenced according to methods known in the art (Argo Bioanalytica, Morris Plains, NJ). The N-terminus yielded the following sequence: ETDEEPEEPGRRGSFVEMVDNLRG... (SEQ ID NO: 55). This corresponds to amino acids 46-69 of the ORF-derived putative sequence. Based on this observation and others described below, the N-terminus of an active, naturally occurring, predominant human brain form of the enzyme is amino acid 46, with respect to SEQ ID NO: 2. Further processing of the purified protein provided the sequence of an internal peptide: IGFAVSACHVHDEFR (SEQ ID NO: 56), which is amino terminal to the putative transmembrane domain, as defined by the ORF. These peptides were used to validate and provide reading frame information for the isolated clones described elsewhere in this application.

20 In additional studies carried out in support of the present invention, N-terminal sequencing of β -secretase isolated from additional cell types revealed that the N-terminus may be amino acid numbers 46, 22, 58, or 63 with respect to the ORF sequence shown in FIG. 2A, depending on the tissue from which the protein is isolated, with the form having as its N-terminus amino acid 46 predominating in the tissues tested. That is, in experiments carried out in support of the present invention, the full-length β -secretase construct (i.e., encoding SEQ ID NO: 2) was transfected into 293T cells and COS A2 cells, using the Fugene technique described in Example 6. β -secretase was isolated from the cells by preparing a crude particulate fraction from the cell pellet, as described in Example 5, followed by extraction with buffer containing 0.2% Triton X-100. The Triton extract was diluted with pH 5.0 buffer and passed through a SP Sepharose column, essentially according to the methods described in Example 5A. This step removed the majority of contaminating proteins. After adjusting the pH to 4.5, β -secretase was further purified and concentrated on P10-P4'staD \rightarrow V Sepharose, as described in Examples 5 and 7. Fractions were analyzed for N-

terminal sequence, according to standard methods known in the art. Results are summarized in Table 3, below.

The primary N-terminal sequence of the 293T cell-derived protein was the same as that obtained from brain. In addition, minor amounts of protein starting just after the signal sequence (at Thr-22) and at the start of the aspartyl protease homology domain (Met-63) were also observed. An additional major form found in Cos A2 cells resulted from a Gly-58 cleavage.

Table 3
N-terminal Sequences and Amounts of β -secretase Forms in Various Cell Types

Source	Est. Amount (pmoles)	N-terminus (Ref: SEQ ID NO: 2)	Sequence
Human brain	1-2	46	ETDEEPEEPGR... (SEQ ID NO: 1)
Recombinant, 293T	~35	46	ETDEEPEEPGR... (SEQ ID NO: 1)
	~7	22	TQHFIRL(P)LR... (SEQ ID NO: 1)
	~5	63	MVDNLRGKS... (SEQ ID NO: 1)
Recombinant, CosA2	~4	46	ETDEEPEEPGR... (SEQ ID NO: 1)
	~3	58	GSFVEMVDNL... (SEQ ID NO: 1)

b. C-terminus

Further experiments carried out in support of the present invention revealed that the C-terminus of the full-length amino acid sequence presented as SEQ ID NO: 2 can also be truncated, while still retaining β -secretase activity of the molecule. More specifically, as described in more detail in Part D below, C-terminal truncated forms of the enzyme ending just before the putative transmembrane region, i.e. at or about 10 amino acids C terminal to amino acid 452 with respect to SEQ ID NO: 2, exhibit β -secretase activity, as evidenced by an ability to cleave APP at the appropriate cleavage site and/or ability to bind SEQ ID NO: 72.

Thus, using the reference amino acid positions provided by SEQ ID NO: 2, one form of β -secretase extends from position 46 to position 501 (β -secretase 46-501; SEQ ID NO: 43). Another form extends from position 46 to any position including and beyond position 452, (β -secretase 4-452+), with a preferred form being β -secretase 46-452 (SEQ ID NO: 58). Other active forms of the β -secretase protein in accordance with the invention begin at amino acid 58 or 63 and may extend to any point including and beyond the cysteine at position 420, and more preferably, including and beyond position

452, while still retaining enzymatic activity (i.e., β -secretase 22-452+, β -secretase 58-452+, β -secretase 63-452+). As described in Part D, below, those forms which are truncated at a C-terminal position at or before about position 452, or even several amino acids thereafter, are particularly useful in crystallization studies, since they lack all or a significant portion of the transmembrane region, which may interfere with protein crystallization. The recombinant protein extending from position 1 to 452 has been affinity purified using the procedures described herein.

C. Crystallization of β -secretase

According to a further aspect, the present invention also includes truncated purified β -secretase in crystallized form, in the absence or presence of binding substrates, such as peptide, modified peptide, or small molecule inhibitors. This section describes methods and utilities of such compositions.

1. Crystallization of the Protein

Truncated β -secretase purified as described above can be used as starting material to determine a crystallographic structure and coordinates for the enzyme. Such structural determinations are particularly useful in defining the conformation and size of the substrate binding site. This information can be used in the design and modelling of substrate inhibitors of the enzyme. As discussed herein, such inhibitors are candidate molecules for therapeutics for treatment of Alzheimer's disease and other amyloid diseases characterized by A β peptide amyloid deposits.

The crystallographic structure of β -secretase is determined by first crystallizing the purified protein. Methods for crystallizing proteins, and particularly proteases, are now well known in the art. The practitioner is referred to Principles of Protein X-ray Crystallography (J. Drenth, Springer Verlag, NY, 1999) for general principles of crystallography. Additionally, kits for generating protein crystals are generally available from commercial providers, such as Hampton Research (Laguna Niguel, CA). Additional guidance can be obtained from numerous research articles that have been written in the area of crystallography of protease inhibitors, especially with respect to HIV-1 and HIV-2 proteases, which are aspartic acid proteases.

Although any of the various truncated forms of β -secretase described herein can be used for crystallization studies, particularly preferred forms lack the first 45 amino acids of the full length sequence shown as SEQ ID NO: 2, since this appears to be the predominant form

(which occurs naturally in human brain. It is thought that some form of post-translational modification, possibly autocatalysis, serves to remove the first 45 amino acids in fairly rapid order, since, to date, virtually no naturally occurring enzyme has been isolated with all of the first 45 amino acids intact. In addition, it is considered preferable to remove the putative transmembrane region from the molecule prior to crystallization, since this region is not necessary for catalysis and potentially could render the molecule more difficult to crystallize.

Thus, a good candidate for crystallization is β -secretase 46-452 (SEQ ID NO: 58), since this is a form of the enzyme that (a) provides the predominant naturally occurring N-terminus, and (b) lacks the "sticky" transmembrane region, while (c) retaining β -secretase activity. Alternatively, forms of the enzyme having extensions that extend part of the way (approximately 10-15 amino acids) into the transmembrane domain may also be used. In general, for determining X-ray crystallographic coordinates of the ligand binding site, any form of the enzyme can be used that either (i) exhibits β -secretase activity, and/or (ii) binds to a known inhibitor, such as the inhibitor ligand P10--P4'staD->V, with a binding affinity that is at least 1/100 the binding affinity of β -secretase [46-501](SEQ ID NO: 43) to P10--P4'staD->V (SEQ ID NO: 72). Therefore, a number of additional truncated forms of the enzyme can be used in these studies. Suitability of any particular form can be assessed by contacting it with the P10--P4'staD->V affinity matrix described above. Truncated forms of the enzyme that bind to the matrix are suitable for such further analysis. Thus, in addition to 46-452, discussed above, experiments in support of the present invention have revealed that a truncated form ending in residue 419, most likely 46-419 (SEQ ID NO: 71), also binds to the affinity matrix and is therefore an alternate candidate protein composition for X-ray crystallographic analysis of β -secretase. More generally, any form of the enzyme that ends before the transmembrane domain, particularly those ending between about residue 419 and 452 are suitable in this regard.

At the N-terminus, as described above, generally the first 45 amino acids will be removed during cellular processing. Other suitable naturally occurring or expressed forms include, for example, one commencing at residue 58 and one commencing at residue 63. However, analysis of the entire enzyme, starting at residue 1, can also provide information about the enzyme. Other forms, such as 1-420 (SEQ ID NO: 60) to 1-452 (SEQ ID NO: 59), including intermediate forms, for example 1-440, can be useful in this regard. In general, it will also be useful to obtain structure on any subdomain of the active enzyme.

(

Methods for purifying the protein, including active forms, are described above. In addition, since the protein is apparently glycosylated in its naturally occurring (and mammalian-expressed recombinant) forms, it may be desirable to express the protein and purify it from bacterial sources, which do not glycosylate mammalian proteins, or express it in sources, such as insect cells, that provide uniform glycosylation patterns, in order to obtain a homogeneous composition. Appropriate vectors and codon optimization procedures for accomplishing this are known in the art.

Following expression and purification, the protein is adjusted to a concentration of about 1-20 mg/ml. In accordance with methods that have worked for other crystallized proteins, the buffer and salt concentrations present in the initial protein solution are reduced to as low a level as possible. This can be accomplished by dialyzing the sample against the starting buffer, using microdialysis techniques known in the art. Buffers and crystallization conditions will vary from protein to protein, and possibly from fragment to fragment of the active β -secretase molecule, but can be determined empirically using, for example, matrix methods for determining optimal crystallization conditions. (Drentz, J., *supra*; Ducruix, A., *et al.*, eds. *Crystallization of Nucleic Acids and Proteins: A Practical Approach*, Oxford University Press, New York, 1992.)

Following dialysis, conditions are optimized for crystallization of the protein. Generally, methods for optimization may include making a "grid" of 1 μ l drops of the protein solution, mixed with 1 μ l well solution, which is a buffer of varying pH and ionic strength. These drops are placed in individual sealed wells, typically in a "hanging drop" configuration, for example in commercially available containers (Hampton Research, Laguna Niguel, CA). Precipitation/crystallization typically occurs between 2 days and 2 weeks. Wells are checked for evidence of precipitation or crystallization, and conditions are optimized to form crystals. Optimized crystals are not judged by size or morphology, but rather by the diffraction quality of crystals, which should provide better than 3 Å resolution. Typical precipitating agents include ammonium sulfate (NH_4SO_4), polyethylene glycol (PEG) and methyl pentane diol (MPD). All chemicals used should be the highest grade possible (e.g., ACS) and may also be re-purified by standard methods known in the art, prior to use.

Exemplary buffers and precipitants forming an empirical grid for determining crystallization conditions are commercially available. For example, the "Crystal Screen" kit (Hampton Research) provides a sparse matrix method of trial conditions that is biased and selected from known crystallization conditions for macromolecules. This provides a "grid"

for quickly testing wide ranges of pH, salts, and precipitants using a very small sample (50 to 100 microliters) of macromolecule. In such studies, 1 μ l of buffer/precipitant(s) solution is added to an equal volume of dialyzed protein solution, and the mixtures are allowed to sit for at least two days to two weeks, with careful monitoring of crystallization. Chemicals can be obtained from common commercial suppliers; however, it is preferable to use purity grades suitable for crystallization studies, such as are supplied by Hampton Research (Laguna Niguel, CA). Common buffers include Citrate, Tris, CHES, Acetate, ADA and the like (to provide a range of pH optima), typically at a concentration of about 100 mM. Typical precipitants include $(\text{NH}_4)_2\text{SO}_4$, MgSO_4 , NaCl, MPD, Ethanol, polyethylene glycol of various sizes, isopropanol, KCl, and the like (Ducruix).

Various additives can be used to aid in improving the character of the crystals, including substrate analogs, ligands, or inhibitors, as discussed in Part 2, below, as well as certain additives, including, but not limited to:

- 5 % Jeffamine
- 5 % Polypropyleneglycol P400
- 5 % Polyethyleneglycol 400
- 5 % ethyleneglycol
- 5 % 2-methyl-2,4-pentanediol
- 5 % Glycerol
- 5 % Dioxane
- 5 % dimethyl sulfoxide
- 5 % n-Octanol
- 100 mM $(\text{NH}_4)_2\text{SO}_4$
- 100 mM CsCl
- 100 mM CoSO_4
- 100 mM MnCl_2
- 100 mM KCl
- 100 mM ZnSO_4
- 100 mM LiCl_2
- 100 mM MgCl_2
- 100 mM Glucose
- 100 mM 1,6-Hexanediol 100 mM Dextran sulfate
- 100 mM 6-amino caproic acid
- 100 mM 1,6 hexane diamine
- 100 mM 1,8 diamino octane
- 100 mM Spermidine
- 100 mM Spermine
- 0.17 mM n-dodecyl- β -D-maltoside NP 40
- 20 mM n-octyl- β -D-glucopyranoside

According to one discovery of the present invention, the full-length β -secretase enzyme contains at least one transmembrane domain, and its purification is aided by the use

of a detergent (Triton X-100). Membrane proteins can be crystallized intact, but may require specialized conditions, such as the addition of a non-ionic detergent, such as C₈G (8-alkyl- β -glucoside) or an n-alkyl-maltoside (C₁₂M). Selection of such a detergent is somewhat empirical, but certain detergents are commonly employed. A number of membrane proteins have been successfully "salted out" by addition of high salt concentrations to the mixture. PEG has also been used successfully to precipitate a number of membrane proteins (Ducruix, *et al.*, *supra*). Alternatively, as discussed above, a C-terminal truncated form of the protein that binds inhibitor but which lacks the transmembrane domain, such as β -secretase 46-452 (SEQ ID NO: 58), is crystallized.

After crystallization conditions are determined, crystallization of a larger amount of the protein can be achieved by methods known in the art, such as vapor diffusion or equilibrium dialysis. In vapor diffusion, a drop of protein solution is equilibrated against a larger reservoir of solution containing precipitant or another dehydrating agent. After sealing, the solution equilibrates to achieve supersaturating concentrations of proteins and thereby induce crystallization in the drop.

Equilibrium dialysis can be used for crystallization of proteins at low ionic strength. Under these conditions, a phenomenon known as "salting in" occurs, whereby the protein molecules achieve balance of electrostatic charges through interactions with other protein molecules. This method is particularly effective when the solubility of the protein is low at the lower ionic strength. Various apparatuses and methods are used, including microdiffusion cells in which a dialysis membrane is attached to the bottom of a capillary tube, which may be bent at its lower portion. The final crystallization condition is achieved by slowly changing the composition of the outer solution. A variation of these methods utilizes a concentration gradient equilibrium dialysis set up. Microdiffusion cells are available from commercial suppliers such as Hampton Research (Laguna Niguel, CA).

Once crystallization is achieved, crystals characterized for purity (e.g., SDS-PAGE) and biological activity. Larger crystals (>0.2 mm) are preferred to increase the resolution of the X-ray diffraction, which is preferably on the order of 10-1.5 Angstroms. The selected crystals are subjected to X-ray diffraction, using a strong, monochromatic X-ray source, such as a Synchrotron source or rotating anode generator, and the resulting X-ray diffraction patterns are analyzed, using methods known in the art.

In one application, β -secretase amino acid sequence and/or X-ray diffraction data is recorded on computer readable medium, by which is meant any medium that can be read and

directly accessed by a computer. These data may be used to model the enzyme, a subdomain thereof, or a ligand thereof. Computer algorithms useful for this application are publicly and commercially available.

5 2. Crystallization of Protein plus Inhibitor

As mentioned above, it is advantageous to co-crystallize the protein in the presence of a binding ligand, such as inhibitor. Generally, the process for optimizing crystallization of the protein is followed, with addition of greater than 1 mM concentration of the inhibitor ligand during the precipitation phase. These crystals are also compared to crystals formed in
10 the absence of ligand, so that measurements of the ligand binding site can be made.

Alternatively, 1-2 μ l of 0.1-25 mM inhibitor compound is added to the drop containing crystals grown in the absence of inhibitor in a process known as "soaking." Based on the coordinates of the binding site, further inhibitor optimization is achieved. Such methods have been used advantageously in finding new, more potent inhibitors for HIV proteases
15 (See, e.g., Viswanadhan, V.N., *et al.* J. Med. Chem. 39: 705-712, 1996; Muegge, I., *et al.* J. Med. Chem. 42: 791-804, 1999).

One inhibitor ligand which is used in these co-crystallization and soaking experiments is P10—P4'staD->V (SEQ ID NO: 72), a statin peptide inhibitor described above. Methods for making the molecule are described herein. The inhibitor is mixed with β -secretase, and
20 the mixture is subjected to the same optimization tests described above, concentrating on those conditions worked out for the enzyme alone. Coordinates are determined and comparisons are made between the free and ligand bound enzyme, according to methods well known in the art. Further comparisons can be made by comparing the inhibitory concentrations of the enzyme to such coordinates, such as described by Viswanadhan, et al,
25 *supra*. Analysis of such comparisons provides guidance for design of further inhibitors, using this method.

D. Biological Activity of β -secretase

1. Naturally occurring β -secretase

30 In studies carried out in support of the present invention, isolated, purified forms of β secretase were tested for enzymatic activity using one or more native or synthetic substrates. For example, as discussed above, when β -secretase was prepared from human brain and

purified to homogeneity using the methods described in Example 5A, a single band was observed by silver stain after electrophoresis of sample fractions from the anion exchange chromatography (last step) on an SDS-polyacrylamide gel under reducing (+B-mercaptoethanol) conditions. As summarized in Table 1, above, this fraction yielded a specific activity of approximately 1.5×10^9 nM/h/mg protein, where activity was measured by hydrolysis of MBP-C125SW.

2. Isolated Recombinant β -secretase

Various recombinant forms of the enzyme were produced and purified from transfected cells. Since these cells were made to overproduce the enzyme, it was found that the purification scheme described with respect naturally occurring forms of the enzyme (e.g., Example 5A) could be shortened, with positive results. For example, as detailed in Example 6, 293T cells were transfected with pCEKclone 27 (FIG. 12 and FIG. 13A-E) (SEQ ID NO: 48) and Cos A2 cells were transfected with pCF β A2 using "FUGENE" 6 Transfection Reagent (Roche Molecular Biochemicals Research, Indianapolis, IN). The vector pCF was constructed from the parent vector pCDNA3, commercially available from Invitrogen, by inserting SEQ ID NO: 80 (FIG. 11A) between the HindIII and EcoRI sites. This sequence encompasses the adenovirus major late promoter tripartite leader sequence, and a hybrid splice created from adenovirus major late region first exon and intron and a synthetically generated IgG variable region splice acceptor.

pCDNA3 was cut with restriction endonucleases HindIII and EcoRI, then blunted by filling in the ends with Klenow fragment of DNA polymerase I. The cut and blunted vector was gel purified, and ligated with isolated fragment from pED.GI. The pED fragment was prepared by digesting with PvuII and SmaI, followed by gel purification of the resulting 419 base-pair fragment, which was further screened for orientation, and confirmed by sequencing.

To create the pCEK expression vector, the expression cassette from pCF was transferred into the EBV expression vector pCEP4 (Invitrogen, Carlsbad, CA). pCEP 4 was cut with BglII and XbaI, filled in, and the large 9.15 kb fragment containing pBR, hygromycin, and EBV sequences) ligated to the 1.9 kb NruI to XmnI fragment of pCF containing the expression cassette (CMV, TPL/MLP/IgG splice, Sp6, SVpolyA, M13 flanking region). pCF β A2 (clone A2) contains full length β -secretase in the vector pCF. pCF vector replicates in COS and 293T cells. In each case, cells were pelleted and a crude particulate fraction was prepared from the pellet. This fraction was extracted with buffer containing 0.2% Triton X-100. The Triton extract was diluted with pH 5.0 buffer and passed

through a SP Sepharose column. After the pH was adjusted to 4.5, β -secretase activity containing fractions were concentrated, with some additional purification on P10-P4'(statine)D->V Sepharose, as described for the brain enzyme. Silver staining of fractions revealed co-purified bands on the gel. Fractions corresponding to these bands were subjected to N-terminal amino acid determination. Results from these experiments revealed some heterogeneity of β -secretase species within the fractions. These species represent various forms of the enzyme; for example, from the 293T cells, the primary N-terminus is the same as that found in the brain, where (with respect to SEQ ID NO: 2) amino acid 46 is at the N-terminus. Minor amounts of protein starting just after the signal sequence (at residue 23) and at the start of the aspartyl protease homology domain (Met-63) were also observed. An additional major form of protein was found in Cos A2 cells, resulting from cleavage at Gly-58. These results are summarized in Table 3, above.

2. Comparison of Isolated, Naturally Occurring β -secretase with Recombinant

β -secretase

As described above, naturally occurring β -secretase derived from human brain as well as recombinant forms of the enzyme exhibit activity in cleaving APP, particularly as evidenced by activity in the MBP-C125 assay. Further, key peptide sequences from the naturally occurring form of the enzyme match portions of the deduced sequence derived from cloning the enzyme. Further confirmation that the two enzymes act identically can be taken from additional experiments in which various inhibitors were found to have very similar affinities for each enzyme, as estimated by a comparison of IC_{50} values measured for each enzyme under similar assay conditions. These inhibitors were discovered in accordance with a further aspect of the invention, which is described below. Significantly, the inhibitors produce near identical IC_{50} values and rank orders of potency in brain-derived and recombinant enzyme preparations, when compared in the same assay.

In further studies, comparisons were made between the full length recombinant enzyme having a C-terminal flag sequence "FLp501" (SEQ ID NO: 2, + SEQ ID NO: 45) and a recombinant enzyme truncated at position 452 "452Stop" (SEQ ID NO: 58 or SEQ ID NO: 59). Both enzymes exhibited activity in cleaving β -secretase substrates such as MBP-C125 as described above. The C-terminal truncated form of the enzyme exhibited activity in cleaving the MBP-C125sw substrate as well as the P26-P4' substrate, with similar rank order

of potency for the various inhibitor drugs tested. In addition, the absolute IC_{50} s were comparable for the two enzymes tested with the same inhibitor. All IC_{50} s were less than 10 μ M.

1. Cellular β -secretase

Further experiments carried out in support of the present invention have revealed that the isolated β -secretase polynucleotide sequences described herein encode β -secretase or β -secretase fragments that are active in cells. This section describes experiments carried out in support of the present invention, cells were transfected with DNA encoding β -secretase alone, or were co-transfected with DNA encoding secretase and DNA encoding wild-type APP as detailed in Example 8.

a. Transfection with β -secretase

In experiments carried out in support of the present invention, clones containing genes expressing the full-length polypeptide (SEQ ID NO: 2) were transfected into COS cells (Fugene and Effectene methods). Whole cell lysates were prepared and various amounts of lysate were tested for β -secretase activity according to standard methods known in the art or described in Example 4 herein. FIG. 14B shows the results of these experiments. As shown, lysates prepared from transfected cells, but not from mock- or control cells, exhibited considerable enzymatic activity in the MPB-C125sw assay, indicating "overexpression" of β -secretase by these cells.

b. Co-transfection of Cells with β -secretase and APP

In further experiments, 293T cells were co-transfected with pCEK clone 27, Figures 12 and 13 or poCK vector containing the full length β -secretase molecule (1-501; SEQ ID NO: 2) and with a plasmid containing either the wild-type or Swedish APP construct pohCK751, as described in Example 8. β -specific cleavage was analyzed by ELISA and Western analyses to confirm that the correct site of cleavage occurs.

Briefly, 293T cells were co-transfected with equivalent amounts of plasmids encoding β APPsw or wt and β -secretase or control β -galactosidase (β -gal) cDNA according to standard methods. β APP and β -secretase cDNAs were delivered via vectors, pohCK or pCEK, which do not replicate in 293T cells (pCEK-clone 27, FIGs. 12 and 13; pohCK751 expressing β APP 751, FIG. 21). Conditioned media and cell lysates were collected 48 hours after transfection. Western assays were carried out on conditioned media and cell lysates. ELISAs for

detection of A β peptide were carried out on the conditioned media to analyze various APP cleavage products.

Western Blot Results

It is known that β -secretase specifically cleaves at the Met-Asp in APPwt and the Leu-Asp in APPsw to produce the A β peptide, starting at position 1 and releasing soluble APP (sAPP β). Immunological reagents, specifically antibody 92 and 92sw (or 192sw), respectively, have been developed that specifically detect cleavage at this position in the APPwt and APPsw substrates, as described in U.S. Patent 5,721,130, incorporated herein by reference. Western blot assays were carried out on gels on which cell lysates were separated. These assays were performed using methods well known in the art, using as primary antibody reagents Ab 92 or Ab92S, which are specific for the C terminus of the N-terminal fragment of APP derived from APPwt and APPsw, respectively. In addition, ELISA format assays were performed using antibodies specific to the N terminal amino acid of the C terminal fragment.

Monoclonal antibody 13G8 (specific for C-terminus of APP-- epitope at positions 675-695 of APP695) was used in a Western blot format to determine whether the transfected cells express APP. FIG. 15A shows that reproducible transfection was obtained with expression levels of APP in vast excess over endogenous levels (triplicate wells are indicated as 1, 2, 3 in FIG. 15A). Three forms of APP-- mature, immature and endogenous-- can be seen in cells transfected with APPwt or APPsw. When β -secretase was co-transfected with APP, smaller C-terminal fragments appeared in triplicate well lanes from co-transfected cells (Western blot FIG. 15A, right-most set of lanes). In parallel experiments, where cells were co-transfected with β -secretase and APPsw substrate, literally all of the mature APP was cleaved (right-most set of lanes labeled "1,2,3" of FIG. 15B). This suggests that there is extensive cleavage by β -secretase of the mature APP (upper band), which results in C-terminal fragments of expected size in the lysate for cleavage at the β -secretase site. Co-transfection with Swedish substrate also resulted in an increase in two different sized CTF fragments (indicated by star). In conjunction with the additional Western and ELISA results described below, these results are consistent with a second cleavage occurring on the APPsw substrate after the initial cleavage at the β -secretase site.

Conditioned medium from the cells was analyzed for reactivity with the 192sw antibody, which is specific for β -s-APPsw. Analysis using this antibody indicated a dramatic increase in β -secretase cleaved soluble APP. This is observed in the gel illustrated in FIG.

16B by comparing the dark bands present in the "APPsw β sec" samples to the bands present in the "APPsw β gal" samples. Antibody specific for β -s-APPwt also indicates an increase in β -secretase cleaved material, as illustrated in FIG. 16A..

Since the antibodies used in these experiments are specific for the β -secretase cleavage site, the foregoing results show that p501 β -secretase cleaves APP at this site, and the overexpression of this recombinant clone leads to a dramatic enhancement of β -secretase processing at the correct β -secretase site in whole cells. This processing works on the wildtype APP substrate and is enhanced substantially on the Swedish APP substrate. Since approximately 20% of secreted APP in 293T cells is β -sAPP, with the increase observed below for APPsw, it is probable that almost all of the sAPP is β -sAPP. This observation was further confirmed by independent Western assays in which alpha and total sAPP were measured.

Monoclonal antibody 1736 is specific for the exposed α -secretase cleaved β -APP (Selkoe, et al.). When Western blots were performed using this antibody as primary antibody, a slight but reproducible decrease in α -cleaved APPwt was observed (FIG. 17A), and a dramatic decrease in α -cleaved APPsw material was also observed (note near absence of reactivity in FIG. 17B in the lanes labeled "APPsw β sec"). These results suggest that the overexpressed recombinant p501 β -secretase cleaves APPsw so efficiently or extensively that there is little or no substrate remaining for α -secretase to cleave. This further indicates that all the sAPP in APPsw β sec samples (illustrated in FIG 16B) is β -sAPP.

A β ELISA Results

Conditioned media from the recombinant cells was collected, diluted as necessary and tested for A β peptide production by ELISA on microtiter plates coated with monoclonal antibody 2G3, which is specific for recognizing the C-terminus of A β (1-40), with the detector reagent biotinylated mAb 3D6, which measures A β (x-40) (i.e., all N-terminus-truncated forms of the A β peptide). Overexpression of β -secretase with APPwt resulted in an approximately 8-fold increase in A β (x-40) production, with 1-40 representing a small percentage of the total. There was also a substantial increase in the production of A β 1-40 (FIG. 18). With APPsw there was an approximate 2-fold increase in A β (x-40). Without adhering to any particular underlying theory, it is thought that the less dramatic increase of A β (x-40) β -sec/APPsw cells in comparison to the β -sec/APPwt cells is due in part to the fact that processing of the APPsw substrate is much more efficient than that of the APPwt substrate. That is, a significant amount of APPsw is processed by endogenous β -secretase, so further increases upon transfection of β -secretase are therefore limited. These data indicate that the expression of recombinant β -secretase increases A β production and that β -secretase is rate limiting for production of A β in cells. This means that β -secretase enzymatic activity is rate limiting for production of A β in cells, and therefore provides a good therapeutic target.

IV. Utility

A. Expression Vectors and Cells Expressing β -secretase

The invention includes further cloning and expression of members of the aspartyl protease family described above, for example, by inserting polynucleotides encoding the proteins into standard expression vectors and transfecting appropriate host cells according to standard methods discussed below. Such expression vectors and cells expressing, for example, the human β -secretase enzyme described herein, have utility, for example, in producing components (purified enzyme or transfected cells) for the screening assays discussed in Part B, below. Such purified enzyme also has utility in providing starting materials for crystallization of the enzyme, as described in Section III, above. In particular, truncated form (s) of the enzyme, such as 46-452 (SEQ ID NO: 58), and the deglycosylated forms of the enzyme described herein are considered to have utility in this regard, as are other forms truncated partway into the transmembrane region, for example amino acid residues 46-458 in reference to SEQ ID NO: 2.

Polynucleotide sequences which encode human β -secretase, splice variants, fragments of the protein, fusion proteins, or functional equivalents thereof, collectively referred to herein as " β -secretase," may be used in recombinant DNA molecules that direct the expression of β -secretase in appropriate host cells. Due to the inherent degeneracy of the genetic code, other nucleic acid sequences that encode substantially the same or a functionally equivalent amino acid sequence may be used to clone and express β -secretase. Such variations will be readily ascertainable to persons skilled in the art.

The polynucleotide sequences of the present invention can be engineered in order to alter a β -secretase coding sequence for a variety of reasons, including but not limited to, alterations that modify the cloning, processing and/or expression of the gene product. For example, alterations may be introduced using techniques which are well known in the art, e.g., site-directed mutagenesis, to insert new restriction sites, to alter glycosylation patterns, to change codon preference, to produce splice variants, etc. For example, it may be advantageous to produce β -secretase -encoding nucleotide sequences possessing non-naturally occurring codons. Codons preferred by a particular prokaryotic or eukaryotic host (Murray, E. *et al.* (1989) *Nuc Acids Res* 17:477-508) can be selected, for example, to increase the rate of β -secretase polypeptide expression or to produce recombinant RNA transcripts having desirable properties, such as a longer half-life, than transcripts produced from naturally occurring sequence. This may be particularly useful in producing recombinant enzyme in non-mammalian cells, such as bacterial, yeast, or insect cells. The present invention also includes recombinant constructs comprising one or more of the sequences as broadly described above. The constructs comprise a vector, such as a plasmid or viral vector, into which a sequence of the invention has been inserted, in a forward or reverse orientation. In a preferred aspect of this embodiment, the construct further comprises regulatory sequences, including, for example, a promoter, operably linked to the sequence. Large numbers of suitable vectors and promoters are known to those of skill in the art, and are commercially available. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are also described in Sambrook, *et al.*, (*supra*).

The present invention also relates to host cells that are genetically engineered with vectors of the invention, and the production of proteins and polypeptides of the invention by recombinant techniques. Host cells are genetically engineered (i.e., transduced, transformed

or transfected) with the vectors of this invention which may be, for example, a cloning vector or an expression vector. The vector may be, for example, in the form of a plasmid, a viral particle, a phage, etc. The engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying the β -secretase gene. The culture conditions, such as temperature, pH and the like, are those previously used with the host cell selected for expression, and will be apparent to those skilled in the art. Exemplary methods for transfection of various types of cells are provided in Example 6, herein.

As described above, according to a preferred embodiment of the invention, host cells can be co-transfected with an enzyme substrate, such as with APP (such as wild type or Swedish mutation form), in order to measure activity in a cell environment. Such host cells are of particular utility in the screening assays of the present invention, particularly for screening for therapeutic agents that are able to traverse cell membranes.

The polynucleotides of the present invention may be included in any of a variety of expression vectors suitable for expressing a polypeptide. Such vectors include chromosomal, nonchromosomal and synthetic DNA sequences, e.g., derivatives of SV40; bacterial plasmids; phage DNA; baculovirus; yeast plasmids; vectors derived from combinations of plasmids and phage DNA, viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudorabies. However, any other vector may be used as long as it is replicable and viable in the host. The appropriate DNA sequence may be inserted into the vector by a variety of procedures. In general, the DNA sequence is inserted into an appropriate restriction endonuclease site(s) by procedures known in the art. Such procedures and related sub-cloning procedures are deemed to be within the scope of those skilled in the art.

The DNA sequence in the expression vector is operatively linked to an appropriate transcription control sequence (promoter) to direct mRNA synthesis. Examples of such promoters include: CMV, LTR or SV40 promoter, the *E. coli* lac or trp promoter, the phage lambda PL promoter, and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or their viruses. The expression vector also contains a ribosome binding site for translation initiation, and a transcription terminator. The vector may also include appropriate sequences for amplifying expression. In addition, the expression vectors preferably contain one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells such as dihydrofolate reductase or

neomycin resistance for eukaryotic cell culture, or such as tetracycline or ampicillin resistance in *E. coli*.

The vector containing the appropriate DNA sequence as described above, as well as an appropriate promoter or control sequence, may be employed to transform an appropriate host to permit the host to express the protein. Examples of appropriate expression hosts include: bacterial cells, such as *E. coli*, *Streptomyces*, and *Salmonella typhimurium*; fungal cells, such as yeast; insect cells such as *Drosophila* and *Spodoptera Sf9*; mammalian cells such as CHO, COS, BHK, HEK 293 or Bowes melanoma; adenoviruses; plant cells, etc. It is understood that not all cells or cell lines will be capable of producing fully functional β -secretase; for example, it is probable that human β -secretase is highly glycosylated in native form, and such glycosylation may be necessary for activity. In this event, eukaryotic host cells may be preferred. The selection of an appropriate host is deemed to be within the scope of those skilled in the art from the teachings herein. The invention is not limited by the host cells employed.

In bacterial systems, a number of expression vectors may be selected depending upon the use intended for β -secretase. For example, when large quantities of β -secretase or fragments thereof are needed for the induction of antibodies, vectors, which direct high level expression of fusion proteins that are readily purified, may be desirable. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as Bluescript(R) (Stratagene, La Jolla, CA), in which the β -secretase coding sequence may be ligated into the vector in-frame with sequences for the amino-terminal Met and the subsequent 7 residues of beta-galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke & Schuster (1989) J Biol Chem 264:5503-5509); pET vectors (Novagen, Madison WI); and the like.

In the yeast *Saccharomyces cerevisiae* a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase and PGH may be used. For reviews, see Ausubel *et al.* (supra) and Grant *et al.* (1987; Methods in Enzymology 153:516-544).

In cases where plant expression vectors are used, the expression of a sequence encoding β -secretase may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV (Brisson *et al.* (1984) Nature 310:511-514) may be used alone or in combination with the omega leader sequence from

C

TMV (Takamatsu *et al.* (1987) EMBO J 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO (Coruzzi *et al.* (1984) EMBO J 3:1671-1680; Broglie *et al.* (1984) Science 224:838-843), or heat shock promoters (Winter J and Sinibaldi RM (1991) Results. Probl. Cell Differ. 17:85-105) may be used. These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. For reviews of such techniques, see Hobbs S or Murry LE (1992) in McGraw Hill Yearbook of Science and Technology, McGraw Hill, New York, N.Y., pp 191-196; or Weissbach and Weissbach (1988) Methods for Plant Molecular Biology, Academic Press, New York, N.Y., pp 421-463.

β -secretase may also be expressed in an insect system. In one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* Sf9 cells or in *Trichoplusia* larvae. The β -secretase coding sequence is cloned into a nonessential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of Kv-SL coding sequence will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein coat. The recombinant viruses are then used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which β -secretase is expressed (Smith *et al.* (1983) J Virol 46:584; Engelhard EK *et al.* (1994) Proc Natl Acad Sci 91:3224-3227).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, a β -secretase coding sequence may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a nonessential E1 or E3 region of the viral genome will result in a viable virus capable of expressing the enzyme in infected host cells (Logan and Shenk (1984) Proc Natl Acad Sci 81:3655-3659). In addition, transcription enhancers, such as the rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

Specific initiation signals may also be required for efficient translation of a β -secretase coding sequence. These signals include the ATG initiation codon and adjacent sequences. In cases where β -secretase coding sequence, its initiation codon and upstream sequences are inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous transcriptional control signals including the ATG initiation codon must be provided. Furthermore, the initiation codon must be in the correct reading

frame to ensure transcription of the entire insert. Exogenous transcriptional elements and initiation codons can be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate to the cell system in use (Scharf D *et al.* (1994) Results Probl Cell Differ 20:125-62; Bittner *et al.* (1987) Methods in Enzymol 153:516-544).

In a further embodiment, the present invention relates to host cells containing the above-described constructs. The host cell can be a higher eukaryotic cell, such as a mammalian cell, or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-Dextran mediated transfection, or electroporation (Davis, L., Dibner, M., and Battey, I. (1986) Basic Methods in Molecular Biology) or newer methods, including lipid transfection with "FUGENE" (Roche Molecular Biochemicals, Indianapolis, IN) or "EFFECTENE" (Quiagen, Valencia, CA), or other DNA carrier molecules. Cell-free translation systems can also be employed to produce polypeptides using RNAs derived from the DNA constructs of the present invention.

A host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the protein include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be important for correct insertion, folding and/or function. For example, in the case of β -secretase, it is likely that the N-terminus of SEQ ID NO: 2 is truncated, so that the protein begins at amino acid 22, 46 or 57-58 of SEQ ID NO: 2. Different host cells such as CHO, HeLa, BHK, MDCK, 293, WI38, etc. have specific cellular machinery and characteristic mechanisms for such post-translational activities and may be chosen to ensure the correct modification and processing of the introduced, foreign protein.

For long-term, high-yield production of recombinant proteins, stable expression may be preferred. For example, cell lines that stably express β -secretase may be transformed using expression vectors which contain viral origins of replication or endogenous expression elements and a selectable marker gene. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells that successfully express the introduced

sequences. Resistant clumps of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type. For example, in experiments carried out in support of the present invention, overexpression of the "452stop" form of the enzyme has been achieved.

5 Host cells transformed with a nucleotide sequence encoding β -secretase may be cultured under conditions suitable for the expression and recovery of the encoded protein from cell culture. The protein or fragment thereof produced by a recombinant cell may be secreted, membrane-bound, or contained intracellularly, depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors
10 containing polynucleotides encoding β -secretase can be designed with signal sequences which direct secretion of β -secretase polypeptide through a prokaryotic or eukaryotic cell membrane.

β -secretase may also be expressed as a recombinant protein with one or more additional polypeptide domains added to facilitate protein purification. Such purification
15 facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp, Seattle, Wash.). The inclusion of a protease-cleavable polypeptide linker sequence between the purification domain and β -
20 secretase is useful to facilitate purification. One such expression vector provides for expression of a fusion protein comprising β -secretase (e.g., a soluble β -secretase fragment) fused to a polyhistidine region separated by an enterokinase cleavage site. The histidine residues facilitate purification on IMIAC (immobilized metal ion affinity chromatography, as
described in Porath *et al.* (1992) Protein Expression and Purification 3:263-281) while the
25 enterokinase cleavage site provides a means for isolating β -secretase from the fusion protein. pGEX vectors (Promega, Madison, Wis.) may also be used to express foreign polypeptides fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to ligand-agarose beads (e.g., glutathione-agarose in the case of GST-fusions) followed by elution in the presence of
30 free ligand.

Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is induced by appropriate means (e.g.,

temperature shift or chemical induction) and cells are cultured for an additional period. Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification. Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents, or other methods, which are well known to those skilled in the art.

β -secretase can be recovered and purified from recombinant cell cultures by any of a number of methods well known in the art, or, preferably, by the purification scheme described herein. Protein refolding steps can be used, as necessary, in completing configuration of the mature protein. Details of methods for purifying naturally occurring as well as purified forms of β -secretase are provided in the Examples.

B. Methods of Selecting β -secretase Inhibitors

The present invention also includes methods for identifying molecules, such as synthetic drugs, antibodies, peptides, or other molecules, which have an inhibitory effect on the activity of β -secretase described herein, generally referred to as inhibitors, antagonists or blockers of the enzyme. Such an assay includes the steps of providing an isolated protein, about 450 amino acid residues in length, which includes an amino acid sequence that is at least 90% identical to SEQ ID NO: 75 [63-423] including conservative substitutions thereof, which exhibits β -secretase activity, as described herein. The β -secretase enzyme is contacted with a test compound to determine whether it has a modulating effect on the activity of the enzyme, as discussed below, and selecting from test compounds capable of modulating β -secretase activity. In particular, inhibitory compounds (antagonists) are useful in the treatment of disease conditions associated with amyloid deposition, particularly Alzheimer's disease. Persons skilled in the art will understand that such assays may be conveniently transformed into kits.

Particularly useful screening assays employ cells which express both β -secretase and APP. Such cells can be made recombinantly by co-transfection of the cells with polynucleotides encoding the proteins, as described in Section III, above, or can be made by transfecting a cell which naturally contains one of the proteins with the second protein. In a

particular embodiment, such cells are grown up in multi-well culture dishes and are exposed to varying concentrations of a test compound or compounds for a pre-determined period of time, which can be determined empirically. Whole cell lysates, cultured media or cell membranes are assayed for β -secretase activity. Test compounds which significantly inhibit activity compared to control (as discussed below) are considered therapeutic candidates.

Isolated β -secretase, its ligand-binding, catalytic, or immunogenic fragments, or oligopeptides thereof, can be used for screening therapeutic compounds in any of a variety of drug screening techniques. The protein employed in such a test may be membrane-bound, free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly.

The formation of binding complexes between β -secretase and the agent being tested can be measured. Compounds that inhibit binding between β -secretase and its substrates, such as APP or APP fragments, may be detected in such an assay. Preferably, enzymatic activity will be monitored, and candidate compounds will be selected on the basis of ability to inhibit such activity. More specifically, a test compound will be considered as an inhibitor of β -secretase if the measured β -secretase activity is significantly lower than β -secretase activity measured in the absence of test compound. In this context, the term "significantly lower" means that in the presence of the test compound the enzyme displays an enzymatic activity which, when compared to enzymatic activity measured in the absence of test compound, is measurably lower, within the confidence limits of the assay method. Such measurements can be assessed by a change in K_m and/or V_{max} , single assay endpoint analysis, or any other method standard in the art. Exemplary methods for assaying β -secretase are provided in Example 4 herein.

For example, in studies carried out in support of the present invention, compounds were selected based on their ability to inhibit β -secretase activity in the MBP-C125 assay. Compounds that inhibited the enzyme activity at a concentration lower than about 50 μM were selected for further screening.

The groups of compounds that are most likely candidates for inhibitor activity comprise a further aspect of the present invention. Based on studies carried out in support of the invention, it has been determined that the peptide compound described herein as P10-P4'staD->V (SEQ ID NO: 72) is a reasonably potent inhibitor of the enzyme. Further studies based on this sequence and peptidomimetics of portions of this sequence have revealed a number of small molecule inhibitors.

Random libraries of peptides or other compounds can also be screened for suitability as β -secretase inhibitors. Combinatorial libraries can be produced for many types of compounds that can be synthesized in a step-by-step fashion. Such compounds include polypeptides, beta-turn mimetics, polysaccharides, phospholipids, hormones, prostaglandins, steroids, aromatic compounds, heterocyclic compounds, benzodiazepines, oligomeric N-substituted glycines and oligocarbamates. Large combinatorial libraries of the compounds can be constructed by the encoded synthetic libraries (ESL) method described in Affymax, WO 95/12608, Affymax, WO 93/06121, Columbia University, WO 94/08051, Pharmacocea, WO 95/35503 and Scripps, WO 95/30642 (each of which is incorporated by reference for all purposes).

A preferred source of test compounds for use in screening for therapeutics or therapeutic leads is a phage display library. See, e.g., Devlin, WO 91/18980; Key, B.K., *et al.*, eds., *Phage Display of Peptides and Proteins, A Laboratory Manual*, Academic Press, San Diego, CA, 1996. Phage display is a powerful technology that allows one to use phage genetics to select and amplify peptides or proteins of desired characteristics from libraries containing 10^8 - 10^9 different sequences. Libraries can be designed for selected variegation of an amino acid sequence at desired positions, allowing bias of the library toward desired characteristics. Libraries are designed so that peptides are expressed fused to proteins that are displayed on the surface of the bacteriophage. The phage displaying peptides of the desired characteristics are selected and can be regrown for expansion. Since the peptides are amplified by propagation of the phage, the DNA from the selected phage can be readily sequenced facilitating rapid analyses of the selected peptides.

Phage encoding peptide inhibitors can be selected by selecting for phage that bind specifically to β -secretase protein. Libraries are generated fused to proteins such as gene II that are expressed on the surface of the phage. The libraries can be composed of peptides of various lengths, linear or constrained by the inclusion of two Cys amino acids, fused to the phage protein or may also be fused to additional proteins as a scaffold. One may start with libraries composed of random amino acids or with libraries that are biased to sequences in the β APP substrate surrounding the β -secretase cleavage site or preferably, to the D \rightarrow V substituted site exemplified in SEQ ID NO: 72. One may also design libraries biased toward the peptidic inhibitors and substrates described herein or biased toward peptide sequences obtained from the selection of binding phage from the initial libraries provide additional test inhibitor compound.

(The β -secretase is immobilized and phage specifically binding to the β -secretase selected for. Limitations, such as a requirement that the phage not bind in the presence of a known active site inhibitor of β -secretase (e.g. the inhibitors described herein), serve to further direct phage selection active site specific compounds. This can be complicated by a differential selection format. Highly purified β -secretase, derived from brain or preferably from recombinant cells can be immobilized to 96-well plastic dishes using standard techniques (reference phage book). Recombinant β -secretase, designed to be fused to a peptide that can bind (e.g. streptavidin binding motifs, His, FLAG or myc tags) to another protein immobilized (such as streptavidin or appropriate antibodies) on the plastic petri dishes can also be used. Phage are incubated with the bound β -secretase and unbound phage removed by washing. The phage are eluted and this selection is repeated until a population of phage binding to β -secretase is recovered. Binding and elution are carried out using standard techniques.

Alternatively β -secretase can be "bound" by expressing it in Cos or other mammalian cells growing on a petri dish. In this case one would select for phage binding to the β -secretase expressing cells, and select against phage that bind to the control cells, that are not expressing β -secretase.

One can also use phage display technology to select for preferred substrates of β -secretase, and incorporate the identified features of the preferred substrate peptides obtained by phage display into inhibitors.

In the case of β -secretase, knowledge of the amino acid sequence surrounding the cleavage site of APP and of the cleavage site of APP^{sw} has provided information for purposes of setting up the phage display screening library to identify preferred substrates of β -secretase. As mentioned above, knowledge of the sequence of a particularly good peptide inhibitor, P10-P4staD->V (SEQ ID NO: 72), as described herein, provides information for setting up a "biased" library toward this sequence.

For example, the peptide substrate library containing 10^8 different sequences is fused to a protein (such as a gene III protein) expressed on the surface of the phage and a sequence that can be used for binding to streptavidin, or another protein, such as His tag and antibody to His. The phage are digested with protease, and undigested phage are removed by binding to appropriate immobilized binding protein, such as streptavidin. This selection is repeated until a population of phage encoding substrate peptide sequences is recovered. The DNA in

(
the phage is sequenced to yield the substrate sequences. These substrates are then used for further development of peptidomimetics, particularly peptidomimetics having inhibitory properties.

Combinatorial libraries and other compounds are initially screened for suitability by
5 determining their capacity to bind to, or preferably, to inhibit β -secretase activity in any of the assays described herein or otherwise known in the art. Compounds identified by such screens are then further analyzed for potency in such assays. Inhibitor compounds can then be tested for prophylactic and therapeutic efficacy in transgenic animals predisposed to an amyloidogenic disease, such as various rodents bearing a human APP-containing transgene,
10 e.g., mice bearing a 717 mutation of APP described by Games et al., *Nature* 373: 523-527, 1995 and Wadsworth et al. (US 5,811,633, US 5,604,131, US 5,720,936), and mice bearing a Swedish mutation of APP such as described by McConlogue et al. (US 5,612,486) and Hsiao et al. (U.S. 5,877,399); Staufenbiel et al., *Proc. Natl. Acad. Sci. USA* 94, 13287-13292 (1997); Sturchler-Pierrat et al., *Proc. Natl. Acad. Sci. USA* 94, 13287-13292 (1997); Borchelt et al.,
15 *Neuron* 19, 939-945 (1997), all of which are incorporated herein by reference.

Compounds or agents found to be efficacious and safe in such animal models will be further tested in standard toxicological assays. Compounds showing appropriate toxicological and pharmacokinetic profiles will be moved into human clinical trials for treatment of Alzheimer's disease and related diseases. The same screening approach can be
20 used on other potential agents such as peptidomimetics described above.

In general, in selecting therapeutic compounds based on the foregoing assays, it is useful to determine whether the test compound has an acceptable toxicity profile, e.g., in a variety of in vitro cells and animal model(s). It may also be useful to search the tested and identified compound(s) against existing compound databases to determine whether the
25 compound or analogs thereof have been previously employed for pharmaceutical purposes, and if so, optimal routes of administration and dose ranges. Alternatively, routes of administration and dosage ranges can be determined empirically, using methods well known in the art (see, e.g., Benet, L.Z., et al. *Pharmacokinetics in Goodman & Gilman's The Pharmacological Basis of Therapeutics*, Ninth Edition, Hardman, J.G., et al., Eds., McGraw-Hill, New York, 1966) applied to standard animal models, such as a transgenic PDAPP
30 animal model (e.g., Games, D., et al. *Nature* 373: 523-527, 1995; Johnson-Wood, K., et al., *Proc. Natl. Acad. Sci. USA* 94: 1550-1555, 1997). To optimize compound activity and/or specificity, it may be desirable to construct a library of near-neighbor analogs to search for

analogues with greater specificity and/or activity. Methods for synthesizing near-neighbor and/or targeted compound libraries are well-known in the combinatorial library field.

C. Inhibitors and Therapeutics

Part B, above, describes method of screening for compounds having β -secretase inhibitory activity. To summarize, guidance is provided for specific methods of screening for potent and selective inhibitors of β -secretase enzyme. Significantly, the practitioner is directed to a specific peptide substrate/inhibitor sequences, such as P10-P4'staD->V (SEQ ID NO: 72), on which drug design can be based and additional sources, such as biased phage display libraries, that should provide additional lead compounds.

The practitioner is also provided ample guidance for further refinement of the binding site of the enzyme, for example, by crystallizing the purified enzyme in accord with the methods provide herein. Noting the success in this area that has been enjoyed in the area of HIV protease inhibitor development, it is contemplated that such efforts will lead to further optimization of the test compounds described herein. With optimized compounds in hand, it is possible to define a compound pharmacophore, and further search existing pharmacophore databases, e.g., as provided by Tripos, to identify other compounds that may differ in 2-D structural formulae with the originally discovered compounds, but which share a common pharmacophore structure and activity. Test compounds are assayed in any of the inhibitor assays described herein, at various stages in development. Therefore, β -secretase inhibitory agents can be discovered by any of the methods described herein, particularly the inhibitor assays and the crystallization/optimization protocols. Such inhibitory agents are therapeutic candidates for treatment of Alzheimer's disease, as well as other amyloidoses characterized by A β peptide deposition. The considerations concerning therapeutic index (toxicology), bioavailability and dosage discussed in Part B above are also important to consider with respect to these therapeutic candidates.

D. Methods of Diagnosis

The present invention also provides methods of diagnosing individuals who carry mutations that provide enhanced β -secretase activity. For example, there are forms of familial Alzheimer's disease in which the underlying genetic disorder has yet to be recognized. Members of families possessing this genetic predisposition can be monitored for alterations in the nucleotide sequence that encodes β -secretase and/or promoter regions

thereof, since it is apparent, in view of the teachings herein, that individuals who overexpress of the enzyme or possess catalytically more efficient forms of the enzyme would be likely to produce relatively more A β peptide. Support for this supposition is provided by the observation, reported herein, that the amount of β -secretase enzyme is rate limiting for production of A β in cells.

More specifically, persons suspected to have a predilection for developing for developing or who already have the disease, as well as members of the general population, may be screened by obtaining a sample of their cells, which may be blood cells or fibroblasts, for example, and testing the samples for the presence of genetic mutations in the β -secretase gene, in comparison to SEQ ID NO: 1 described herein, for example. Alternatively or in addition, cells from such individuals can be tested for β -secretase activity. According to this embodiment, a particular enzyme preparation might be tested for increased affinity and/or Vmax with respect to a β -secretase substrate such as MBP-C125, as described herein, with comparisons made to the normal range of values measured in the general population.

Individuals whose β -secretase activity is increased compared to normal values are susceptible to developing Alzheimer's disease or other amyloidogenic diseases involving deposition of A β peptide.

E. Therapeutic Animal Models

A further utility of the present invention is in creation of certain transgenic and/or knockout animals that are also useful in the screening assays described herein. Of particular use is a transgenic animal that overexpresses the β -secretase enzyme, such as by adding an additional copy of the mouse enzyme or by adding the human enzyme. Such an animal can be made according to methods well known in the art (e.g., Cordell, U.S. Patent 5,387,742; Wadsworth et al., US 5,811,633, US 5,604,131, US 5,720,936; McConlogue et al., US 5,612,486; Hsiao et al., U.S. 5,877,399; and "Manipulating the Mouse Embryo, A Laboratory Manual," B. Hogan, F. Costantini and E. Lacy, Cold Spring Harbor Press, 1986)), substituting the one or more of the constructs described with respect to β -secretase, herein, for the APP constructs described in the foregoing references, all of which are incorporated by reference.

An overexpressing β -secretase transgenic mouse will make higher levels of A β and s β APP from APP substrates than a mouse expressing endogenous β -secretase. This would

(facilitate analysis of APP processing and inhibition of that processing by candidate therapeutic agents. The enhanced production of A β peptide in mice transgenic for β -secretase would allow acceleration of AD-like pathology seen in APP transgenic mice. This result can be achieved by either crossing the β -secretase expressing mouse onto a mouse displaying
5 AD-like pathology (such as the PDAPP or Hsiao mouse) or by creating a transgenic mouse expressing both the β -secretase and APP transgene.

Such transgenic animals are used to screen for β -secretase inhibitors, with the advantage that they will test the ability of such inhibitors to gain entrance to the brain and to effect inhibition *in vivo*.

10 A so-called "knock-out mouse" in which the endogenous enzyme is either permanently (as described in US Patent Nos. 5,464,764, 5,627,059 and 5,631,153, which are incorporated by reference in their entirety) or inducibly deleted (as described in US Patent No. 4,959,317, which is incorporated by reference in its
15 entirety), or which is inactivated, is described in further detail below. Such mice serve as controls for β -secretase activity and/or can be crossed with APP mutant mice, to provide validation of the pathological sequelae. Such mice can also provide a screen for other drug targets, such as drugs specifically directed at A β deposition events.

β -secretase knockout mice provide a model of the potential effects of β -secretase inhibitors *in vivo*. Comparison of the effects of β -secretase test inhibitors *in vivo* to the
20 phenotype of the β -secretase knockout can help guide drug development. For example, the phenotype may or may not include pathologies seen during drug testing of β -secretase inhibitors. If the knockout does not show pathologies seen in the drug-treated mice, one could infer that the drug is interacting non-specifically with another target in addition to the β -secretase target. Tissues from the knockout can be used to set up drug binding assays or to
25 carry out expression cloning to find the targets that are responsible for these toxic effects. Such information can be used to design further drugs that do not interact with these undesirable targets. The knockout mice will facilitate analyses of potential toxicities that are inherent to β -secretase inhibition. Knowledge of potential toxicities will help guide the design of design drugs or drug-delivery systems to reduce such toxicities. Inducible knockout
30 mice are particularly useful in distinguishing toxicity in an adult animal from embryonic

effects seen in the standard knockout. If the knockout confers fetal-lethal effects, the inducible knockout will be advantageous.

Methods and technology for developing knock-out mice have matured to the point that a number of commercial enterprises generate such mice on a contract basis (e.g., Lexicon Genetics, Woodland TX; Cell & Molecular Technologies, Lavallete, NJ; Crysalis, DNX Transgenic Sciences, Princeton, NJ). Methodologies are also available in the art. (See Galli-Taliadoros, L.A., *et al.*, J. Immunol. Meth. 181: 1-15, 1995). Briefly, a genomic clone of the enzyme of interest is required. Where, as in the present invention, the exons encoding the regions of the protein have been defined, it is possible to achieve inactivation of the gene without further knowledge of the regulatory sequences controlling transcription. Specifically, a mouse strain 129 genomic library can be screened by hybridization or PCR, using the sequence information provided herein, according to methods well known in the art. (Ausubel; Sambrook) The genomic clone so selected is then subjected to restriction mapping and partial exonic sequencing for confirmation of mouse homologue and to obtain information for knock-out vector construction. Appropriate regions are then sub-cloned into a "knock-out" vector carrying a selectable marker, such as a vector carrying a *neo*^r cassette, which renders cells resistant to aminoglycoside antibiotics such as gentamycin. The construct is further engineered for disruption of the gene of interest, such as by insertion of a sequence replacement vector, in which a selectable marker is inserted into an exon of the gene, where it serves as a mutagen, disrupting the coordinated transcription of the gene. Vectors are then engineered for transfection into embryonic stem (ES) cells, and appropriate colonies are isolated. Positive ES cell clones are micro-injected into isolated host blastocysts to generate chimeric animals, which are then bred and screened for germline transmission of the mutant allele.

β -secretase knock-out mice can be generated such that the mutation is inducible, such as by inserting in the knock-out mice a lox region flanking the β -secretase gene region. Such mice are then crossed with mice bearing a "Cre" gene under an inducible promoter, resulting in at least some off-spring bearing both the "Cre" and the lox constructs. When expression of "Cre" is induced, it serves to disrupt the gene flanked by the lox constructs. Such a "Cre-lox" mouse is particularly useful, when it is suspected that the knock-out mutation may be lethal. In addition, it provides the opportunity for knocking out the gene in selected tissues, such as the brain. Methods for generating Cre-

(lox constructs are provided by U.S. Patent 4,959,317, incorporated herein by reference, and are made on a contractual basis by Lexicon Genetics, Woodlands, TX, among others.

The following examples illustrate, but in no way are intended to limit the present invention.

5 Example 1

Isolation of Coding Sequences for Human β -secretase

A. PCR Cloning

(Poly A⁺ RNA from IMR human neuroblastoma cells was reverse transcribed using the Perkin-Elmer kit. Eight degenerate primer pools, each 8 fold degenerate, encoding the
10 N and C terminal portions of the amino acid sequence obtained from the purified protein were designed (shown in Table 4; oligos 3407 through 3422)(SEQ ID NOS: 3-21). PCR reaction composed of cDNA from 10 ng of RNA, 1.5 mM MgCl₂, 0.125 μ l AmpliTaq® Gold, 160 μ M each dNTP (plus 20 μ M additional from the reverse transcriptase reaction), Perkin-Elmer TAQ buffer (from AmpliTaq® Gold kit, Perkin-Elmer, Foster City, CA), in a 25 μ l reaction
15 volume. Each of oligonucleotide primers 3407 through 3414 was used in combination with each of oligos 3415 through 3422 for a total for 64 reactions. Reactions were run on the Perkin-Elmer 7700 Sequence Detection machine under the following conditions: 10 min at 95°C, 4 cycles of, 45 °C annealing for 15 second, 72 °C extension for 45 second and 95 °C denaturation for 15 seconds followed by 35 cycles under the same conditions with the
20 exception that the annealing temperature was raised to 55 °C. (The foregoing conditions are referred to herein as "Reaction 1 conditions.") PCR products were visualized on 4% agarose gel (Northern blots) and a prominent band of the expected size (68 bp) was seen in reactions, particularly with the primers 3515-3518. The 68 kb band was sequenced and the internal region coded for the expected amino acid sequence. This gave the exact DNA sequence for
25 22 bp of the internal region of this fragment.

Additional sequence was deduced from the efficiency of various primer pools of discrete sequence in generating this PCR product. Primer pools 3419 to 3422 (SEQ ID NOS: 15-18) gave very poor or no product, whereas pools 3415 to 3418 (SEQ ID NOS: 11-14 respectively) gave robust signal. The difference between these pools is a CTC (3415
30 to 3418) (SEQ ID NOS: 11-14) vs TTC (3419 to 3422) (SEQ ID NOS: 15-18) in the 3' most end of the pools. Since CTC primed more efficiently we can conclude that the reverse complement GAG is the correct codon. Since Met coding is unique it was concluded that the following codon is ATG. Thus the exact DNA sequence obtained is:

CCC.GGC.CGG.AGG.GGC.AGC.TTT.GTG.GAG.ATG.GT (SEQ ID NO: 49)
encoding the amino acid sequence P G R R G S F V E M V (SEQ ID NO: 50). This
sequence can be used to design exact oligonucleotides for 3 and 5' RACE PCR on either
cDNA or libraries or to design specific hybridization probes to be used to screen libraries.

5 Since the degenerate PCR product was found to be so robust, this reaction may also be
used as a diagnostic for the presence of clones containing this sequence. Pools of libraries
can be screened using this PCR product to indicate the presence of a clone in the pool. The
pools can be broken out to identify individual clones. Screening pools of known complexity
and or size can provide information on the abundance of this clone in a library or source and
10 can approximate the size of the full length clone or message.

For generation of a probe, PCR reactions using oligonucleotides 3458 (SEQ ID
NO: 19) and 3469 (SEQ ID NO: 21) or 3458 (SEQ ID NO: 19) and 3468 (SEQ ID
NO: 20) (Table 4) can be carried out using the 23 RACE product, clone 9C7E.35 (30
ng, clone 9C7E.35 was isolated from origene library, see Example 2), or cDNA
15 generated from brain, using the standard PCR conditions (Perkin-Elmer, rtPCR and
AmpliTaq® Gold kits) with the following: 25 µl reaction volume 1.5 mM MgCl₂,
0.125 µl of AmpliTaq® Gold (Perkin-Elmer), initial 95° for 10 min to activate the
AmpliTaq® Gold, 36 cycles of 65° 15 sec 72° 45 sec 95° for 15 sec, followed by 3
min at 72°. Product was purified on a Quiagen PCR purification kit and used as a
20 substrate for randompriming to generate a radiolabelled probe (Sambrook, *et al.*,
supra; Amersham RediPrime® kit). This probe was used to isolate full length clone
pCEK clone 27 shown in FIGS. 12 and 13 (A-E) (SEQ ID NO: 48).

Derivation of full length clone pCEK clone 27

A human primary neuronal cell library in the mammalian expression vector pCEK2
25 vector was generated using size selected cDNA, and pools of clones generated from different
sized inserts. The cDNA library for β-secretase screening was made with poly(A)⁺ RNA
isolated from primary human neuronal cells. The cloning vector was pCEK2 (FIG. 12).

30

pCEK2

Double-stranded cDNA inserts were synthesized using the cDNA Synthesis Kit from
Stratagene with some modifications. The inserts were then fractionated according to their
sizes. A total of five fractions were individually ligated with double-cut (NotI and XhoI)

(pCEK2 and subsequently transformed into the E. Coli strain XL-10 Gold which is designed to accept very large plasmids.

The fractions of transformed E. Coli were plated on Terrific Broth agar plates containing ampicillin and let grown for 18 hours. Each fraction yielded about 200,000 colonies to give a total of one million colonies. The colonies were then scraped from the plates and plasmids isolated from them in pools of approximately 70,000 clones/pool. 70,000 clones from each pool of the library was screened for the presence of the putative β -secretase gene using the diagnostic PCR reaction (degenerate primers 3411 (SEQ ID NO: 7) and 3417 (SEQ ID NO: 13) shown above).

10 Clones from the 1.5 kb pool were screened using a radiolabeled probe generated from a 390 b.p. PCR product generated from clone 9C7E.35. For generation of a probe, PCR product was generated using 3458 (SEQ ID NO: 19) and 3468 (SEQ ID NO: 20) as primers and clone 9C7E.35 (30 ng) as substrate.

15 PCR product was used as a substrate for random priming to generate a radiolabeled probe. 180,000 clones from the 1.5 kb pool (70,000 original clones in this pool), were screened by hybridization with the PCR probe and 9 positive clones identified. Four of these clones were isolated and by restriction mapping these appear to encode two independent clones of 4 to 5 kb (clone 27) and 6 to 7 kb (clone 53) length. Sequencing of clone 27 verified that it contains a coding region of 1.5 kb. FIG. 13 (A-F) shows the sequence of pCEK clone27 (clone 27) (SEQ ID NO: 48).

Table 4

SEQ ID NO.	Pool No.	Nucleotide Sequence (Degenerate substitutions are shown in parentheses)
3	3407	G.AGA.GAC.GA(GA).GA(GA).CC(AT).GAG.GAG.CC
4	3408	G.AGA.GAC.GA(GA).GA(GA).CC(AT).GAA.GAG.CC
5	3409	G.AGA.GAC.GA(GA).GA(GA).CC(AT).GAA.GAA.CC
6	3410	G.AGA.GAC.GA(GA).GA(GA).CC(AT).GAG.GAA.CC
7	3411	AGA.GAC.GA(GA).GA(GA).CC(CG).GAG.GAG.CC
8	3412	AGA.GAC.GA(GA).GA(GA).CC(CG).GAA.GAG.CC
9	3413	AGA.GAC.GA(GA).GA(GA).CC(CG).GAA.GAA.CC
10	3414	AGA.GAC.GA(GA).GA(GA).CC(CG).GAG.GAA.CC

11	3415	CG.TCA.CAG.(GA)TT.(GA)TC.AAC.CAT.CTC
12	3416	CG.TCA.CAG.(GA)TT.(GA)TC.TAC.CAT.CTC
13	3417	CG.TCA.CAG.(GA)TT.(GA)TC.CAC.CAT.CTC
14	3418	CG.TCA.CAG.(GA)TT.(GA)TC.GAC.CAT.CTC
15	3419	CG.TCA.CAG.(GA)TT.(GA)TC.AAC.CAT.TTC
16	3420	CG.TCA.CAG.(GA)TT.(GA)TC.TAC.CAT.TTC
17	3421	CG.TCA.CAG.(GA)TT.(GA)TC.CAC.CAT.TTC
18	3422	CG.TCA.CAG.(GA)TT.(GA)TC.GAC.CAT.TTC
19	3458	GAG GGG CAG CTT TGT GGA GA
20	3468	CAG.CAT.AGG.CCA.GCC.CCA.GGA.TGC.CT
21	3469	GTG.ATG.GCA.GCA.ATG.TTG.GCA.CGC

Example 2

Screening of human fetal brain cDNA library

The Origene human fetal brain Rapid-Screen™ cDNA Library Panel is provided as a 96-well format array consisting of 5000 clones (plasmid DNA) per well from a human fetal brain library. Subplates are available for each well consisting of 96 wells of 50 clones each in *E. coli*. This is an oligo-dT primed library, size-selected and unidirectionally inserted into the vector pCMV-XL3.

- 94 wells from the master plate were screened using PCR. The Reaction 1
- 10 Conditions described in Example 1, above, were followed, using only primers 3407 (SEQ ID NO: 3) and 3416 (SEQ ID NO: 12) with 30ng of plasmid DNA from each well. Two pools showed the positive 70bp band. The same primers and conditions were used to screen 1µl *E. coli* from each well of one of the subplates. *E. coli* from the single positive well was then plated onto LB/amp plates and single colonies screened using the same PCR
- 15 conditions. The positive clone, about 1Kb in size, was labeled 9C7E.35. It contained the original peptide sequence as well as 5' sequence that included a methionine. The 3' sequence did not contain a stop codon, suggesting that this was not a full-length clone, consistent with Northern blot data.

Example 3

PCR Cloning Methods

3'RACE was used in experiments carried out in support of the present invention to elucidate the polynucleotide encoding human β -secretase. Methods and conditions appropriate for replicating the experiments described herein and/or determining polynucleotide sequences encoding additional members of the novel family of aspartyl proteases described herein may be found, for example, in White, B.A., ed., PCR Cloning Protocols; Humana Press, Totowa, NJ, 1997, or Ausubel, *supra*, both of which are incorporated herein by reference.

RT-PCR

For reverse transcription polymerase chain reaction (RT-PCR), two partially degenerate primer sets used for RT-PCR amplification of a cDNA fragment encoding this peptide. Primer set 1 consisted of DNA's #3427-3434 (SEQ ID NOS: 22-29 respectively), the sequences of which are shown in Table 5, below. Matrix RT-PCR using combinations of primers from this set with cDNA reverse transcribed from primary human neuronal cultures as template yielded the predicted 54 bp cDNA product with primers #3428 + 3433 (SEQ ID NOS: 23 + 28). All RT-PCR reactions employed 10-50 ng input poly-A+ RNA equivalents per reaction, and were carried out for 35 cycles employing step cycle conditions with a 95°C denaturation for 1 minute, 50°C annealing for 30 sec, and a 72°C extension for 30 sec.

The degeneracy of primers #3428 + 3433 (SEQ ID NOS: 23 + 28) was further broken down, resulting in primer set 2, comprising DNAs #3448-3455 (SEQ ID NOS: 30-37) (Table 5). Matrix RT-PCR was repeated using primer set 2, and cDNA reverse transcribed from poly-A+ RNA from IMR-32 human neuroblastoma cells (American Type Culture Collection, Manassas, VA), as well as primary human neuronal cultures, as template for amplification. Primers #3450 (SEQ ID NO: 32) and 3454 (SEQ ID NO: 36) from set 2 most efficiently amplified a cDNA fragment of the predicted size (72 bp), although primers 3450+3453 (SEQ ID NOS: 32 and 35), and 3450+3455 (SEQ ID NOS: 32 and 37) also amplified the same product, albeit at lower efficiency. A 72 bp PCR product was obtained by amplification of cDNA from IMR-32 cells and primary human neuronal cultures with primers 3450 (SEQ ID NO: 32) and 3454 (SEQ ID NO: 36).

5' and 3' RACE-PCR

Internal primers matching the upper (coding) strand for 3' Rapid Amplification of 5' Ends (RACE) PCR, and lower (non-coding) strand for 5' RACE PCR were designed and made according to methods known in the art (e.g., Frohman, M. A., M. K. Dush and G. R.

Martin (1988). "Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene specific oligo-nucleotide primer." Proc. Natl. Acad. Sci. U.S.A. 85(23): 8998-9002.) The DNA primers used for this experiment (#3459 & #3460) (SEQ ID NOS: 36 and 39) are illustrated schematically in Table 4 and the exact sequence of these primers is presented in Table 3. These primers can be utilized in standard RACE-PCR methodology employing commercially available templates (e.g. Marathon Ready cDNA®, Clontech Labs), or custom tailored cDNA templates prepared from RNAs of interest as described by Frohman *et al.* (ibid.).

In experiments carried out in support of the present invention, a variation of RACE was employed to exploit an IMR-32 cDNA library cloned in the retrovirus expression vector pLPCXlox, a derivative of pLNCX. As the vector junctions provide unique anchor sequences abutting the cDNA inserts in this library, they serve the purpose of 5' and 3' anchor primers in RACE methodology. The sequences of the specific 5' and 3' anchor primers we employed to amplify β -secretase cDNA clones from the library, primers #3475 (SEQ ID NO: 40) and #3476 (SEQ ID NO: 41), are derived from the DNA sequence of the vector provided by Clontech Labs, Inc., and are shown in Table 3.

Primers #3459 (SEQ ID NO: 38) and #3476 (SEQ ID NO: 41) were used for 3' RACE amplification of downstream sequences from our IMR-32 cDNA library in the vector pLPCXlox. The library had previously been sub-divided into 100 pools of 5,000 clones per pool, and plasmid DNA was isolated from each pool. A survey of the 100 pools with the primers identified as diagnostic for presence of the β -secretase clone, according to methods described in Example 1, above, provided individual pools from the library for RACE-PCR. 100 ng template plasmid from pool 23 was used for PCR amplification with primers 3459 + 3476 (SEQ ID NOS: 38 and 41 respectively). Amplification was carried out for 40 cycles using ampli-Taq Gold®, under the following conditions: denaturation at 95°C for 1 min, annealing at 65°C for 45 sec., and extension at 72°C for 2 min. Reaction products were fractionated by agarose gel chromatography, according to methods known in the art (Ausubel; Sambrook).

An approximately 1.8 Kb PCR fragment was revealed by agarose gel fractionation of the reaction products. The PCR product was purified from the gel and subjected to DNA sequence analysis using primer #3459 (SEQ ID NO: 38). The resulting sequence, designated 23A, and the predicted amino acid sequence deduced from the DNA sequence are shown in FIG. 5. Six of the first seven deduced amino-acids from one of the reading frames of 23A were an exact match with the last 7 amino-acids of the N-terminal sequence determined from the purified

protein, purified and sequenced in further experiments carried out in support of the present invention, from natural sources.

Table 5

SEQ ID NO.	DNA #	NUCLEOTIDE SEQUENCE	COMMENTS
22	3427	GAY GAR GAG CCN GAG GA	
23	3428	GAY GAR GAG CCN GAa GA	
24	3429	GAY GAR GAa CCN GAg GA	
25	3430	GAY GAR GAa CCN GAa GA	
26	3431	RTT RTC NAC CAT TTC	
27	3432	RTT RTC NAC CAT cTC	
28	3433	TCN ACC ATY TCN ACA AA	
29	3434	TCN ACC ATY TCN ACG AA	
30	3448	ata <u>ttc tag a</u> GAY GAR GAg CCa GAa GA	5' primer, break down of 3428 w/ 5' XbaI tail, 1 of 4
31	3449	ata <u>ttc tag a</u> GAY GAR GAg CCg GAa GA	5' primer, break down of 3428 w/ 5' XbaI tail, 2 of 4
32	3450	ata <u>ttc tag a</u> GAY GAR GAg CCc GAa GA	5' primer, break down of 3428 w/ 5' XbaI tail, 3 of 4
33	3451	ata <u>ttc tag a</u> GAY GAR GAg CcI GAa GA	5' primer, break down of 3428 w/ 5' XbaI tail, 4 of 4
34	3452	aca <u>cga att c</u> TT RTC NAC CAT YTC aAC AAA	breakdown of 3433, 1 of 4; tm = 50
35	3453	aca <u>cga att c</u> TT RTC NAC CAT YTC gAC AAA	breakdown of 3433 w/ 5' Eco RI tail, 2 of 4; tm = 50
36	3454	aca <u>cga att c</u> TT RTC NAC CAT YTC cAC AAA	breakdown of 3433 w/ 5' Eco RI tail, 3 of 4; tm = 50
37	3455	aca <u>cga att c</u> TT RTC NAC CAT YTC IAC AAA	breakdown of 3433 w/ 5' Eco RI tail, 4 of 4; tm = 50
38	3459	aa gaG CCC GGC CGG AGG GGC A	5' upper strand primer for 3' race encodes eEPGRRG
39	3460	aaa GCT GCC CCT CCG GCC GGG	3' lower strand primer for 5' RACE
40	3475	AGC TCG TTT AGT GAA CCG TCA GAT CG	pLNCX 5' primer
41	3476	ACC TAC AGG TGG GGT CTT TCA TTC CC	pLNCX, 3' primer

Example 4

β -secretase Inhibitor Assays

Assays for measuring β -secretase activity are well known in the art. Particularly useful assays, summarized below, are detailed in allowed U.S. Patent 5,744,346, incorporated herein by reference.

A. Preparation of MBP-C125sw

1. Preparation of cells

Two 250 ml cell culture flasks containing 50 ml LBamp100 per flask were seeded with one colony per flask of E. coli pMAL-C125SW cl. 2 (E. coli expressing MBP-C125sw fusion protein). Cells were allowed to grow overnight at 37°C. Aliquots (25 ml) were seeded in 500 ml per flask of LBamp100 in 2 liter flasks, which were then allowed to grow at 30°. Optical densities were measured at 600 nm (OD_{600}) vs LB broth; 1.5 ml 100mM IPTG was added when the OD was ~0.5. At this point, a pre-incubation aliquot was removed for SDS-PAGE ("I"). Of this aliquot, 0.5 ml was centrifuged for 1 min in a Beckman microfuge, and the resulting pellet was dissolved in 0.5 ml 1 x LSB. The cells were incubated/induced for 5-6 hours at 30 C, after which a post-incubation aliquot ("I") was removed. Cells were then centrifuged at 9,000 rpm in a KA9.1 rotor for 10 min at 4° C. Pellets were retained and stored at -20 C.

2. Extraction of bacterial cell pellets

Frozen cell pellets were resuspended in 50 ml 0.2 M NaCl, 50mM Tris, pH 7.5, then sonicated in rosette vessal for 5 x 20 sec bursts, with 1min rests between bursts. The extract was centrifuged at 16,500 rpm in a KA18.5 rotor 30 min (39,000 x g). Using pipette as a pestle, the sonicated pellet was suspended in 50 ml urea extraction buffer (7.6 M urea, 50 mM Tris pH 7.5, 1 mM EDTA, 0.5% TX-100). The total volume was about 25 ml per flask. The suspension was then sonicated 6x 20 sec, with 1 min rests between bursts. The suspension was then centrifuged again at 16,500 rpm 30 min in the KA18.5 rotor. The resulting supernatant was added to 1.5 L of buffer consisting of 0.2 M NaCl 50 mM Tris buffer, pH 7.5, with 1% Triton X-100 (0.2M NaCl-Tris-1%Tx), and was stirred gently at 4 degrees C for 1 hour, followed by centrifugation at 9,000 rpm in KA9.1 for 30 min at 4°C. The supernatant was loaded onto a column of washed amylose (100 ml of 50% slurry; New England BioLabs). The column was washed with 0.2 M NaCl-Tris-1%TX to baseline (+10 column volumes), then with 2 column volumes 0.2M NaCl-Tris-1% reduced Triton X-100.

The protein was then eluted with 10 mM maltose in the same buffer. An equal volume of 6 M guanidine HCl/0.5% TX-100 was added to each fraction. Peak fractions were pooled and diluted to a final concentration of about 2 mg/ml. The fractions were stored at -40 degrees C, before dilution (20-fold, to 0.1 mg/ml in 0.15% Triton X-100). Diluted aliquots were also stored at -40 C.

B. Antibody-based Assays

The assays described in this section are based on the ability of certain antibodies, hereinafter "cleavage-site antibodies," to distinguish cleavage of APP by β -secretase, based on the unique cleavage site and consequent exposure of a specific C-terminus formed by the cleavage. The recognized sequence is a sequence of usually about 3-5 residues is immediately amino terminal of the β amyloid peptide (β AP) produced by β -secretase cleavage of β -APP, such as Val-Lys-Met in wild-type or Val-Asn-Leu- in the Swedish double mutation variant form of APP. Recombinantly-expressed proteins, described below, were used as substrates for β -secretase.

MBP-C125 Assay: MBP-C125 substrates were expressed in *E. coli* as a fusion protein of the last 125 amino acids of APP fused to the carboxy-terminal end of maltose-binding protein (MBP), using commercially available vectors from New England Biolabs. The β -cleavage site was thus 26 amino acids downstream of the start of the C-125 region. This latter site is recognized by monoclonal antibody SW192.

Recombinant proteins were generated with both the wild-type APP sequence (MBP-C125 wt) at the cleavage site (..Val-Lys-Met-Asp-Ala..) (SEQ ID NO: 54) or the "Swedish" double mutation (MBP-C125.sw) (..Val-Asn-Leu-Asp-Ala..) (SEQ ID NO: 51). As shown schematically in FIG. 19A, cleavage of the intact MBP-fusion protein results in the generation of a truncated amino-terminal fragment, with the new SW-192 Ab-positive epitope uncovered at the carboxy terminus. This amino-terminal fragment can be recognized on Western blots with the same Ab, or, quantitatively, using an anti-MBP capture-biotinylated SW-192 reporter sandwich format, as shown in FIG. 19A. Anti-MBP polyclonal antibodies were raised in rabbits (Josman Labs, Berkeley) by immunization with purified recombinantly expressed MBP (New England Biolabs). Antisera were affinity purified on a column of immobilized MBP. MBP-C125 SW and WT substrates were expressed in *E. coli*, then purified as described above.

Microtiter 96-well plates were coated with purified anti-MBP antibody (at a concentration of 5-10 μ g/ml), followed by blocking with 2.5g/liter human serum albumin in

1 g/liter sodium phosphate monobasic, 10.8 g/liter sodium phosphate dibasic, 25 g/liter sucrose, 0.5 g/liter sodium azide, pH 7.4. Appropriately diluted β -secretase enzyme (5 μ l) was mixed with 2.5 μ l of 2.2 μ M MBP-C125sw substrate stock, in a 50 μ l reaction mixture with a final buffer concentration of 20 mM acetate buffer, pH 4.8, 0.06% Triton X-100, in individual wells of a 96-well microtiter plate, and incubated for 1 hour at 37 degrees C. Samples were then diluted 5-fold with Specimen Diluent (0.2 g/l sodium phosphate monobasic, 2.15 g/l sodium phosphate dibasic, 0.5 g/l sodium azide, 8.5 g/l sodium chloride, 0.05% Triton X-405, 6 g/l BSA), further diluted 5-10 fold into Specimen Diluent on anti-MBP coated plates, and incubated for 2 hours at room temperature. Following incubations with samples or antibodies, plates were washed at least four times in TTBS (0.15 M NaCl, 50 mM Tris, pH 7.5, 0.05% Tween-20). Biotinylated SW192 antibodies were used as the reporter. SW192 polyclonal antibodies were biotinylated using NHS-biotin (Pierce), following the manufacturer's instruction. Usually, the biotinylated antibodies were used at about 240 ng/ml, the exact concentration varying with the lot of antibodies used. Following incubation of the plates with the reporter, the ELISA was developed using streptavidin-labeled alkaline phosphatase (Boehringer-Mannheim) and 4-methyl-umbelliferyl phosphate as fluorescent substrate. Plates were read in a Cytofluor 2350-Fluorescent Measurement System. Recombinantly generated MBP-26SW (product analog) was used as a standard to generate a standard curve, which allowed the conversion of fluorescent units into amount of product generated.

This assay protocol was used to screen for inhibitor structures, using "libraries" of compounds assembled onto 96-well microtiter plates. Compounds were added, in a final concentration of 20 μ g/ml in 2% DMSO, in the assay format described above, and the extent of product generated compared with control (2% DMSO only) β -secretase incubations, to calculate "% inhibition." "Hits" were defined as compounds which result in >35% inhibition of enzyme activity at test concentration. This assay can also be used to provide IC_{50} values for inhibitors, by varying the concentration of test compound over a range to calculate from a dose-response curve the concentration required to inhibit the activity of the enzyme by 50%.

Generally, inhibition is considered significant as compared to control activity in this assay if it results in activity that is at least 1 standard deviation, and preferably 2 standard deviations lower than a mean activity value determined over a range of samples. In addition

a reduction of activity that is greater than about 25%, and preferably greater than about 35% of control activity may also be considered significant.

Using the foregoing assay system, 24 "hits" were identified (>30% inhibition at 50 μ M concentration) from the first 6336 compounds tested (0.4% hit rate). Of these 12 compounds had IC_{50} s less than 50 μ M, including re-screening in the P26-P4'sw assay, below.

P26-P4'sw assay. The P26-P4'sw substrate is a biotin-linked peptide of the sequence (biotin)CGGADRGLTTRPGSGLTNIKTEEISEVNLD AEF (SEQ ID NO: 63). The P26-P1 standard has the sequence (biotin)CGGADRGLTTRPGSGLTNIKTEEISEVN L (SEQ ID NO: 64), where the N-terminal "CGG" serves as a linker between biotin and the substrate in both cases. Peptides were prepared by Anaspec, Inc. (San Jose, CA) using solid phase synthesis with boc-amino acids. Biotin was coupled to the terminal cysteine sulfhydryl by Anaspec, Inc. after synthesis of the peptide, using EZ-link Iodoacetyl-LC-Biotin (Pierce). Peptides are stored as 0.8-1.0 mM stocks in 5 mM Tris, with the pH adjusted to around neutral (pH 6.5-7.5) with sodium hydroxide.

For the enzyme assay, the substrate concentration can vary from 0 – 200 μ M. Specifically for testing compounds for inhibitory activity, substrate concentration is 1.0 μ M. Compounds to be tested were added in DMSO, with a final DMSO concentration of 5%; in such experiments, the controls also receive 5% DMSO. Concentration of enzyme was varied, to give product concentrations within the linear range of the ELISA assay (125 – 2000 pM, after dilution). These components were incubated in 20 mM sodium acetate, pH 4.5, 0.06% Triton X-100, at 37 °C for 1 to 3 hours. Samples were diluted 5-fold in specimen diluent (145.4 mM sodium chloride, 9.51 mM sodium phosphate, 7.7 mM sodium azide, 0.05% Triton X-405, 6 gm/liter bovine serum albumin, pH 7.4) to quench the reaction, then diluted further for the ELISA as needed. For the ELISA, Costar High Binding 96-well assay plates (Corning, Inc., Corning, NY) were coated with SW 192 monoclonal antibody from clone 16A7, or a clone of similar affinity. Biotin-P26-P4' standards were diluted in specimen diluent to a final concentration of 0 to 2 nM. Diluted samples and standards (100 μ l) are incubated on the SW192 plates at 4 °C for 24 hours. The plates are washed 4 times in TTBS buffer (150 mM sodium chloride, 25 mM Tris, 0.05 % Tween 20, pH 7.5), then incubated with 0.1 ml/well of streptavidin – alkaline phosphatase (Roche Molecular Biochemicals, Indianapolis, IN) diluted 1:3000 in specimen diluent. After incubating for one hour at room temperature, the plate was washed 4 times in TTBS, as described in the previous section, and

incubated with fluorescent substrate solution A (31.2 gm/liter 2-amino-2-methyl-1-propanol, 30 mg/liter, adjusted to pH 9.5 with HCl). Fluorescent values were read after 30 minutes.

C. Assays using Synthetic Oligopeptide Substrates

5 This assay format is particularly useful for measuring activity of partially purified β -secretase preparations. Synthetic oligopeptides are prepared which incorporate the known cleavage site of β -secretase, and optional detectable tags, such as fluorescent or chromogenic moieties. Examples of such peptides, as well as their production and detection methods are described in allowed U.S. Patent 5,942,400, herein incorporated by reference. Cleavage
10 products can be detected using high performance liquid chromatography, or fluorescent or chromogenic detection methods appropriate to the peptide to be detected, according to methods well known in the art. By way of example, one such peptide has the sequence SEVNL DAEF (SEQ ID NO: 52), and the cleavage site is between residues 5 and 6. Another preferred substrate has the sequence ADRGLTTRPGSGLTNIKTEEISEVNLDAE F
15 (SEQ ID NO: 53), and the cleavage site is between residues 26 and 27.

D. β -secretase Assays of Crude Cell or Tissue Extracts

Cells or tissues were extracted in extraction buffer (20 mM HEPES, pH 7.5, 2 mM
20 EDTA, 0.2% Triton X-100, 1 mM PMSF, 20 μ g/ml pepstatin, 10 μ g/ml E-64). The volume of extraction buffer will vary between samples, but should be at least 200 μ l per 10^6 cells. Cells can be suspended by trituration with a micropipette, while tissue may require homogenization. The suspended samples were incubated for 30 minutes on ice. If necessary to allow pipetting, unsolubilized material was removed by centrifugation at 4 degrees C,
25 16,000 x g (14,000 rpm in a Beckman microfuge) for 30 minutes. The supernate was assayed by dilution into the final assay solution. The dilution of extract will vary, but should be sufficient so that the protein concentration in the assay is not greater than 60 μ g/ml. The assay reaction also contained 20 mM sodium acetate, pH 4.8, and 0.06% Triton X-100 (including Triton contributed by the extract and substrate), and 220 – 110 nM MBP-C125 (a 1:10 or
30 1:20 dilution of the 0.1 mg/ml stock described in the protocol for substrate preparation). Reactions were incubated for 1 – 3 hours at 37 degrees C before quenching with at least 5 – fold dilution in specimen diluent and assaying using the standard protocol.

Example 5
Purification of β -secretase

A. Purification of Naturally Occurring β -secretase

5 Human 293 cells were obtained and processed as described in U.S. Patent 5,744,346, incorporated herein by reference. (293 cells are available from the American Type Culture Collection, Manassas, VA). Frozen tissue (293 cell paste or human brain) was cut into pieces and combined with five volumes of homogenization buffer (20 mM Hepes, pH 7.5, 0.25 M sucrose, 2 mM EDTA). The suspension was homogenized using a blender and centrifuged at
10 16,000 x g for 30 min at 4°C. The supernatants were discarded and the pellets were suspended in extraction buffer (20 mM MES, pH 6.0, 0.5% Triton X-100, 150 mM NaCl, 2 mM EDTA, 5 μ g/ml leupeptin, 5 μ g/ml E64, 1 μ g/ml pepstatin, 0.2 mM PMSF) at the original volume. After vortex-mixing, the extraction was completed by agitating the tubes at 4°C for a period of one hour. The mixtures were centrifuged as above at 16,000 x g, and the
15 supernatants were pooled. The pH of the extract was adjusted to 7.5 by adding ~1% (v/v) of 1 M Tris base (not neutralized).

The neutralized extract was loaded onto a wheat germ agglutinin-agarose (WGA-agarose) column pre-equilibrated with 10 column volumes of 20 mM Tris, pH 7.5, 0.5% Triton X-100, 150 mM NaCl, 2 mM EDTA, at 4°C. One milliliter of the agarose resin was
20 used for every 1 g of original tissue used. The WGA-column was washed with 1 column volume of the equilibration buffer, then 10 volumes of 20 mM Tris, pH 7.5, 100 mM NaCl, 2 mM NaCl, 2 mM EDTA, 0.2% Triton X-100 and then eluted as follows. Three-quarter column volumes of 10% chitin hydrolysate in 20 mM Tris, pH 7.5, 0.5%, 150 mM NaCl, 0.5% Triton X-100, 2 mM EDTA were passed through the column after which the flow was
25 stopped for fifteen minutes. An additional five column volumes of 10% chitin hydrolysate solution were then used to elute the column. All of the above eluates were combined (pooled WGA-eluate).

The pooled WGA-eluate was diluted 1:4 with 20 mM NaOAc, pH 5.0, 0.5% Triton X-100, 2 mM EDTA. The pH of the diluted solution was adjusted to 5.0 by adding a few
30 drops of glacial acetic acid while monitoring the pH. This "SP load" was passed through a 5-ml Pharmacia HiTrap SP-column equilibrated with 20 mM NaOAc, pH 5.0, 0.5% Triton X-100, 2 mM EDTA, at 4 ml/min at 4°C.

C

The foregoing methods provided peak activity having a specific activity of greater than 253 nM product/ml/h/ μ g protein in the MBP-C125-SW assay, where specific activity is determined as described below, with about 1500-fold purification of the protein. Specific activity of the purified β -secretase was measured as follows. MBP C125-SW substrate was combined at approximately 220 nM in 20 mM sodium acetate, pH 4.8, with 0.06% Triton X-100. The amount of product generated was measured by the β -secretase assay, also described below. Specific activity was then calculated as:

$$\text{Specific Activity} = \frac{(\text{Product conc. nM})(\text{Dilution factor})}{(\text{Enzyme sol. vol})(\text{Incub. time h})(\text{Enzyme conc. mg/vol})}$$

The Specific Activity is thus expressed as pmoles of product produced per μ g of β -secretase per hour. Further purification of human brain enzyme was achieved by loading the SP flow through fraction on to the P10-P4'sta D->V affinity column, according to the general methods described below. Results of this purification step are summarized in Table 1, above.

B. Purification of β -secretase from Recombinant Cells

Recombinant cells produced by the methods described herein generally were made to over-express the enzyme; that is, they produced dramatically more enzyme per cell than is found to be endogenously produced by the cells or by most tissues. It was found that some of the steps described above could be omitted from the preparation of purified enzyme under these circumstances, with the result that even higher levels of purification were achieved.

CosA2 or 293 T cells transfected with β -secretase gene construct (see Example 6) were pelleted, frozen and stored at -80 degrees until use. The cell pellet was resuspended by homogenizing for 30 seconds using a handheld homogenizer (0.5 ml/pellet of approximately 10^6 cells in extraction buffer consisting of 20 mM TRIS buffer, pH 7.5, 2 mM EDTA, 0.2% Triton X-100, plus protease inhibitors: 5 μ g/ml E-64, 10 μ g/ml pepstatin, 1 mM PMSF), centrifuged as maximum speed in a microfuge (40 minutes at 4 degrees C). Pellets were suspended in original volume of extraction buffer, then stirred at 1 hour at 4 degrees C with rotation, and centrifuged again in a microfuge at maximum speed for 40 minutes. The resulting supernatant was saved as the "extract." The extract was then diluted with 20 mM sodium acetate, pH 5.0, 2 mM EDTA and 0.2% Triton X-100 (SP buffer A), and 5M NaCl was added to a final concentration of 60 mM NaCl. The pH of the solution was then adjusted

()

to pH 5.0 with glacial acetic acid diluted 1:10 in water. Aliquots were saved ("SP load"). The SP load was passed through a 1 ml SP HiTrap column which was pre-washed with 5 ml SP buffer A, 5 ml SP buffer B (SP buffer A with 1 M NaCl) and 10 ml SP buffer A. An additional 2 ml of 5% SP buffer B was passed through the column to displace any remaining sample from the column. The pH of the SP flow-through was adjusted to pH 4.5 with 10X diluted acetic acid. This flow-through was then applied to a P10-P4'staD->V-Sepharose Affinity column, as described below. The column (250 μ l bed size) was pre-equilibrated with at least 20 column volumes of equilibration buffer (25 mM NaCl, 0.2% Triton X-100, 0.1 mM EDTA, 25 mM sodium acetate, pH 4.5), then loaded with the diluted supernatant. After loading, subsequent steps were carried out at room temperature. The column was washed with washing buffer (125 mM NaCl, 0.2% Triton X-100, 25 mM sodium acetate, pH 4.5) before addition of 0.6 column bed volumes of borate elution buffer (200 mM NaCl, 0.2% reduced Triton X-100, 40 mM sodium borate, pH 9.5). The column was then capped, and an additional 0.2 ml elution buffer was added. The column was allowed to stand for 30 minutes. Two bed volumes elution buffer were added, and column fractions (250 μ l) were collected. The protein peak eluted in two fractions. 0.5 ml of 10 mg/ml peptstatin was added per milliliter of collected fractions.

Cell extracts made from cells transfected with full length clone 27 (encoding SEQ ID NO: 2; 1-501), 419stop (SEQ ID NO: 57) and 452stop (SEQ ID NO: 59) were detected by Western blot analysis using antibody 264A (polyclonal antibody directed to amino acids 46-67 of β -secretase with reference to SEQ ID NO: 2).

Example 6

Preparation of Heterologous Cells Expressing Recombinant β -secretase

25

Two separate clones (pCEKclone27 and pCEKclone53) were transfected into 293T or COS(A2) cells using Fugene and Effectene methods known in the art. 293T cells were obtained from Edge Biosystems (Gaithersburg, MD). They are KEK293 cells transfected with SV40 large antigen. COSA2 are a subclone of COS1 cells; subcloned in soft agar.

30

FuGENE Method: 293T cells were seeded at 2×10^5 cells per well of a 6 well culture plate. Following overnight growth, cells were at approximately 40-50% confluency. Media was changed a few hours before transfection (2 ml/well). For each sample, 3 μ l of FuGENE 6 Transfection Reagent (Roche Molecular Biochemicals, Indianapolis, IN) was diluted into

0.1 ml of serum-free culture medium (DME with 10 mM Hepes) and incubated at room temperature for 5 min. One microgram of DNA for each sample (0.5-2 mg/ml) was added to a separate tube. The diluted FuGENE reagent was added drop-wise to the concentrated DNA. After gentle tapping to mix, this mixture was incubated at room temperature for 15 minutes.

5 The mixture was added dropwise onto the cells and swirled gently to mix. The cells were then incubated at 37 degrees C, in an atmosphere of 7.5% CO₂. The conditioned media and cells were harvested after 48 hours. Conditioned media was collected, centrifuged and isolated from the pellet. Protease inhibitors (5 µg/ml E64, 2 µg/ml pepstatin, 0.2 mM PMSF) were added prior to freezing. The cell monolayer was rinsed once with PBS, then 0.5

10 ml of lysis buffer (1 mM HIPIS, pH 7.5, 1 mM EDTA, 0.5% Triton X-100, 1 mM PMSF, 10 µg/ml E64) was added. The lysate was frozen and thawed, vortex mixed, then centrifuged, and the supernatant was frozen until assayed.

Effective Method. DNA (0.6µg) was added with "EFFECTENE" reagent (Qiagen, Valencia, CA) into a 6-well culture plate using a standard transfection protocol according to manufacturer's instructions. Cells were harvested 3 days after transfection and the cell pellets

15 were snap frozen. Whole cell lysates were prepared and various amounts of lysate were tested for β-secretase activity using the MBP-C125sw substrate. FIG. 14B shows the results of these experiments, in which picomoles of product formed is plotted against micrograms of COS cell lysate added to the reaction. The legend to the figure describes the enzyme source, where activity from cells transfected with DNA from pCEKclone27 and

20 PCEKclone53 (clones 27 and 53) using Effective are shown as closed diamonds and solid squares, respectively, activity from cells transfected with DNA from clone 27 prepared with FuGENE are shown as open triangles, and mock transfected and control plots show no activity (closed triangles and "X" markers). Values greater than 700 pM product are out of

25 the linear range of the assay.

Example 7

Preparation of P10-P4'sta(D->V) Sepharose Affinity Matrix

30 A. Preparation of P10-P4'sta(D->V) inhibitor peptide

P10-P4'sta(D->V) has the sequence NH₂-KTEEISEVN[sta]VAEF-COOH (SEQ ID NO: 72), where "sta" represents a statine moiety. The synthetic peptide was synthesized in a peptide synthesizer using boc-protected amino acids for chain assembly. All chemicals,

reagents, and boc amino acids were purchased from Applied Biosystems (ABI; Foster City, CA) with the exception of dichloromethane and N,N-dimethylformamide which were from Burdick and Jackson. The starting resin, boc-Phe-OCH₂-Pam resin was also purchased from ABI. All amino acids were coupled following preactivation to the corresponding HOBT ester using 1.0 equivalent of 1-hydroxybenzotriazole (HOBT), and 1.0 equivalent of N,N-dicyclohexylcarbodiimide (DCC) in dimethylformamide. The boc protecting group on the amino acid α -amine was removed with 50% trifluoroacetic acid in dichloromethane after each coupling step and prior to Hydrogen Fluoride cleavage.

Amino acid side chain protection was as follows: Glu(Bzl), Lys(CI-CBZ), Ser(Obzl), Thr(Obzl). All other amino acids were used with no further side chain protection including boc-Statine.

[(Bzl) benzyl, (CBZ) carbobenzoxy, (CI-CBZ) chlorocarbobenzoxy, (Obzl) O-benzyl]

The side chain protected peptide resin was deprotected and cleaved from the resin by reacting with anhydrous hydrogen fluoride (HF) at 0°C for one hour. This generates the fully deprotected crude peptide as a C-terminal carboxylic acid.

Following HF treatment, the peptide was extracted from the resin in acetic acid and lyophilized. The crude peptide was then purified using preparative reverse phase HPLC on a Vydac C4, 330Å, 10µm column 2.2cm I.D. x 25cm in length. The solvent system used with this column was 0.1% TFA / H₂O ([A] buffer) and 0.1% TFA / CH₃CN ([B] buffer) as the mobile phase. Typically the peptide was loaded onto the column in 2% [B] at 8-10 mL/min. and eluted using a linear gradient of 2% [B] to 60% [B] in 174 minutes.

The purified peptide was subjected to mass spectrometry, and analytical reverse phase HPLC to confirm its composition and purity.

B. Incorporation into Affinity Matrix

All manipulations were carried out at room temperature. 12.5 ml of 80% slurry of NHS-Sepharose (i.e. 10 ml packed volume; Pharmacia, Piscataway, NJ) was poured into a Bio-Rad EconoColumn (BioRad, Richmond, CA) and washed with 165 ml of ice-cold 1.0 mM HCl. When the bed was fully drained, the bottom of the column was closed off, and 5.0 ml of 7.0 mg/ml P10-P4'sta (D->V) peptide (SEQ ID NO: 72) (dissolved in 0.1 M HEPES, pH 8.0) was added. The column was capped and incubated with rotation for 24 hours. After incubation, the column was allowed to drain, then washed with 8 ml of 1.0 M ethanolamine, pH 8.2. An additional 10 ml of the ethanolamine solution was added, and the column was again capped.

and incubated overnight with rotation. The column bed was washed with 20 ml of 1.5 M sodium chloride, 0.5 M Tris, pH 7.5, followed by a series of buffers containing 0.1 mM EDTA, 0.2% Triton X-100, and the following components; 20 mM sodium acetate, pH 4.5 (100 ml); 20 mM sodium acetate, pH 4.5, 1.0 M sodium chloride (100 ml); 20 mM sodium borate, pH 9.5, 1.0 M sodium chloride (200 ml); 20 mM sodium borate, pH 9.5 (100 ml). Finally, the column bed was washed with 15 ml of 2 mM Tris, 0.01% sodium azide (no Triton or EDTA), and stored in that buffer, at 4°C.

Example 8

Co-Transfection of Cells with β -secretase and APP

293T cells were co-transfected with equivalent amounts plasmids encoding APP_{sw} or wt and β -secretase or control β -galactoside (β -gal) cDNA using FuGene 6 Reagent, as described in Example 4, above. Either pCEKclone27 or pohCJ containing full length β -secretase were used for expression of β -secretase. The plasmid construct pohCK751 used for the expression of APP in these transfections was derived as described in Dugan et al., JBC, 270(18) 10982-10989(1995) and shown schematically in FIG. 21. A β -gal control plasmid was added so that the total amount of plasmid transfected was the same for each condition. β -gal expressing pCEK and pohCK vectors do not replicate in 293T or COS cells. Triplicate wells of cells were transfected with the plasmid, according to standard methods described above, then incubated for 48 hours, before collection of conditioned media and cells. Whole cell lysates were prepared and tested for the β -secretase enzymatic activity. The amount of β -secretase activity expressed by transfected 293T cells was comparable to or higher than that expressed by CosA2 cells used in the single transfection studies. Western blot assays were carried out on conditioned media and cell lysates, using the antibody 13G8, and A β ELISAs carried out on the conditioned media to analyze the various APP cleavage products.

While the invention has been described with reference to specific methods and embodiments, it will be appreciated that various modifications and changes may be made without departing from the invention. All patent and literature references referred to herein are herein incorporated by reference.

Claims

1. A protein purified to apparent homogeneity comprising a segment of a β -secretase enzyme protein lacking the signal sequence (amino acid residues 1-21 with respect to SEQ ID NO:2) and lacking the putative pro region (amino acid residues 22-45 with respect to SEQ ID NO:2).
2. The protein of claim 1, wherein the enzyme has been purified sufficiently so that its activity in cleaving the 695-amino acid isotype of β -amyloid precursor protein (B-APP) between amino acids 596 and 597 thereof is at least 10,000-fold greater than an activity exhibited by a solubilised but unenriched membrane fraction from human 293 cells.
3. The protein of claim 1, characterised by a specific activity of at least about 0.2×10^5 nM/h/ μ g protein in an MBP-C125sw substrate assay.
4. The protein of claim 3, wherein said specific activity is at least 1.0×10^5 nM/h/ μ g protein.
5. The protein of any one of claims 1-4, wherein said protein has an N-terminal residue corresponding to a residue selected from the group consisting of residues 46, 58 and 63 with respect to SEQ ID NO: 2 and a C-terminus selected from a residue between positions 452 and 501 with respect to SEQ ID NO: 2.
6. The protein of claim 5, wherein said C-terminus is between residue positions 452 and 470 with respect to SEQ ID NO: 2.
7. The protein of any of claims 1-5, wherein said protein consists of a polypeptide having the amino acid sequence SEQ ID NO: 43 [46-501], SEQ ID NO: 70 [63-452], SEQ ID NO: 69 [63-501], SEQ ID NO: 67 [58-501], or SEQ ID NO: 68 [58-452].

8. The protein of any one of claims 1-5, wherein said protein comprises or consists of a polypeptide having the amino acid sequence SEQ ID NO: 58 [46-452].
- 5 9. A crystalline protein composition formed from the protein of any one of claims 1 and 5-8.
- 10 10. The composition of claim 9, wherein said purified protein is characterised by a binding affinity for the β -secretase inhibitor substrate P10-P4' sta D \rightarrow V which is at least 1/100 of an affinity exhibited by a protein having the amino acid sequence SEQ ID NO: 43 [46-501], when said proteins are tested for binding to said substrate under the same conditions.
- 15 11. The composition of claim 9 or claim 10 when appended to claim 1, wherein said protein has the sequence SEQ ID NO: 71 [46-419] or SEQ ID NO: 75 [63-423].
12. The composition of claim 9, 10 or 11, wherein said protein is glycosylated or deglycosylated.
- 20 13. The composition of any one of claims 9-12, further comprising a β -secretase substrate selected from the group consisting of MBP-C125wt, MBP-C125sw, APP, APPsw, and β -secretase-cleavable fragments thereof or a β -secretase inhibitor molecule.
- 25 14. The composition of claim 13, wherein said β -secretase inhibitor is a peptide having fewer than about 15 amino acids and comprises the sequence SEQ ID NO: 78 (VMXVAEF; P3-P4'X D \rightarrow V), including conservative substitutions thereof.
15. The composition of claim 13, wherein said β -secretase inhibitor has the

sequence SEQ ID NO: 72 [P10-P4'sta D→V], including conservative substitutions thereof.

16. The composition of any one of claims 13-15, wherein said β -secretase inhibitor
5 has the sequence SEQ ID NO: 81 [EVMXVAEF], wherein X is hydroxyethylene or statine.

17. The composition of claim 13, wherein said β -secretase inhibitor is
characterised by a K_i of no more than about 0.5 mM.

10

18. The composition of claim 17, wherein said β -secretase inhibitor is
characterised by a K_i of no more than about 50 μ M.

19. An isolated protein, comprising a polypeptide that (i) is fewer than about 450
15 amino acid residues in length, (ii) includes an amino acid sequence that is at least 90%
identical to SEQ ID NO: 75 [63-423] including conservative substitutions thereof, and
(iii) exhibits β -secretase activity, as evidenced by an ability to cleave a substrate
selected from the group consisting of the 695 amino acid isotype of beta amyloid
precursor protein (β APP) between amino acids 596 and 597 thereof, MBP-C125wt and
20 MBP-C125sw.

20. The protein of claim 19, wherein said polypeptide includes or has the amino
acid sequence of SEQ ID NO: 75 [63-423].

25 21. The protein of claim 19, wherein said amino acid sequence is at least 95%
identical to SEQ ID NO: 58 [46-452].

22. The protein of claim 21, wherein said polypeptide has the sequence SEQ ID
NO: 58 [46-452].

30

23. The protein of any one of claims 1-8 and 19-22, wherein said protein is expressed by a heterologous cell.
24. A composition comprising the protein of any one of claims 19 to 23 and a β -secretase substrate selected from the group consisting of MBP-C125wt, MBP-C125sw, APP, APPsw, and β -secretase-cleavable fragments thereof or a β -secretase inhibitor molecule.
25. The composition of claim 24, wherein said β -secretase-cleavable fragment is selected from the group consisting of SEVKMDAEF (P5-P4'wt), SEVNLDAEF (sw), SEVKLDAEF, SEVKFDAEF, SEVNFDAEF, SEVKMAAEF, SEVNLAEEF, SEVKLAAEF, SEVKMLAEF, SEVNLEAEF, SEVKLLAEF, SEVKFAAEF, SEVNFAAEF, SEVKFLAEF, and SEVNFLAEF.
26. The composition of claim 24, wherein said β -secretase inhibitor is a peptide having fewer than about 15 amino acids and comprises the sequence SEQ ID NO: 78 (VM[X]VAEF, where X is hydroxyethylene or statine), including conservative substitutions thereof.
27. The composition of claim 26, wherein said β -secretase inhibitor has the sequence SEQ ID NO: 81 (VM[X]VAEF, where X is hydroxyethylene or statine).
28. The composition of claim 25, wherein said β -secretase inhibitor has the sequence SEQ ID NO: 72 (P10-P4'sta D \rightarrow V), including conservative substitutions thereof.
29. The composition of any one of claims 24, 26, 27 and 28, wherein said β -secretase inhibitor has a K_i of no more than about 1 μ M.
30. The composition of any one of claims 24, 26, 27 and 28, wherein said β -

secretase inhibitor is labelled with a detectable reporter molecule.

31. An antibody raised against a purified β -secretase protein comprising a polypeptide that includes an amino acid sequence that is at least 90% identical to SEQ ID NO: 75 [63-423] including conservative substitutions thereof, wherein said antibody further lacks significant immunoreactivity with a protein having a sequence selected from the group consisting of SEQ ID NO: 2 [1-501] and SEQ ID NO: 43 [46-501].
32. The antibody of claim 31, wherein said antibody is reactive with a protein selected from the group consisting of SEQ ID NO: 67 [58-501], SEQ ID NO: 69 [63-501], SEQ ID NO: 58 [46-452], SEQ ID NO: 68 [58-452] and SEQ ID NO: 70 [63-452].
33. An isolated nucleic acid, comprising a sequence of nucleotides that encodes a β -secretase protein that is at least 95% identical to a protein selected from the group consisting of SEQ ID NO: 43 [46-501], SEQ ID NO: 58 [46-452], SEQ ID NO: 67 [58-501], SEQ ID NO: 68 [58-452], SEQ ID NO: 69 [63-501], SEQ ID NO: 70 [63-452], SEQ ID NO: 75 [63-423], and SEQ ID NO: 71 [46-419], or a complementary sequence of any of such nucleotides, and specifically excluding a nucleic acid encoding a protein having the sequence SEQ ID NO: 2 [1-501].
34. The isolated nucleic acid of claim 33, wherein said sequence of nucleotides encodes a protease having an amino acid sequence SEQ ID NO: 58 [46-452] or SEQ ID NO: 43 [46-501].
35. A expression vector, comprising the isolated nucleic acid of claim 33 or claim 34, and operably linked to said nucleic acid, regulatory sequences effective for expression of the nucleic acid in a selected host cell.

36. The expression vector of claim 35, wherein said vector is suitable for transfection of a bacterial cell.
- 5 37. A heterologous cell transfected with the vector of claim 35 or claim 36, wherein said cell expresses a biologically active β -secretase.
38. The cell of claim 37, wherein said cell is a eukaryotic or bacterial cell.
- 10 39. The cell of claim 38, wherein said eukaryotic cell is an insect or yeast cell.
40. A method of producing a recombinant β -secretase enzyme, comprising culturing a cell according to claim 37, 38 or 39 under conditions to promote growth of said cell, and subjecting an extract or cultured medium from said cell to an affinity matrix
- 15 containing a β -secretase inhibitor molecule.
41. The method of claim 40, wherein said inhibitor molecule is SEQ ID NO: 72 [P10-P4'staD \rightarrow V].
- 20 42. The method of claim 40, wherein said matrix contains an antibody characterised by an ability to bind β -secretase.
43. The method of claim 42, wherein said antibody is according to claim 31 or claim 32.
- 25 44. A heterologous cell, comprising
- (i) a nucleic acid molecule encoding an active β -secretase protein according to any one of claims 19-22 or a nucleic acid molecule encoding the active β -secretase protein of SEQ ID NO: 2 [1-501];
- 30 (ii) a nucleic acid molecule encoding a β -secretase substrate molecule selected

from the group consisting of MBP-C125wt, MBP-C125sw, APPwt, APPsw, and β -secretase cleavable fragments thereof; and

(iii) operatively linked to (i) and (ii), a regulatory sequence effective for expression of said nucleic acid molecules in said cell.

5

45. The cell of claim 44, wherein said nucleic acid encoding said β -secretase protein is heterologous to said cell.

46. The cell of claim 45, wherein said nucleic acid encoding said β -secretase substrate molecule is heterologous to said cell.

10

47. The cell of claim 43, wherein said β -secretase-cleavable fragment is selected from the group consisting of SEVKMDAEF (P5-P4' wt), SEVNLDAEF (sw), SEVKLDAEF, SEVKFDAEF, SEVNFDAEF, SEVKMAAEF, SEVNLAEEF, SEVKLAAEF, SEVKMLAEF, SEVNLLAEF, SEVKLLAEF, SEVKFAAEF, SEVNFAAEF, SEVKFLAEF, and SEVNFLAEF.

15

48. A method of screening for compounds that inhibit A β production, comprising contacting an isolated β -secretase polypeptide according to claim 19 with (i) a test compound and (ii) a β -secretase substrate, and selecting the test compound as capable of inhibiting A β production if said β -secretase polypeptide exhibits less β -secretase activity in the presence of said compound than in the absence of said compound.

20

49. The method of claim 48, wherein said β -secretase polypeptide has a sequence selected from the group consisting of SEQ ID NO: 43 [46-501] and SEQ ID NO: 58 [46-452].

25

50. The method of claim 48, wherein said β -secretase polypeptide and said substrate are produced by a cell according to claim 44.

30

SUBSTITUTE SEQUENCE LISTING

<110> Anderson, John P.
 Basi, Guriqbal
 Doane, Minh Tam
 Frigon, Normand
 John, Varghese
 Power, Michael
 Sinha, Sukanto
 Tatsuno, Gwen
 Tung, Jay
 Wang, Shuwen
 McConlogue, Lisa

<120> Beta-Secretase Enzyme Compositions and
 Methods

<130> 228-US-NEW2

<140> 09/501,708

<141> 2000-02-10

<150> 60/119,571

<151> 1999-02-10

<150> 60/139,172

<151> 1999-06-15

<160> 104

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1503

<212> DNA

<213> Homo sapiens

<400> 1

atggcccaag ccctgccttg gctcctgctg tggatgggcg cgggagtgct gcctgcccac	60
ggcaccagc acggcatccg gctgcccctg cgcagcggcc tggggggcgc ccccctgggg	120
ctgcggtgctg cccgggagac cgcagaagag cccgaggagc ccggccggag gggcagcttt	180
gtggagatgg tggacaacct gaggggcaag tcggggcagg gctactacgt ggagatgacc	240
gtgggcagcc ccccgagac gctcaacatc ctggtggata caggcagcag taactttgca	300
gtgggtgctg ccccccaccc ctctctgcat cgtactacc agaggcagct gtccagcaca	360
taccgggacc tccggaaggg tgtgtatgtg ccctacaccc agggcaagtg ggaaggggag	420
ctgggcaccg acctggttaag catcccccat ggccccaacg tcaactgtgc tgccaacatt	480
gctgccatca ctgaatcaga caagttcttc atcaacggct ccaactggga aggcattcctg	540
gggctggcct atgctgagat tgccaggcct gacgactccc tggagccttt ctttgactct	600
ctggtaaacg agaccacgt tcccaacctc ttctccctgc agctttgtgg tgctggcttc	660
cccctcaacc agtctgaagt gctggcctct gtcggaggga gcatgatcat tggaggtatc	720
gaccactcgc tgtacacagg cagtctctgg tatacaccca tccggcgagg gtggtattat	780
gaggtgatca ttgtgcgggt ggagatcaat ggacaggatc tgaaaatgga ctgcaaggag	840
tacaactatg acaagagcat tgtggacagt ggcaccacca accttcgttt gcccaagaaa	900
gtgtttgaag ctgcagtcaa atccatcaag gcagcctcct ccacggagaa gttccctgat	960
ggttttctggc taggagagca gctggtgtgc tggcaagcag gcaccacccc ttgggaacatt	1020
ttcccagtca tctactcta cctaattgggt gaggttacca accagtcctt ccgcatacc	1080
atccttcgc agcaatacct gcggccagtg gaagatgtgg ccacgtccca agacgactgt	1140
tacaagtttg ccatctcaca gtcacccacg ggcagctgta tgggagctgt tatcatggag	1200
ggcttctacg ttgtctttga tcgggcccga aaacgaattg gctttgctgt cagcgcttgc	1260
catgtgcacg atgagttcag gacggcagcg gtggaaggcc cttttgtcac cttggacatg	1320

gaagactgtg gctacaacat tccacagaca gatgagtcaa cctcatgac catagcctat 1380
gtcatggctg ccatctgcgc cctcttcagt ctgccactct gcctcatggg gtgtcagtg 1440
cgctgcctcc gctgcctgcg ccagcagcat gatgactttg ctgatgacat ctccctgctg 1500
aag 1503

<210> 2

<211> 501

<212> PRT

<213> Homo sapiens

<400> 2

Met Ala Gln Ala Leu Pro Trp Leu Leu Leu Trp Met Gly Ala Gly Val
1 5 10 15
Leu Pro Ala His Gly Thr Gln His Gly Ile Arg Leu Pro Leu Arg Ser
20 25 30
Gly Leu Gly Gly Ala Pro Leu Gly Leu Arg Leu Pro Arg Glu Thr Asp
35 40 45
Glu Glu Pro Glu Glu Pro Gly Arg Arg Gly Ser Phe Val Glu Met Val
50 55 60
Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu Met Thr
65 70 75 80
Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp Thr Gly Ser
85 90 95
Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu His Arg Tyr
100 105 110
Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val
115 120 125
Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu Gly Thr Asp
130 135 140
Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg Ala Asn Ile
145 150 155 160
Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser Asn Trp
165 170 175
Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg Pro Asp Asp
180 185 190
Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr His Val Pro
195 200 205
Asn Leu Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro Leu Asn Gln
210 215 220
Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile Gly Gly Ile
225 230 235 240
Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro Ile Arg Arg
245 250 255
Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile Asn Gly Gln
260 265 270
Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys Ser Ile Val
275 280 285
Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val Phe Glu Ala
290 295 300
Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys Phe Pro Asp
305 310 315 320
Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala Gly Thr Thr
325 330 335
Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met Gly Glu Val
340 345 350
Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln Tyr Leu Arg
355 360 365
Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr Lys Phe Ala
370 375 380
Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val Ile Met Glu
385 390 395 400

Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile Gly Phe Ala
 405 410 415
 Val Ser Ala Cys His Val His Asp Glu Phe Arg Thr Ala Ala Val Glu
 420 425 430
 Gly Pro Phe Val Thr Leu Asp Met Glu Asp Cys Gly Tyr Asn Ile Pro
 435 440 445
 Gln Thr Asp Glu Ser Thr Leu Met Thr Ile Ala Tyr Val Met Ala Ala
 450 455 460
 Ile Cys Ala Leu Phe Met Leu Pro Leu Cys Leu Met Val Cys Gln Trp
 465 470 475 480
 Arg Cys Leu Arg Cys Leu Arg Gln Gln His Asp Asp Phe Ala Asp Asp
 485 490 495
 Ile Ser Leu Leu Lys
 500

<210> 3

<211> 24

<212> DNA

<213> Homo sapiens

<400> 3

gagagacgar garccwgagg agcc

24

<210> 4

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 4

gagagacgar garccwgaag agcc

24

<210> 5

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 5

gagagacgar garccwgaag aacc

24

<210> 6

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 6

gagagacgar garccwgagg aacc

24

<210> 7

<211> 23

<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 7
agagacgarg arccsgagga gcc

23

<210> 8
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 8
agagacgarg arccsgaaga gcc

23

<210> 9
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 9
agagacgarg arccsgaaga acc

23

<210> 10
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 10
agagacgarg arccsgagga acc

23

<210> 11
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 11
cgtcacagrt trtcaacct ctc

23

<210> 12
<211> 23
<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 12

cgtcacagrt trtctacat ctc

23

<210> 13

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 13

cgtcacagrt trtccacat ctc

23

<210> 14

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 14

cgtcacagrt trtcgacat ctc

23

<210> 15

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 15

cgtcacagrt trtcaacat ttc

23

<210> 16

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 16

cgtcacagrt trtctacat ttc

23

<210> 17

<211> 23

<212> DNA

<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 17
cgtcacagrt trtcaccat ttc

23

<210> 18
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 18
cgtcacagrt trtcgacat ttc

23

<210> 19
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 19
gaggggcagc tttgtggaga

20

<210> 20
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 20
cagcataggc cagccccagg atgcct

26

<210> 21
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer derived from SEQ
ID NO: 2

<400> 21
gtgatggcag caatgttggc acgc

24

<210> 22
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer

<221> misc_feature
<222> (1)...(17)
<223> n = A,T,C or G

<400> 22
gaygargagc cngagga

<210> 23
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer

<221> misc_feature
<222> (1)...(17)
<223> n = A,T,C or G

<400> 23
gaygargagc cngaaga

<210> 24
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer

<221> misc_feature
<222> (1)...(17)
<223> n = A,T,C or G

<400> 24
gaygargaac cngagga

<210> 25
<211> 17
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer

<221> misc_feature
<222> (1)...(17)
<223> n = A,T,C or G

<400> 25
gaygargaac cngaaga

<210> 26
<211> 15
<212> DNA
<213> Artificial Sequence

<220>

<223> Degenerate oligonucleotide primer

<221> misc_feature

<222> (1)...(15)

<223> n = A,T,C or G

<400> 26

rttrtcnacc atttc

15

<210> 27

<211> 15

<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerate oligonucleotide primer

<221> misc_feature

<222> (1)...(15)

<223> n = A,T,C or G

<400> 27

rttrtcnacc atctc

15

<210> 28

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerate oligonucleotide primer

<221> misc_feature

<222> (1)...(17)

<223> n = A,T,C or G

<400> 28

tcnaccatyt cnacaaa

17

<210> 29

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerate oligonucleotide primer

<221> misc_feature

<222> (1)...(17)

<223> n = A,T,C or G

<400> 29

tcnaccatyt cnacgaa

17

<210> 30

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerate oligonucleotide primer

<400> 30
 atattctaga gaygargagc cagaaga 27

 <210> 31
 <211> 27
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Degenerate oligonucleotide primer

 <400> 31
 atattctaga gaygargagc cggaaga 27

 <210> 32
 <211> 27
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Degenerate oligonucleotide primer

 <400> 32
 atattctaga gaygargagc ccgaaga 27

 <210> 33
 <211> 27
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Degenerate oligonucleotide primer

 <400> 33
 atattctaga gaygargagc ctgaaga 27

 <210> 34
 <211> 30
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Degenerate oligonucleotide primer

 <221> misc_feature
 <222> (1)...(30)
 <223> n = A,T,C or G

 <400> 34
 acacgaattc ttrtcnacca tytcaacaaa 30

 <210> 35
 <211> 30
 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Degenerate oligonucleotide primer

 <221> misc_feature

<222> (1)...(30)
<223> n = A,T,C or G

<400> 35
acacgaattc ttrtcnacca tytcgacaaa

30

<210> 36
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer

<221> misc_feature
<222> (1)...(30)
<223> n = A,T,C or G

<400> 36
acacgaattc ttrtcnacca tytccacaaa

30

<210> 37
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer

<221> misc_feature
<222> (1)...(30)
<223> n = A,T,C or G

<400> 37
acacgaattc ttrtcnacca tytctaataa

30

<210> 38
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer

<400> 38
aagagcccg cccgaggggc a

21

<210> 39
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Degenerate oligonucleotide primer

<400> 39
aaagctgccc ctccggccgg g

21

<210> 40
<211> 26
<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerate oligonucleotide primer

<400> 40

agctcgttta gtgaaccgtc agatcg

26

<210> 41

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Degenerate oligonucleotide primer

<400> 41

acctacaggt ggggtctttc attccc

26

<210> 42

<211> 2348

<212> DNA

<213> Homo sapiens

<400> 42

ccatgccggc	ccctcacagc	cccgcggga	gcccagagccc	gctgcccagg	ctggccgccc	60
csgtgccgat	gtagcgggct	ccggatccca	gcctctcccc	tgctcccgtg	ctctgccgat	120
ctcccctgac	cgctctccac	agccccgacc	cgggggctgg	cccagggccc	tgcaggccct	180
ggcgctcctga	tccccccaag	ctccctctcc	tgagaagcca	ccagcaccac	ccagacttgg	240
gggcaggcgc	cagggacgga	cgtgggccag	tgcgagccca	gagggcccga	aggccggggc	300
ccaccatggc	ccaagccctg	ccctggctcc	tgctgtggat	gggcgcggga	gtgctgcctg	360
cccacggcac	ccagcacggc	atccggctgc	ccctgcgcag	cggcctgggg	ggcgccccc	420
tggggctgcg	gctgcccccg	gagaccgacg	aagagcccga	ggagcccggc	cggaggggca	480
gctttgtgga	gatggtggac	aacctgaggg	gcaagtcggg	gcagggctac	tacgtggaga	540
tgaccgtggg	cagccccccg	cagacgctca	acatcctggg	ggatacaggc	agcagtaact	600
ttgcagtggg	tgctgcccc	cacccttcc	tgcâtcgcta	ctaccagagg	cagctgtcca	660
gcacataccg	ggacctccgg	aagggtgtgt	atgtgcccta	caccagggc	aagtgggaag	720
gggagctggg	caccgacctg	gtaagcatcc	cccatggccc	caacgtcact	gtgcgtgcca	780
acattgtctg	catcactgaa	tcagacaagt	tcttcatcaa	cggctccaac	tgggaaggca	840
tcctggggct	ggcctatgct	gagattgcca	ggcctgacga	ctccctggag	cctttctttg	900
actctctggt	aaagcagacc	cacgttccca	acctcttctc	cctgcagctt	tgtggtgctg	960
gcttccccct	caaccagtct	gaagtctgg	cctctgtcgg	agggagcatg	atcattggag	1020
gtatcgacca	ctcgctgtac	acaggcagtc	tctggtatac	acccatccgg	cgggagtggg	1080
attatgaggt	gatcattgtg	cgggtggaga	tcaatggaca	ggatctgaaa	atggactgca	1140
aggagtacaa	ctatgacaag	agcattgtgg	acagtggcac	caccaacctt	cgtttgccca	1200
âgaaagtgtt	tgaagctgca	gtcaaatcca	tcaaggcagc	ctcctccacg	gagaagtccc	1260
ctgatggttt	ctggctagga	gagcagctgg	tgtgctggca	agcaggcacc	accccttgga	1320
acattttccc	agtcattctca	ctctacctaa	tgggtgaggt	taccaaccag	tccttcgcga	1380
tcaccatcct	tcgcagcaa	tacctgcggc	cagtgggaaga	tgtggccacg	tcccaagacg	1440
actgttacaa	gtttgccatc	tcacagtcac	ccacgggcac	tgttatggga	gctgttatca	1500
tggagggcct	ctacgttgtc	tttgatcggg	cccgaâaacg	aattggcttt	gctgtcagcg	1560
cttgccatgt	gcacgatgag	ttcaggacgg	cagcggtgga	aggccctttt	gtcaccttgg	1620
acatggaaga	ctgtggctac	aacattccac	agacagatga	gtcaaccctc	atgaccatag	1680
cctatgtcat	ggctgccatc	tgcgccctct	tcâtgtgtcc	actctgcctc	atgggtgtgc	1740
agtggcgctg	cctccgctgc	ctgcgccagc	agcatgatga	ctttgctgat	gacatctccc	1800
tgctgaagtg	aggaggccca	tgggcagaag	atagagattc	ccctggacca	cacctccgtg	1860
gttcactttg	gtcacaagta	ggagacacag	atggcacctg	tggccagagc	acctcaggac	1920
cctcccacc	caccaaâtgc	ctctgccttg	atggagaagg	aaaaggctgg	caaggtgggt	1980
tccagggact	gtacctgtag	gaaacagaaa	agagaagaaa	gaagcactct	gctggcggga	2040
atactcttgg	tcacctcaaa	tttaagtctg	gaaattctgc	tgcttgaâac	ttcagccctg	2100
aacctttgtc	caccattcct	ttaaattctc	caacccaaag	tattcttctt	ttcttagttt	2160

cagaagtact ggcatacacac gcaggttacc ttggcgtgtg tccctgtggt accctggcag 2220
 agaagagacc aagcttggtt ccctgctggc caaagtcagt aggagaggat gcacagtttg 2280
 ctatttgctt tagagacagg gactgtataa acaagcctaa cattggtgca aagattgcct 2340
 cttgaatt 2348

<210> 43
 <211> 456
 <212> PRT
 <213> Homo sapiens

<400> 43
 Glu Thr Asp Glu Glu Pro Glu Glu Pro Gly Arg Arg Gly Ser Phe Val
 1 5 10 15
 Glu Met Val Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val
 20 25 30
 Glu Met Thr Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp
 35 40 45
 Thr Gly Ser Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu
 50 55 60
 His Arg Tyr Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg
 65 70 75 80
 Lys Gly Val Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu
 85 90 95
 Gly Thr Asp Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg
 100 105 110
 Ala Asn Ile Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly
 115 120 125
 Ser Asn Trp Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg
 130 135 140
 Pro Asp Asp Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr
 145 150 155 160
 His Val Pro Asn Leu Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro
 165 170 175
 Leu Asn Gln Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile
 180 185 190
 Gly Gly Ile Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro
 195 200 205
 Ile Arg Arg Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile
 210 215 220
 Asn Gly Gln Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys
 225 230 235 240
 Ser Ile Val Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val
 245 250 255
 Phe Glu Ala Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys
 260 265 270
 Phe Pro Asp Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala
 275 280 285
 Gly Thr Thr Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met
 290 295 300
 Gly Glu Val Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln
 305 310 315 320
 Tyr Leu Arg Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr
 325 330 335
 Lys Phe Ala Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val
 340 345 350
 Ile Met Glu Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile
 355 360 365
 Gly Phe Ala Val Ser Ala Cys His Val His Asp Glu Phe Arg Thr Ala
 370 375 380
 Ala Val Glu Gly Pro Phe Val Thr Leu Asp Met Glu Asp Cys Gly Tyr
 385 390 395 400

Asn Ile Pro Gln Thr Asp Glu Ser Thr Leu Met Thr Ile Ala Tyr Val
 405 410 415
 Met Ala Ala Ile Cys Ala Leu Phe Met Leu Pro Leu Cys Leu Met Val
 420 425 430
 Cys Gln Trp Arg Cys Leu Arg Cys Leu Arg Gln Gln His Asp Asp Phe
 435 440 445
 Ala Asp Asp Ile Ser Leu Leu Lys
 450 455

<210> 44
 <211> 2348
 <212> DNA
 <213> Homo sapiens

<400> 44
 ccatgccggc ccctcacagc ccgcgcggga gcccgagccc gctgcccagg ctggccgccc 60
 csgtgccgat gtagcgggct ccggatccca gcctctcccc tgctcccgtg ctctgccgat 120
 ctccctgac cgctctccac agcccgagac cgggggctgg cccaggggccc tgcaggccct 180
 ggcgtcctga tgccccaag ctccctctcc tgagaagcca ccagcaccac ccagacttgg 240
 gggcaggcgc cagggacgga cgtgggcccag tgcgagccca gagggcccga aggcgggggc 300
 ccaccatggc ccaagccctg ccctggctcc tgcgtgggat gggcgcgga gtgctgcctg 360
 cccacggcac ccagcacggc atccggctgc cctgctgag cggcctgggg ggcgcccccc 420
 tggggctgcg gctgcccgg gagaccgacg aagagcccga ggagcccggc cggaggggca 480
 gctttgtgga gatggtggac aaactgaggg gcaagtccgg gcagggttac tacgtggaga 540
 tgaccgtggg cagcccccg cagacgctca acatcctggt ggatacaggc agcagtaact 600
 ttgcagtggg tgcgtcccc cacccttcc tgcctcgtca ctaccagagg cagctgtcca 660
 gcacataccg ggacctccgg aagggtgtgt atgtgcccta caccagggc aagtgggaag 720
 gggagctggg caccgacctg gtaagcatcc cccatggccc caacgtcact gtgcgtgcca 780
 acattgctgc catcactgaa tcagacaagt tcttcatcaa cggctccaac tgggaaggca 840
 tccctggggt ggcctatgct gagattgcca ggcctgacga ctccctggag cctttctttg 900
 actctctggt aaagcagacc cactgtccca acctctctc cctgcagctt tgtggtgctg 960
 gcttccccct caaccagtct gaagtgtggt cctctgtcgg agggagcatg atcattggag 1020
 gtatcgacca ctgctgtac acaggcagtc tctgtgtatc acccatccgg cgggagtggg 1080
 attatgaggt gatcattgtg cgggtggaga tcaatggaca ggatctgaaa atggactgca 1140
 aggagtacaa ctatgacaag agcattgtgg acagtggcac caccaacctt cgtttgccca 1200
 agaaaagtgt tgaagctgca gtcaaatcca tcaaggcagc ctctccacg gagaagtcc 1260
 ctgatgggtt ctggctagga gagcagctgg tgtgctggca agcaggcacc accccttggg 1320
 acattttccc agtcatctca ctctacttaa tgggtgaggt taccaaccag tccttccgca 1380
 tcaccatcct tccgcagcaa tacctgcggc cagtggaga tgtggccacg tcccaagacg 1440
 actgttacia gtttgccatc tcacagtcac ccacgggcac tgttatggga gctgttatca 1500
 tggagggtt ctacgtgtg tttgatcggg cccgaaaacg aattggcttt gctgtcagcg 1560
 cttgccatgt gcacgatgag ttccaggacgg cagcgggtga aggccctttt gtcaccttgg 1620
 acatggaaga ctgtggctac aacattccac agacagatga gtcaaccctc atgaccatag 1680
 cctatgtcat ggctgccatc tgcgcctctc tcatgtgcc actetgctc atgggtgtgtc 1740
 agtggcgctg cctccgctgc ctgcgcccag agcatgatga ctttctgat gacatctccc 1800
 tgcgaagtg aggaggccca tgggcagaag atagagattc ccctggacca cactccgtg 1860
 gttcactttg gtcacaagta ggagacacag atggcacctg tggccagagc acctcaggac 1920
 cctccccacc caccaaatgc ctctgccttg atggagaagg aaaaggctgg caaggtgggt 1980
 tccagggact gtacctgtag gaadacagaaa agagaagaaa gaagcactct gctggcggga 2040
 atactcttgg tcacctcaaa tttaagtcgg gaaattctgc tgcctgaaac ttcagccctg 2100
 aacctttgtc caccattcct ttaaattctc caaccacaaag tattcttctt tcttagttt 2160
 cagaagtact ggcacacac gcaggttacc ttggcgtgtg tccctgtggt accctggcag 2220
 agaagagacc aagcttggtt cctgctggc caaagtcagt aggagaggat gcacagtgtg 2280
 ctatttgctt tagagacagg gactgtataa acaagcctaa cattggtgca aagattgcct 2340
 cttgaatt 2348

<210> 45
 <211> 7
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Flag sequence

<400> 45
 Tyr Lys Asp Asp Asp Lys
 1 5

<210> 46
 <211> 22
 <212> PRT
 <213> Homo sapiens

<400> 46
 Met Ala Gln Ala Leu Pro Trp Leu Leu Leu Trp Met Gly Ala Gly Val
 1 5 10 15
 Leu Pro Ala His Gly Thr
 20

<210> 47
 <211> 23
 <212> PRT
 <213> Homo sapiens

<400> 47
 Gln His Gly Ile Arg Leu Pro Leu Arg Ser Gly Leu Gly Gly Ala Pro
 1 5 10 15
 Leu Gly Leu Arg Leu Pro Arg
 20

<210> 48
 <211> 16080
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Expression Vector pCEK

<221> misc_feature
 <222> (1) ... (16080)
 <223> n = A,T,C or G

<400> 48
 ttctcatgtt tgacagctta tcacgcaga tccgggcaac gttgttgcatt tgctgcaggc 60
 gcagaactgg taggtatgga agatccgatg tacgggccag atatacgcgt tgacattgat 120
 tattgactag ttattaatag taatcaatta cggggtcatt agtteatagc ccatatatgg 180
 agttccgcgt tacataactt acggtaaatg gccgcctgg ctgaccgcc aacgaccocc 240
 gccattgac gtcaataatg acgtatgttc ccatagtaac gccaataggg actttccatt 300
 gacgtcaatg ggtggactat ttacggtaaa ctgccactt ggcagtacat caagtgtatc 360
 atatgccaaag tacgcccctt attgacgtca atgacggtaa atggcccgcc tggcattatg 420
 ccagtaecat gaccttatgg gactttccta ctgggcagta catctacgta ttagtcatcg 480
 ctattaccat ggtgatgcgg ttttggcagt acatcaatgg gcgtggatag cggtttgact 540
 cacggggatt tccaagtctc caccattg acgtcaatgg gagtttgttt tggcaccaaa 600
 atcaacggga ctttccaaaa tgctgtaaca actcgcgcc attgacgcaa atgggcggta 660
 ggcgtgtacg gtgggaggtc tatataagca gagctctctg gctaactaga gaaccactg 720
 ettactggct tatcgaaatt aatacgactc actataggga gacccaagct ctgttgggct 780
 cgcggttgag gacaaactct tcgcggtctt tccagtactc ttggatcgga aaccgcgcg 840
 cctccgaacg gtactccgcc accgagggac ctgagcgagt ccgcacgcac cggatcgga 900
 aacctctcga ctgttggggt gagtactccc tctcaaaagc gggcatgact tctgcgctaa 960
 gatgtcagt ttccaaaaac gaggaggatt tgatattcac ctggcccgcg gtgatgcctt 1020
 tgagggtggc cgcgtccatc tggtcagaaa agacaatctt tttgttgtca agcttgagg 1080
 gtggcaggct tgagatctgg ccatacactt gagtgacaat gacatccact ttgcctttct 1140

ctccacaggt	gtccactccc	aggteccaact	gcaggtcgac	tctagaccgg	gggaattctg	1200
cagatatcca	tcacactggc	cgcactcgtc	cccagcccgc	ccgggagctg	cgagccgcga	1260
gctggattat	ggtggcctga	gcagccaacg	cagccgcagg	agcccgagc	ccttgccctt	1320
gcccgcgccc	ccgccgcggc	gggggaccag	ggaagccgcc	accggccgcg	catgcccgcc	1380
cctcccagcc	ccgccgggag	ccgcgcggcg	ctgcccaggc	tggccgcggc	cgtgccgatg	1440
tagcgggctc	cggatcccag	cctctcccct	gctcccgctg	tctgcggatc	ccccctgacc	1500
gctctccaca	gcccggaccc	gggggctggc	ccagggccct	gcagggccctg	gcgtccctgat	1560
gcccccaagc	tcctctctct	gagaagccac	cagcaccacc	cagacttggg	ggcaggcgcc	1620
agggacggac	gtgggcccag	gcgagcccag	agggcccga	ggccggggcc	caccatggcc	1680
caagccctgc	cctggctcct	gctgtggatg	ggcgcgggag	tgctgcctgc	ccacggcacc	1740
cagcacggca	tccggctgcc	cctgcgcagc	ggcctggggg	gcgccccctt	ggggtgcgg	1800
ctgccccggg	agaccgacga	agagcccag	gagcccgcc	ggaggggag	ccttggtggag	1860
atggtggaca	acctgagggg	caagtcgggg	cagggtact	acgtggagat	gaccgtgggc	1920
agccccccgc	agacgctcaa	catcctgggt	gatacagga	gcagtaactt	tgcagtgggt	1980
gctgcccccc	accccttctt	gcategctac	taccagaggc	agctgtccag	cacataccgg	2040
gacctccgga	agggtgtgta	tgtgccctac	acccagggca	agtgggaagg	ggagtggggc	2100
accgacctgg	taagcatccc	ccatggcccc	aacgtcactg	tgcgtgccaa	cattgctgcc	2160
atcactgaat	cagacaagtt	cttcatcaac	ggctccaact	gggaaggcat	cctggggctg	2220
gcctatgctg	agattgccag	gcctgacgac	tccttgagac	ctttctttga	ctctctggtta	2280
aagcagaccc	acgttcccaa	cctcttctcc	ctgcagcttt	gtggtgctgg	cttccccctt	2340
aaccagtctg	aagtgtctgc	ctctgtcggg	ctcagcatga	tcattggagg	tatcgaccac	2400
tcgctgtaca	caggcagttc	ctggtataca	cccatccggc	gggagtggta	ttatgaggtc	2460
atcattgtgc	gggtggagat	caatggacag	gatctgaaaa	tggactgcaa	ggagtacaac	2520
tatgacaaga	gcattgtgga	cagtggcacc	accaaccttc	gtttgcccaa	gaaagtgttt	2580
gaagctgcag	tcaaatccat	caaggcagcc	tcctccacgg	agaagttccc	tgatggtttc	2640
tggctaggag	agcagctggt	gtgctggcaa	gcaggcacca	ccccttgga	cattttccca	2700
gtcatctcac	tctacctaat	gggtgaggtt	accaaccagt	ccttcgcgat	caccatcctt	2760
ccgcagcaat	acctgcggcc	agtggaagat	gtggccacgt	cccaagacga	ctgttacaag	2820
tttgcctatc	cacagtcac	cacgggcact	gttatgggag	ctgttatcat	ggagggcttc	2880
tacgttgtct	ttgatcgggc	ccgaaaacga	attggctttg	ctgtcagcgc	ttgccatgtg	2940
cacgatgagt	tcaggacggc	agcgggtgaa	ggcccttttg	tcaccttgga	catggaagac	3000
tgtggctaca	acattccaca	gacagatgag	tcaacctca	tgaccatag	ctatgtcatg	3060
gctgccatct	gcgccctctt	catgctgcca	ctctgcctca	tgggtgtgta	gtggcgctgc	3120
ctccgctgcc	tgcgccagca	gcattgatgac	tttgcctgat	acatctccct	gctgaagtga	3180
ggagggcccat	gggcagaaga	tagagattcc	cctggaccac	acctccgtgg	ttcacttttg	3240
tcacaagtag	gagacacaga	tggcacctgt	ggccagagca	cctcaggacc	ctccccaccc	3300
accaaatgcc	tctgccttga	tggagaagga	aaaggctggc	aagggtgggt	ccagggactg	3360
tacctgtagg	aaacagaaaa	gagaagaaag	aagcactctg	ctggcgggga	tactcttggt	3420
cacctcaaat	ttaagtcggg	aaattctgct	gcttgaacct	tcagccctga	acctttgtcc	3480
accattcctt	ttaattctcc	aaccctaaag	attctctctt	tcttagtttc	agaagtactg	3540
gcacacacg	caggttacct	tggcgtgtgt	ccctgtggta	ccctggcaga	gaagagacca	3600
agcttggttc	cctgctggcc	aaagtcagta	ggagaggatg	cacagtttgc	tatttgcttt	3660
agagacaggg	actgtataaa	caagcctaac	attggtgcaa	agattgcctc	ttgaattaaa	3720
aaaaaaaaact	agattgacta	tttatacaaa	tgggggaggc	tggaaagagg	agaaggagag	3780
ggagtacaaa	gacagggaat	agtgggatca	aagctaggaa	aggcagaaac	acaaccactc	3840
accagtccta	gttttagacc	tcattctcaa	gatagcatcc	catctcagaa	gatgggtgtt	3900
gttttcaatg	ttttcttttc	tgtggttgca	gcctgaccaa	aagtggatg	ggaagggtct	3960
atctagccaa	agagctcttt	tttagctctc	ttaaagtgaag	tgccactaa	gaagttccac	4020
tttaacacatg	aatttctgcc	atattaattt	cattgtctct	atctgaacca	ccctttatct	4080
tacatatgat	aggcagcact	gaaatatcct	aaccctctaa	gctccagggtg	ccctgtggga	4140
gagcaactgg	actatagcag	ggctgggctc	tgtcttctct	gtcataggct	cactctttcc	4200
cccaaatcct	cctctggagc	tttgagccca	aggtgtctaa	aggaataggt	aggagacctc	4260
ttctatctaa	tccttaaaag	cataatgttg	aacattcatt	caacagctga	tgcctataaa	4320
ccctgectg	gatttcttcc	tattaggcta	taagaagtag	caagatcttt	acataattca	4380
gagtggtttc	attgccttcc	tacctctctt	aatggccctt	ccattttatt	gactaaagca	4440
tcacacagtg	gcactagcat	tataccaaga	gtatgagaaa	tacagtgtct	tatggctcta	4500
acattactgc	cttcagtatc	aaggctgctt	ggagaaagga	tggcagcctc	agggcttcc	4560
tatgtcctcc	accacaagag	ctccttgatg	aaggctcatc	ttttccctta	tcctgttctt	4620
ccctccccc	ctcctaattg	tacgtggggt	ccagggctgg	ttcttgggct	aggtagtggtg	4680
gaccaagttc	attacctccc	tatcagttct	agcatagtaa	actacggtac	cagtgttagt	4740
gggaagagct	gggttttctt	agtataccca	ctgcatccta	ctcctacctg	gtcaacccgc	4800

tgcctccagg	tatgggacct	gctaagtgtg	gaattacctg	ataagggaga	gggaaataca	4860
aggagggcct	ctggtgttcc	tggcctcagc	cagctgcca	caagccataa	accaataaaa	4920
caagaatact	gagtcagttt	tttatctggg	ttctcttcat	tcccactgca	cttgggtctg	4980
ctttggctga	ctgggaacac	cccataacta	cagagtctga	caggaagact	ggagactgtc	5040
cacttctagc	tgggaactta	ctgtgtaaat	aaactttcag	aactgctacc	atgaagtga	5100
aatgccacat	tttgccttat	aatttctacc	catgttggga	aaaactggct	ttttcccagc	5160
cctttccagg	gcataaaact	caacccttc	gatagcaagt	cccatcagcc	tattattttt	5220
ttaaagaaaa	cttgcaactg	tttttctttt	tacagttaact	tccttctctg	cccaaaatta	5280
taaaactctaa	gtgtaaaaaa	aagtcttaac	aacagcttct	tgcttgtaaa	aatatgtatt	5340
atacatctgt	atttttaaat	tctgtctctg	aaaaatgact	gtcccatctt	ccactcactg	5400
catttggggc	ctttcccat	ggtctgcatg	tcttttatca	ttgcaggcca	gtggacagag	5460
ggagaagggg	gaacaggggt	gcgcaacact	tgtgttgctt	tctgactgat	cctgaacaag	5520
aaagagttaac	actgaggcgc	tgcctcccat	gcacaactct	ccaaaacact	tatcctcctg	5580
caagagtggg	ctttccgggt	ctttactggg	aagcagttaa	gccccctcct	caccccttcc	5640
ttttttcttt	ctttactcct	ttggcttcaa	aggattttgg	aaaagaaaca	atatgcttta	5700
cactcatttt	caatttctaa	atttgcaggg	gatactgaaa	aatcaggcag	gtggcctaag	5760
gctgctgtaa	agttgagggg	agaggaaatc	ttaagattac	aagataaaaa	acgaatcccc	5820
taaacaaaaa	gaacaataga	actggtcttc	cattttgcca	cctttctgt	tcattgacagc	5880
tactaacctg	gagacagtaa	catttctatta	accaaagaaa	gtgggtcacc	tgacctctga	5940
agagctgagt	actcaggcca	ctccaatcac	cctacaagat	gccaaggagg	tcccagggaag	6000
tccagctcct	taaaactgacg	ctagtcaata	aacctgggca	agtgaggcaa	gagaaatgag	6060
gaagaatcca	tctgtgaggt	gacaggcacg	gatgaaagac	aaagacggaa	aagagtatca	6120
aaggcagaaa	ggagatcatt	tagttgggtc	tgaaaggaaa	agtnnttgct	atccgacatg	6180
tactgctagt	wcctgtaagc	attttaggtc	ccagaatgga	aaaaaaaatc	aagctatnng	6240
ttatataata	atgnnnnnnn	nnnnnnnnnn	nntcgagcat	gcacttagag	ggccctattc	6300
tatagtgtca	cctaaatgct	agagctcgct	gatcagcctc	gactgtgcct	tctagtgtcc	6360
agccatctgt	tgtttgccc	tccccctg	cttcttgac	cctggaagg	gccactccca	6420
ctgtcctttc	ctaataaaat	gaggaaattg	catgcgattg	tctgagtagg	tgctattcta	6480
ttctgggggg	tggggtgggg	caggacagca	agggggagga	ttggggaagc	aatagcaggc	6540
atgctgggga	tgcggtgggc	tctatggctt	ctgaggcgga	aagaaccagc	tggggctcta	6600
gggggtatcc	ccacgcgccc	tgtagcggcg	cattaagcgc	ggcgggtgtg	gtggttacgc	6660
gcagcgtgac	cgctacactt	gccagcgccc	tagecgccgc	tcctttcgct	ttcttccctt	6720
cctttctcgc	cacgttcgcc	ggctttcccc	gtcaagctct	aaatcggggc	atccctttag	6780
ggttccgatt	tagtgcttta	cggcacctcg	accccaaaaa	acttgattag	ggtgatgggt	6840
cactgagtgg	gccatcgccc	tgatagacgg	tttttcgccc	tttgacgttg	gagtcacagt	6900
tctttaatag	tggactcttg	ttccaaactg	gaacaacact	caaccctatc	tcggtctatt	6960
cttttgattt	ataagggatt	ttggggattt	cggcctattg	gttaaaaaat	gagctgattt	7020
aacaaaaatt	taacgcgaat	tctagagccc	cggccgcgga	cgaactaaac	ctgactacgg	7080
catctctgcc	ccttcttcgc	ggggcagtg	atgtaatdec	ttcagttgg	tggtacaact	7140
tgccaactgg	gccctgttcc	acatgtgaca	cgggggggga	ccaaacacaa	aggggttctc	7200
tgactgtagt	tgacatcctt	ataaatggat	gtgcacattt	gccaacactg	agtggctttc	7260
atcctggagc	agactttgca	gtctgtggac	tgcaacacaa	cattgctttt	atgtgtaact	7320
cttggtgaa	gctcttacac	caatgctggg	ggacatgtac	ctcccagggg	cccaggaaga	7380
ctacgggagg	ctacaccaac	gtcaatcaga	ggggcctgtg	tagctaccga	taagcggacc	7440
ctcaagaggg	cattagcaat	agtgtttata	aggccccctt	gttaacccta	aacgggtagc	7500
atatgcttcc	cgggtagtag	tatatactat	ccagactaac	cctaattcaa	tagcatatgt	7560
tacccaacgg	gaagcatatg	ctatcgaatt	agggttagta	aaagggtcct	aaggaaacagc	7620
gatattctcc	accccatgag	ctgtcacggt	tttatttaca	tggggtcagg	attccacgag	7680
ggtagtgaac	catttttagtc	acaaggcgag	tggctgaaga	tcaaggagcg	ggcagtgaac	7740
tctectgaat	cttcgcctgc	ttcttcatte	tecttcgttt	agctaataga	ataactgctg	7800
agttgtgaac	agtaagggtg	atgtgaggtg	ctcgaaaaca	aggtttcagg	tgacgcccc	7860
agaataaaaat	ttggacgggg	ggttcagtg	tggcattgtg	ctatgacacc	aatataaccc	7920
tcacaaaccc	cttgggcaat	aaatactagt	gtaggaatga	aacattctga	atatctttaa	7980
caatagaaat	ccatgggggtg	gggacaagcc	gtaaagactg	gatgtccatc	tcacacgaat	8040
ttatggctat	gggcaacaca	taatcctagt	gcaatatgat	actgggggtta	ttaagatgtg	8100
tcccaggcag	ggaccaagac	agggtgaacca	tgttggtaca	ctctatttgt	aacaagggga	8160
aaagagagtgg	acgcccagacg	cagcggactc	cactgggtgtg	ctctaaccac	cccgaataat	8220
aaacggggct	ccacgccaat	ggggcccata	aacaaagaca	agtggccact	cttttttttg	8280
aaattgtgga	gtgggggcac	gcgtcagccc	ccacacgccc	ccctcggtt	ttggactgta	8340
aaataagggt	gtaataactt	ggctgattgt	aaccccgcta	accactgcg	tcaaacact	8400
tgcccacaaa	accactaatg	gcaccccg	gaatacctgc	ataagtaggt	ggcggggcca	8460

agataggggc	gcgattgctg	cgatctggag	gacaaattac	acacacttgc	gcctgagcgc	8520
caagcacagg	gttggttgct	ctcatattca	cgaggctcgt	gagagcacgg	tgggctaattg	8580
ttgccatggg	tagcatatac	tacccaaata	tctggatagc	atatgctatc	ctaattctata	8640
tctgggtagc	ataggctatc	ctaattctata	tctgggtagc	atatgctatc	ctaattctata	8700
tctgggtagt	atatgctatc	ctaattctata	tctgggtagc	atatgctatc	ctaattctata	8760
tctgggtagc	atatgctatc	ctaattctata	tctgggtagt	atatgctatc	ctaattctata	8820
tccgggtagc	atatgctatc	ctaattctata	ttagggtagt	atatgctatc	ctaattctata	8880
tctgggtagc	atatactacc	caaatatctg	gatagcatat	gctatcctaa	tctatattctg	8940
ggtagcatat	gctatcctaa	tctatattctg	ggtagcatat	gctatcctaa	tctatattctg	9000
ggtagcatat	gctatcctaa	tctatattctg	ggtagcatat	gctatcctaa	tctatattctg	9060
ggtagcatat	gctatcctaa	tctatattctg	ggtagcatat	gctatcctaa	tctatattctg	9120
ggtagcatat	gctatcctaa	tctatattctg	ggtagcatat	gctatcctaa	tctatattctg	9180
gtcagcatat	gataccaggt	agtagagtg	gagtgctatc	ctttgcatat	gcegcacact	9240
cccaaggggg	cgtgaatttt	cgctgcttgt	cttttctctg	catgctggtt	gctcccatte	9300
ttagggtgaat	ttaaggaggc	caggctaaag	ccgtcgcag	tctgattgct	caccaggtaa	9360
atgtcgctaa	tgttttccaa	cgcgagaagg	tgttgagcgc	ggagctgagt	gacgtgacaa	9420
catgggtatg	cccaattgcc	ccatgttggg	aggacgaaaa	tggtgacaag	acagatggcc	9480
agaaatacac	caacagcacg	catgatgtct	gggtgggatt	tattctttag	tgcgggggaa	9540
tacacggctt	ttaatacgat	tgagggcgtc	tccaaacaag	ttacatcact	cctgcccttc	9600
ctcaccctca	tctccatcac	ctccttcac	tccgtcatct	ccgtcatcac	cctccgcggc	9660
agcccccctc	accatagggtg	gaaaccaggg	aggcaaatct	actccatcgt	caaagctgca	9720
cacagtcacc	ctgatattgc	aggtaggagc	gggctttgtc	ataacaagg	ccttaatcgc	9780
atccttcaaa	acctcagcaa	atatatgagt	ttgfaaaaag	accatgaaat	aacagacaa	9840
ggactccctt	agcggggcag	gttctggggc	gggtccagg	gccattccaa	aggggagacg	9900
actcaatggt	gtaagacgac	attgtggaat	agcaagggca	gttctctgcc	ttagggttga	9960
aagggagggtc	ttactacctc	catatacgaa	cacaccggcg	acccaagttc	cttcgtcggt	10020
agtcctttct	acgtgactcc	tagccaggag	agctcttaaa	ccttctgcaa	tgttctcaaa	10080
tttcgggttg	gaacctcctt	gaccacgatg	ctttccaaac	cacctcctt	ttttgcgcct	10140
gcctccatca	ccctgacccc	gggtccaggt	gcttgggctt	tctctgggtg	catctgcggt	10200
gccttgcctt	atcgctcccg	gggtccaggt	aggtccacca	tctggggcac	cttcttggtg	10260
gtattcaaaa	taatcggtt	ccctacaggt	gtggaaaaat	ggccttctac	ctggaggggg	10320
cctgcgcggt	ggagaccggg	atgatgatga	ctgactactg	ggactcctgg	gcctcttttc	10380
tccacgtcca	cgacctctcc	ccctggctct	ttcacgactt	ccccccctgg	ctctttcacg	10440
tctcttacct	cggtggcctc	gacctccctc	tgcacctcct	gctcctgccc	ctcctgctcc	10500
cggtccctca	ctgcctcctc	gacctccctc	ccctcctgct	cctgccccct	ctgccccctc	10560
tgccccctct	cctgctcctg	ccctcctgct	ccctcctgct	cctgccccct	ctgccccctc	10620
tgctcctgcc	cctcctgccc	ctcctgctcc	tgccccctct	gccccctcct	ctgctcctgc	10680
ccctcctgcc	cctcctcctg	ctcctgcccc	tcttgcccct	cctgctcctg	ccctcctgct	10740
ccctcctgct	cctgccccct	ctgccccctc	tgctcctgct	cctcctgctc	ctgccccctc	10800
tgctcctgcc	cctcctgctc	ctgccccctc	tgccccctct	gccccctcct	ctgctcctgc	10860
ccctcctgct	cctgccccct	ctgccccctc	tgccccctct	gccccctcct	ctgctcctgc	10920
tcttgccccct	cctgccccct	ctgccccctc	tcttgctcct	gccccctcct	ccctcctcct	10980
tgctcctgcc	cctcctcctg	ctcctgcccc	tcttgccccct	cctgccccct	ctcctgctcc	11040
tgccccctct	gccccctcct	ctgctcctgc	ccctcctcct	gctcctgccc	ctcctgcccc	11100
tcttgccccct	cctcctgctc	ctgccccctc	tcttgctcct	gccccctcct	ccctcctgct	11160
ccctcctgct	cctcctcctg	ctcctgcccc	tcttgctcct	cctgccccct	ctgctcctgc	11220
ccctcctgct	cctgctcctg	ctcctggttc	accgtgggtc	cctttgcagc	caatgcaact	11280
tggacgtttt	tgggtgtctc	ggacaccatc	tctatgtctt	ggccttgatc	ctgagccgcc	11340
cgggggtcct	ggtcttccgc	ctcctcgctc	tctgctcctt	ccctgctcct	gtccatgggt	11400
atcacccctt	cttcttttag	gtccactgct	gcccggagct	tctggtccag	atgtgtctcc	11460
cttctctcct	aggccatttc	caggctcctg	acctggcccc	tctgtagaca	tgattcacac	11520
taaaagagat	caatagacat	ctttattaga	cgacgctcag	tgaatacagg	gagtgacag	11580
tcttgccccct	tccaacagcc	ccccccacct	catccccctt	atggctcgct	tcagacagat	11640
ccagggtctga	aaattcccca	tctccgaac	catcctcgct	ctcatcacca	attactcgca	11700
gcccggaaaa	ctcccgctga	acatcctcaa	gatttgctgc	ctgagcctca	agccaggcct	11760
caaattcctc	gtcccccttt	ttgctggagc	gtagggtatg	ggattctcgg	gacccctcct	11820
cttctctctc	aaggtcacca	gacagagatg	ctactggggc	aacgggaaga	aagctgggtg	11880
cggcctgtga	ggatcagctt	atcgatgata	agctgtcaaa	catgagaatt	cttgaagacg	11940
aaagggcctc	gtgatacgcc	tatttttata	ggttaatgtc	atgataataa	tggtttctta	12000
gacgtcaggt	ggcacttttc	ggggaaatgt	gcgcggaacc	cctatttggt	tatttttcta	12060
aatacatcca	aatatgtatc	cgctcatgag	acaataaacc	tgataaatgc	ttcaataata	12120

ttgaaaaagg	aagagtatga	gtattcaaca	tttccgtgtc	gcccttattc	ccttttttgc	12180
ggcattttgc	cttctgtttt	ttgctcacc	agaaacgctg	gtgaaagtta	aagatgctga	12240
agatcagttg	gggtcacgag	tggtttacat	cgaactggat	ctcaacagcg	gtaagatcct	12300
tgagagtttt	cgccccgaag	aacgttttcc	aatgatgagc	acttttaaa	ttctgtatg	12360
tggtcggtta	ttatcccgtg	ttgacgccgg	gcaagagcaa	ctcggtcgcc	gcatacacta	12420
ttctcagaat	gacttgggtg	agtactcacc	agtcacagaa	aagcatctta	cggatggcat	12480
gacagtaaga	gaattatgca	gtgctgccat	aaccatgagt	gataacactg	cggccaactt	12540
acttctgaca	acgatcggag	gaccgaagga	gctaaccgct	tttttgaca	acatggggga	12600
tcattgtaact	cgccttgatc	gttgggaacc	ggagctgaat	gaagccatac	caaacgacga	12660
gctgtacacc	acgatgcctg	cagcaatggc	aacaacgttg	cgcaaaactat	taactggcga	12720
actacttact	ctagcttccc	ggcaacaatt	aatagactgg	atggaggcgg	ataaagtttg	12780
aggaccactt	ctgcgtcggg	cccttccggc	tggtctggtt	attgctgata	aacttggagc	12840
cgggtgagct	gggtctcggg	gtatcattgc	agcactgggg	ccagatggta	agccctcccg	12900
tatctgtagtt	atctacacga	cggggagtc	ggcaactatg	gatgaacgaa	atagacagat	12960
cgctgagata	gggtgcctcac	tgattaaagca	ttgtaactg	tcagaccaag	tttactcata	13020
tatactttag	attgatttaa	aacttcat	ttattttaa	aggatctagg	tgaagatcct	13080
ttttgataat	ctcatgacca	aaatccctta	acgtgagttt	tcgttccact	gagcgtcaga	13140
ccccgtagaa	aagatcaaag	gatcttcttg	agatcctttt	ttctgctg	taattctgtg	13200
cttgcaaaaca	aaaaaacccac	cgtaccagc	gggtgtttgt	ttgccggat	aagagctacc	13260
aaactttttt	ccgaaggtaa	ctggttccag	cagagcgag	ataccaaata	ctgtccttct	13320
agtgtagccg	tagttaggcc	accacttcaa	gaactctgta	gcaccgccta	catacctcgc	13380
tctgtctaata	ctgttaccag	tggtctgctg	cagtggcgat	aagtcgtgtc	ttaccgggtt	13440
ggactcaaga	cgatagttac	cggataaggc	gcagcggtcg	ggctgaacgg	ggggttcgtg	13500
cacacagccc	agcttgagc	gaacgacct	caccgaactg	agatacctac	agcgtgagct	13560
atgagaaagc	gccacgcttc	cgaaggag	aaaggcggac	aggtatccgg	taagcggcag	13620
ggtcggaaca	ggagagcgca	cgaaggagct	tcagggggga	aacgcctggg	atctttatag	13680
tcctgtcggg	tttcgccacc	tctgacttga	gcgtcgattt	ttgtgatgct	cgtcaggggg	13740
gcggagccta	tgaaaaaacg	ccagcaacgc	ggccttttta	cggttcctgg	ccttttctgt	13800
cgccgcgtgc	ggctgctgga	gatggcggac	gcgatggata	tggtctgcca	aggggtgggt	13860
tgcgcatcca	cagttctccg	caagaattga	ttggtcccaa	ttcttggagt	gggtaattccg	13920
ttagcgaggt	gccgcccgtt	tccattcagg	tcgaggtggc	cgggtcccat	gcaccgcgac	13980
gcaacgcggg	gagggcagaca	aggtataggg	cggcgccctac	aatccatgcc	aaccctgtcc	14040
atgtgtctgc	cgaggcggca	taaactcgccg	tgacgatcag	cgggtccagt	atcgaagtta	14100
ggctggtaag	agccgcgagc	gatccttgaa	gctgtccctg	atggctcgta	tctacttgc	14160
tgagacagcat	ggcctgcaac	gcgggcattcc	cgatgccgccc	ggaagcgaga	agaatcataa	14220
tggggaaggc	catccagcct	cgcgtcgcga	acgccagcaa	gacgtagccc	agcgcgtcgg	14280
ccgccatgcc	ctgcttcate	cccgtggccc	gttgctcgcg	tttgcgtggg	gtgtccccgg	14340
aagaaatata	tttgcatgtc	tttagttcta	tgatgacaca	aaccccgccc	agcgtcttgt	14400
cattggcgaa	ttcgaacacg	cagatgcagt	cggggcgggc	cgggtcccagg	tccacttccg	14460
atattaaggt	gacgcgtgtg	gcctcgaaaca	ctgcagcgacc	ctgcagcgac	ccgcttaaca	14520
gcgtcaacag	cgtgccgcag	atcccgggca	atgagatatg	aaaaagcctg	aactcaccgc	14580
gacgtctgtc	gagaagtttc	tgatcgaaaa	gttcgacagc	gtctccgacc	tgatgcagct	14640
ctcgaggggc	gaagaatctc	gtgctttcag	cttcgatgta	ggaggggctg	gatatgtcct	14700
gcgggtaaat	agctgcgcgg	atggtttcta	caaagatcgt	tagtgggagc	ggcactttgc	14760
atcgggccgg	ctccccgatt	ccggaagtgc	ttgacattgg	ggaattcagc	gagagcctga	14820
cctattgcat	ctcccgccgt	gcacaggggtg	tcacgttgca	agacctgcct	gaaaccgaac	14880
tgcccgctgt	tctgcagccg	gtcgcggagg	ccatggatgc	gatcgtctgc	gccgatctta	14940
gccagacgag	cgggttcggc	ccattcggac	cgcaaggaa	cgggtcaatac	actacatggc	15000
gtgatttcat	atgcgcgatt	gctgatcccc	atgtgtatca	ctggcaaaact	gtgatggacg	15060
acaccgtcag	tgctgcccgc	gcgcaggctc	tcgatgagct	gatgctttgg	gccgaggact	15120
gccccgaagt	ccggcacctc	gtgcacggcg	atctcggctc	caacaatgtc	ctgacggaca	15180
atggccgcat	aacagcggtc	attgactgga	cgcaggcgat	gttcggggat	tcccaatac	15240
aggtcgccaa	catcttcttc	tgaggccgt	ggttggcggg	tatggagcag	cagacgcgct	15300
acttcgagcg	gaggcatccg	gagcttgcag	gatcgccg	gctccggcg	tatatgctcc	15360
gcattgggtct	tgaccaactc	tatcagagct	tggttgacgg	caatttcgat	gatgcagctt	15420
gggagcaggg	tcgatgcgac	gcaatcgtcc	gatccggagc	cgggactgtc	ggcggtacac	15480
aaatcgcccc	cagaagcggg	gccgtctgga	ccgatggctg	tgtagaagta	ctcgccgata	15540
gtggaacggg	gagatggggg	aggctaactg	aaacacggaa	ggagacaata	ccggaaggaa	15600
cccgcgctat	gacggcaata	aaaagacaga	ataaaacgca	cgggtgttgg	gtcgtttgtt	15660
cataaacggg	gggttcgggtc	ccagggtctg	cactctgtcg	ataccccacc	gagaccccat	15720
tggggccaat	acggcccgct	ttcttctttt	tccccacccc	accccccaag	ttcgggtgaa	15780

ggcccagggc tcgcagccaa cgtcggggcg gcaggccctg ccatagccac tggcccctg 15840
 ggtagggac ggggtcctcc atggggaatg gtttatggtt cgtgggggtt attattttg 15900
 gcgttgctg ggggtctgtc catgactgga ctgagcagac agaccatgg tttttggatg 15960
 gcctgggcat ggaccgcatg tactggcgcg acacgaacac cgggcgtctg tggctgcaa 16020
 acacccccga cccccaaaaa ccaccgcgcg gatttctggc gtgccaagct agtcgaccaa 16080

<210> 49
 <211> 32
 <212> DNA
 <213> Homo sapiens

<400> 49
 cccggccgga ggggcagctt tgggagatg gt 32

<210> 50
 <211> 11
 <212> PRT
 <213> Homo sapiens

<400> 50
 Pro Gly Arg Arg Gly Ser Phe Val Glu Met Val
 1 5 10

<210> 51
 <211> 5
 <212> PRT
 <213> Homo sapiens

<400> 51
 Val Asn Leu Asp Ala
 1 5

<210> 52
 <211> 9
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic oligopeptide substrate

<400> 52
 Ser Glu Val Asn Leu Asp Ala Glu Phe
 1 5

<210> 53
 <211> 30
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic oligopeptide substrate

<400> 53
 Ala Asp Arg Gly Leu Thr Thr Arg Pro Gly Ser Gly Leu Thr Asn Ile
 1 5 10 15
 Lys Thr Glu Glu Ile Ser Glu Val Asn Leu Asp Ala Glu Phe
 20 25 30

<210> 54
 <211> 5
 <212> PRT
 <213> Homo sapiens

 <220>
 <223> Wild type Amyloid Precursor Protein cleavage site
 (fragment)

 <400> 54
 Val Lys Met Asp Ala
 1 5

 <210> 55
 <211> 24
 <212> PRT
 <213> Homo sapiens

 <400> 55
 Glu Thr Asp Glu Glu Pro Glu Glu Pro Gly Arg Arg Gly Ser Phe Val
 1 5 10 15
 Glu Met Val Asp Asn Leu Arg Gly
 20

 <210> 56
 <211> 15
 <212> PRT
 <213> Homo sapiens

 <400> 56
 Ile Gly Phe Ala Val Ser Ala Cys His Val His Asp Glu Phe Arg
 1 5 10 15

 <210> 57
 <211> 419
 <212> PRT
 <213> Homo sapiens

 <400> 57
 Met Ala Gln Ala Leu Pro Trp Leu Leu Leu Trp Met Gly Ala Gly Val
 1 5 10 15
 Leu Pro Ala His Gly Thr Gln His Gly Ile Arg Leu Pro Leu Arg Ser
 20 25 30
 Gly Leu Gly Gly Ala Pro Leu Gly Leu Arg Leu Pro Arg Glu Thr Asp
 35 40 45
 Glu Glu Pro Glu Glu Pro Gly Arg Arg Gly Ser Phe Val Glu Met Val
 50 55 60
 Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu Met Thr
 65 70 75 80
 Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp Thr Gly Ser
 85 90 95
 Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu His Arg Tyr
 100 105 110
 Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val
 115 120 125
 Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu Gly Thr Asp
 130 135 140
 Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg Ala Asn Ile
 145 150 155 160
 Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser Asn Trp
 165 170 175

Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg Pro Asp Asp
 180 185 190
 Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr His Val Pro
 195 200 205
 Asn Leu Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro Leu Asn Gln
 210 215 220
 Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile Gly Gly Ile
 225 230 235 240
 Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro Ile Arg Arg
 245 250 255
 Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile Asn Gly Gln
 260 265 270
 Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys Ser Ile Val
 275 280 285
 Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val Phe Glu Ala
 290 295 300
 Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys Phe Pro Asp
 305 310 315 320
 Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala Gly Thr Thr
 325 330 335
 Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met Gly Glu Val
 340 345 350
 Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln Tyr Leu Arg
 355 360 365
 Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr Lys Phe Ala
 370 375 380
 Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val Ile Met Glu
 385 390 395 400
 Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile Gly Phe Ala
 405 410 415
 Val Ser Ala

<210> 58
 <211> 407
 <212> PRT
 <213> Homo sapiens

<400> 58
 Glu Thr Asp Glu Glu Pro Glu Glu Pro Gly Arg Arg Gly Ser Phe Val
 1 5 10 15
 Glu Met Val Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val
 20 25 30
 Glu Met Thr Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp
 35 40 45
 Thr Gly Ser Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu
 50 55 60
 His Arg Tyr Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg
 65 70 75 80
 Lys Gly Val Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu
 85 90 95
 Gly Thr Asp Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg
 100 105 110
 Ala Asn Ile Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly
 115 120 125
 Ser Asn Trp Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg
 130 135 140
 Pro Asp Asp Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr
 145 150 155 160
 His Val Pro Asn Leu Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro
 165 170 175

Leu Asn Gln Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile
 180 185 190
 Gly Gly Ile Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro
 195 200 205
 Ile Arg Arg Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile
 210 215 220
 Asn Gly Gln Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys
 225 230 235 240
 Ser Ile Val Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val
 245 250 255
 Phe Glu Ala Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys
 260 265 270
 Phe Pro Asp Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala
 275 280 285
 Gly Thr Thr Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met
 290 295 300
 Gly Glu Val Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln
 305 310 315 320
 Tyr Leu Arg Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr
 325 330 335
 Lys Phe Ala Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val
 340 345 350
 Ile Met Glu Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile
 355 360 365
 Gly Phe Ala Val Ser Ala Cys His Val His Asp Glu Phe Arg Thr Ala
 370 375 380
 Ala Val Glu Gly Pro Phe Val Thr Leu Asp Met Glu Asp Cys Gly Tyr
 385 390 395 400
 Asn Ile Pro Gln Thr Asp Glu
 405

<210> 59
 <211> 452
 <212> PRT
 <213> Homo sapiens

<400> 59
 Met Ala Gln Ala Leu Pro Trp Leu Leu Leu Trp Met Gly Ala Gly Val
 1 5 10 15
 Leu Pro Ala His Gly Thr Gln His Gly Ile Arg Leu Pro Leu Arg Ser
 20 25 30
 Gly Leu Gly Gly Ala Pro Leu Gly Leu Arg Leu Pro Arg Glu Thr Asp
 35 40 45
 Glu Glu Pro Glu Glu Pro Gly Arg Arg Gly Ser Phe Val Glu Met Val
 50 55 60
 Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu Met Thr
 65 70 75 80
 Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp Thr Gly Ser
 85 90 95
 Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu His Arg Tyr
 100 105 110
 Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val
 115 120 125
 Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu Gly Thr Asp
 130 135 140
 Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg Ala Asn Ile
 145 150 155 160
 Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser Asn Trp
 165 170 175
 Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg Pro Asp Asp
 180 185 190

Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr His Val Pro
 195 200 205
 Asn Leu Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro Leu Asn Gln
 210 215 220
 Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile Gly Gly Ile
 225 230 235 240
 Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro Ile Arg Arg
 245 250 255
 Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile Asn Gly Gln
 260 265 270
 Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys Ser Ile Val
 275 280 285
 Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val Phe Glu Ala
 290 295 300
 Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys Phe Pro Asp
 305 310 315 320
 Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala Gly Thr Thr
 325 330 335
 Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met Gly Glu Val
 340 345 350
 Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln Tyr Leu Arg
 355 360 365
 Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr Lys Phe Ala
 370 375 380
 Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val Ile Met Glu
 385 390 395 400
 Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile Gly Phe Ala
 405 410 415
 Val Ser Ala Cys His Val His Asp Glu Phe Arg Thr Ala Ala Val Glu
 420 425 430
 Gly Pro Phe Val Thr Leu Asp Met Glu Asp Cys Gly Tyr Asn Ile Pro
 435 440 445
 Gln Thr Asp Glu
 450

<210> 60
 <211> 420
 <212> PRT
 <213> Homo sapiens

<400> 60
 Met Ala Gln Ala Leu Pro Trp Leu Leu Leu Trp Met Gly Ala Gly Val
 1 5 10 15
 Leu Pro Ala His Gly Thr Gln His Gly Ile Arg Leu Pro Leu Arg Ser
 20 25 30
 Gly Leu Gly Gly Ala Pro Leu Gly Leu Arg Leu Pro Arg Glu Thr Asp
 35 40 45
 Glu Glu Pro Glu Glu Pro Gly Arg Arg Gly Ser Phe Val Glu Met Val
 50 55 60
 Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu Met Thr
 65 70 75 80
 Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp Thr Gly Ser
 85 90 95
 Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu His Arg Tyr
 100 105 110
 Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val
 115 120 125
 Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu Gly Thr Asp
 130 135 140
 Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg Ala Asn Ile
 145 150 155 160

Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser Asn Trp
 165 170 175
 Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg Pro Asp Asp
 180 185 190
 Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr His Val Pro
 195 200 205
 Asn Leu Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro Leu Asn Gln
 210 215 220
 Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile Gly Gly Ile
 225 230 235 240
 Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro Ile Arg Arg
 245 250 255
 Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile Asn Gly Gln
 260 265 270
 Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys Ser Ile Val
 275 280 285
 Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val Phe Glu Ala
 290 295 300
 Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys Phe Pro Asp
 305 310 315 320
 Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala Gly Thr Thr
 325 330 335
 Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met Gly Glu Val
 340 345 350
 Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln Tyr Leu Arg
 355 360 365
 Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr Lys Phe Ala
 370 375 380
 Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val Ile Met Glu
 385 390 395 400
 Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile Gly Phe Ala
 405 410 415
 Val Ser Ala Cys
 420

<210> 61

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Synthetic peptide inhibitor

<221> MOD_RES

<222> 4

<223> Xaa = hydroxyethylene

<400> 61

Glu Val Met Xaa Ala Glu Phe

1

5

<210> 62

<211> 26

<212> PRT

<213> Homo sapiens

<400> 62

Leu Met Thr Ile Ala Tyr Val Met Ala Ala Ile Cys Ala Leu Phe Met

1

5

10

15

Leu Pro Leu Cys Leu Met Val Cys Gln Trp

20

25

<210> 63
 <211> 33
 <212> PRT
 <213> Homo sapiens

<220>
 <223> P26-P4'sw peptide substrate

<400> 63
 Cys Gly Gly Ala Asp Arg Gly Leu Thr Thr Arg Pro Gly Ser Gly Leu
 1 5 10 15
 Thr Asn Ile Lys Thr Glu Glu Ile Ser Glu Val Asn Leu Asp Ala Glu
 20 25 30
 Phe

<210> 64
 <211> 29
 <212> PRT
 <213> Homo sapiens

<220>
 <223> P26-P1' peptide substrate with CGG linker

<400> 64
 Cys Gly Gly Ala Asp Arg Gly Leu Thr Thr Arg Pro Gly Ser Gly Leu
 1 5 10 15
 Thr Asn Ile Lys Thr Glu Glu Ile Ser Glu Val Asn Leu
 20 25

<210> 65
 <211> 501
 <212> PRT
 <213> Mus musculus

<400> 65
 Met Ala Pro Ala Leu His Trp Leu Leu Leu Trp Val Gly Ser Gly Met
 1 5 10 15
 Leu Pro Ala Gln Gly Thr His Leu Gly Ile Arg Leu Pro Leu Arg Ser
 20 25 30
 Gly Leu Ala Gly Pro Pro Leu Gly Leu Arg Leu Pro Arg Glu Thr Asp
 35 40 45
 Glu Glu Ser Glu Glu Pro Gly Arg Arg Gly Ser Phe Val Glu Met Val
 50 55 60
 Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu Met Thr
 65 70 75 80
 Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp Thr Gly Ser
 85 90 95
 Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu His Arg Tyr
 100 105 110
 Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val
 115 120 125
 Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu Gly Thr Asp
 130 135 140
 Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg Ala Asn Ile
 145 150 155 160
 Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser Asn Trp
 165 170 175
 Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg Pro Asp Asp
 180 185 190

Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr His Ile Pro
 195 200 205
 Asn Ile Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro Leu Asn Gln
 210 215 220
 Thr Glu Ala Leu Ala Ser Val Gly Gly Ser Met Ile Ile Gly Gly Ile
 225 230 235 240
 Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro Ile Arg Arg
 245 250 255
 Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile Asn Gly Gln
 260 265 270
 Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys Ser Ile Val
 275 280 285
 Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val Phe Glu Ala
 290 295 300
 Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys Phe Pro Asp
 305 310 315 320
 Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala Gly Thr Thr
 325 330 335
 Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met Gly Glu Val
 340 345 350
 Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln Tyr Leu Arg
 355 360 365
 Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr Lys Phe Ala
 370 375 380
 Val Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val Ile Met Glu
 385 390 395 400
 Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile Gly Phe Ala
 405 410 415
 Val Ser Ala Cys His Val His Asp Glu Phe Arg Thr Ala Ala Val Glu
 420 425 430
 Gly Pro Phe Val Thr Ala Asp Met Glu Asp Cys Gly Tyr Asn Ile Pro
 435 440 445
 Gln Thr Asp Glu Ser Thr Leu Met Thr Ile Ala Tyr Val Met Ala Ala
 450 455 460
 Ile Cys Ala Leu Phe Met Leu Pro Leu Cys Leu Met Val Cys Gln Trp
 465 470 475 480
 Arg Cys Leu Arg Cys Leu Arg His Gln His Asp Asp Phe Gly Asp Asp
 485 490 495
 Ile Ser Leu Leu Lys
 500

<210> 66
 <211> 480
 <212> PRT
 <213> Homo sapiens

<400> 66
 Thr Gln His Gly Ile Arg Leu Pro Leu Arg Ser Gly Leu Gly Gly Ala
 1 5 10 15
 Pro Leu Gly Leu Arg Leu Pro Arg Glu Thr Asp Glu Glu Pro Glu Glu
 20 25 30
 Pro Gly Arg Arg Gly Ser Phe Val Glu Met Val Asp Asn Leu Arg Gly
 35 40 45
 Lys Ser Gly Gln Gly Tyr Tyr Val Glu Met Thr Val Gly Ser Pro Pro
 50 55 60
 Gln Thr Leu Asn Ile Leu Val Asp Thr Gly Ser Ser Asn Phe Ala Val
 65 70 75 80
 Gly Ala Ala Pro His Pro Phe Leu His Arg Tyr Tyr Gln Arg Gln Leu
 85 90 95
 Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val Tyr Val Pro Tyr Thr
 100 105 110

Gln Gly Lys Trp Glu Gly Glu Leu Gly Thr Asp Leu Val Ser Ile Pro
 115 120 125
 His Gly Pro Asn Val Thr Val Arg Ala Asn Ile Ala Ala Ile Thr Glu
 130 135 140
 Ser Asp Lys Phe Phe Ile Asn Gly Ser Asn Trp Glu Gly Ile Leu Gly
 145 150 155 160
 Leu Ala Tyr Ala Glu Ile Ala Arg Pro Asp Asp Ser Leu Glu Pro Phe
 165 170 175
 Phe Asp Ser Leu Val Lys Gln Thr His Val Pro Asn Leu Phe Ser Leu
 180 185 190
 Gln Leu Cys Gly Ala Gly Phe Pro Leu Asn Gln Ser Glu Val Leu Ala
 195 200 205
 Ser Val Gly Gly Ser Met Ile Ile Gly Gly Ile Asp His Ser Leu Tyr
 210 215 220
 Thr Gly Ser Leu Trp Tyr Thr Pro Ile Arg Arg Glu Trp Tyr Tyr Glu
 225 230 235 240
 Val Ile Ile Val Arg Val Glu Ile Asn Gly Gln Asp Leu Lys Met Asp
 245 250 255
 Cys Lys Glu Tyr Asn Tyr Asp Lys Ser Ile Val Asp Ser Gly Thr Thr
 260 265 270
 Asn Leu Arg Leu Pro Lys Lys Val Phe Glu Ala Ala Val Lys Ser Ile
 275 280 285
 Lys Ala Ala Ser Ser Thr Glu Lys Phe Pro Asp Gly Phe Trp Leu Gly
 290 295 300
 Glu Gln Leu Val Cys Trp Gln Ala Gly Thr Thr Pro Trp Asn Ile Phe
 305 310 315 320
 Pro Val Ile Ser Leu Tyr Leu Met Gly Glu Val Thr Asn Gln Ser Phe
 325 330 335
 Arg Ile Thr Ile Leu Pro Gln Gln Tyr Leu Arg Pro Val Glu Asp Val
 340 345 350
 Ala Thr Ser Gln Asp Asp Cys Tyr Lys Phe Ala Ile Ser Gln Ser Ser
 355 360 365
 Thr Gly Thr Val Met Gly Ala Val Ile Met Glu Gly Phe Tyr Val Val
 370 375 380
 Phe Asp Arg Ala Arg Lys Arg Ile Gly Phe Ala Val Ser Ala Cys His
 385 390 395 400
 Val His Asp Glu Phe Arg Thr Ala Ala Val Glu Gly Pro Phe Val Thr
 405 410 415
 Leu Asp Met Glu Asp Cys Gly Tyr Asn Ile Pro Gln Thr Asp Glu Ser
 420 425 430
 Thr Leu Met Thr Ile Ala Tyr Val Met Ala Ala Ile Cys Ala Leu Phe
 435 440 445
 Met Leu Pro Leu Cys Leu Met Val Cys Gln Trp Arg Cys Leu Arg Cys
 450 455 460
 Leu Arg Gln Gln His Asp Asp Phe Ala Asp Asp Ile Ser Leu Leu Lys
 465 470 475 480

<210> 67

<211> 444

<212> PRT

<213> Homo sapiens

<400> 67

Gly Ser Phe Val Glu Met Val Asp Asn Leu Arg Gly Lys Ser Gly Gln
 1 5 10 15
 Gly Tyr Tyr Val Glu Met Thr Val Gly Ser Pro Pro Gln Thr Leu Asn
 20 25 30
 Ile Leu Val Asp Thr Gly Ser Ser Asn Phe Ala Val Gly Ala Ala Pro
 35 40 45

His Pro Phe Leu His Arg Tyr Tyr Gln Arg Gln Leu Ser Ser Thr Tyr
 50 55 60
 Arg Asp Leu Arg Lys Gly Val Tyr Val Pro Tyr Thr Gln Gly Lys Trp
 65 70 75 80
 Glu Gly Glu Leu Gly Thr Asp Leu Val Ser Ile Pro His Gly Pro Asn
 85 90 95
 Val Thr Val Arg Ala Asn Ile Ala Ala Ile Thr Glu Ser Asp Lys Phe
 100 105 110
 Phe Ile Asn Gly Ser Asn Trp Glu Gly Ile Leu Gly Leu Ala Tyr Ala
 115 120 125
 Glu Ile Ala Arg Pro Asp Asp Ser Leu Glu Pro Phe Phe Asp Ser Leu
 130 135 140
 Val Lys Gln Thr His Val Pro Asn Leu Phe Ser Leu Gln Leu Cys Gly
 145 150 155 160
 Ala Gly Phe Pro Leu Asn Gln Ser Glu Val Leu Ala Ser Val Gly Gly
 165 170 175
 Ser Met Ile Ile Gly Gly Ile Asp His Ser Leu Tyr Thr Gly Ser Leu
 180 185 190
 Trp Tyr Thr Pro Ile Arg Arg Glu Trp Tyr Tyr Glu Val Ile Ile Val
 195 200 205
 Arg Val Glu Ile Asn Gly Gln Asp Leu Lys Met Asp Cys Lys Glu Tyr
 210 215 220
 Asn Tyr Asp Lys Ser Ile Val Asp Ser Gly Thr Thr Asn Leu Arg Leu
 225 230 235 240
 Pro Lys Lys Val Phe Glu Ala Ala Val Lys Ser Ile Lys Ala Ala Ser
 245 250 255
 Ser Thr Glu Lys Phe Pro Asp Gly Phe Trp Leu Gly Glu Gln Leu Val
 260 265 270
 Cys Trp Gln Ala Gly Thr Thr Pro Trp Asn Ile Phe Pro Val Ile Ser
 275 280 285
 Leu Tyr Leu Met Gly Glu Val Thr Asn Gln Ser Phe Arg Ile Thr Ile
 290 295 300
 Leu Pro Gln Gln Tyr Leu Arg Pro Val Glu Asp Val Ala Thr Ser Gln
 305 310 315 320
 Asp Asp Cys Tyr Lys Phe Ala Ile Ser Gln Ser Ser Thr Gly Thr Val
 325 330 335
 Met Gly Ala Val Ile Met Glu Gly Phe Tyr Val Val Phe Asp Arg Ala
 340 345 350
 Arg Lys Arg Ile Gly Phe Ala Val Ser Ala Cys His Val His Asp Glu
 355 360 365
 Phe Arg Thr Ala Ala Val Glu Gly Pro Phe Val Thr Leu Asp Met Glu
 370 375 380
 Asp Cys Gly Tyr Asn Ile Pro Gln Thr Asp Glu Ser Thr Leu Met Thr
 385 390 395 400
 Ile Ala Tyr Val Met Ala Ala Ile Cys Ala Leu Phe Met Leu Pro Leu
 405 410 415
 Cys Leu Met Val Cys Gln Trp Arg Cys Leu Arg Cys Leu Arg Gln Gln
 420 425 430
 His Asp Asp Phe Ala Asp Asp Ile Ser Leu Leu Lys
 435 440

<210> 68
 <211> 395
 <212> PRT
 <213> Homo sapiens

<400> 68
 Gly Ser Phe Val Glu Met Val Asp Asn Leu Arg Gly Lys Ser Gly Gln
 1 5 10 15
 Gly Tyr Tyr Val Glu Met Thr Val Gly Ser Pro Pro Gln Thr Leu Asn
 20 25 30

Ile Leu Val Asp Thr Gly Ser Ser Asn Phe Ala Val Gly Ala Ala Pro
 35 40 45
 His Pro Phe Leu His Arg Tyr Tyr Gln Arg Gln Leu Ser Ser Thr Tyr
 50 55 60
 Arg Asp Leu Arg Lys Gly Val Tyr Val Pro Tyr Thr Gln Gly Lys Trp
 65 70 75 80
 Glu Gly Glu Leu Gly Thr Asp Leu Val Ser Ile Pro His Gly Pro Asn
 85 90 95
 Val Thr Val Arg Ala Asn Ile Ala Ala Ile Thr Glu Ser Asp Lys Phe
 100 105 110
 Phe Ile Asn Gly Ser Asn Trp Glu Gly Ile Leu Gly Leu Ala Tyr Ala
 115 120 125
 Glu Ile Ala Arg Pro Asp Asp Ser Leu Glu Pro Phe Phe Asp Ser Leu
 130 135 140
 Val Lys Gln Thr His Val Pro Asn Leu Phe Ser Leu Gln Leu Cys Gly
 145 150 155 160
 Ala Gly Phe Pro Leu Asn Gln Ser Glu Val Leu Ala Ser Val Gly Gly
 165 170 175
 Ser Met Ile Ile Gly Gly Ile Asp His Ser Leu Tyr Thr Gly Ser Leu
 180 185 190
 Trp Tyr Thr Pro Ile Arg Arg Glu Trp Tyr Tyr Glu Val Ile Ile Val
 195 200 205
 Arg Val Glu Ile Asn Gly Gln Asp Leu Lys Met Asp Cys Lys Glu Tyr
 210 215 220
 Asn Tyr Asp Lys Ser Ile Val Asp Ser Gly Thr Thr Asn Leu Arg Leu
 225 230 235 240
 Pro Lys Lys Val Phe Glu Ala Ala Val Lys Ser Ile Lys Ala Ala Ser
 245 250 255
 Ser Thr Glu Lys Phe Pro Asp Gly Phe Trp Leu Gly Glu Gln Leu Val
 260 265 270
 Cys Trp Gln Ala Gly Thr Thr Pro Trp Asn Ile Phe Pro Val Ile Ser
 275 280 285
 Leu Tyr Leu Met Gly Glu Val Thr Asn Gln Ser Phe Arg Ile Thr Ile
 290 295 300
 Leu Pro Gln Gln Tyr Leu Arg Pro Val Glu Asp Val Ala Thr Ser Gln
 305 310 315 320
 Asp Asp Cys Tyr Lys Phe Ala Ile Ser Gln Ser Ser Thr Gly Thr Val
 325 330 335
 Met Gly Ala Val Ile Met Glu Gly Phe Tyr Val Val Phe Asp Arg Ala
 340 345 350
 Arg Lys Arg Ile Gly Phe Ala Val Ser Ala Cys His Val His Asp Glu
 355 360 365
 Phe Arg Thr Ala Ala Val Glu Gly Pro Phe Val Thr Leu Asp Met Glu
 370 375 380
 Asp Cys Gly Tyr Asn Ile Pro Gln Thr Asp Glu
 385 390 395

<210> 69

<211> 439

<212> PRT

<213> Homo sapiens

<400> 69

Met Val Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu
 1 5 10 15
 Met Thr Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp Thr
 20 25 30
 Gly Ser Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu His
 35 40 45
 Arg Tyr Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys
 50 55 60

Gly Val Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu Gly
 65 70 75 80
 Thr Asp Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg Ala
 85 90 95
 Asn Ile Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser
 100 105 110
 Asn Trp Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg Pro
 115 120 125
 Asp Asp Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr His
 130 135 140
 Val Pro Asn Leu Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro Leu
 145 150 155 160
 Asn Gln Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile Gly
 165 170 175
 Gly Ile Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro Ile
 180 185 190
 Arg Arg Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile Asn
 195 200 205
 Gly Gln Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys Ser
 210 215 220
 Ile Val Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val Phe
 225 230 235 240
 Glu Ala Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys Phe
 245 250 255
 Pro Asp Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala Gly
 260 265 270
 Thr Thr Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met Gly
 275 280 285
 Glu Val Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln Tyr
 290 295 300
 Leu Arg Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr Lys
 305 310 315 320
 Phe Ala Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val Ile
 325 330 335
 Met Glu Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile Gly
 340 345 350
 Phe Ala Val Ser Ala Cys His Val His Asp Glu Phe Arg Thr Ala Ala
 355 360 365
 Val Glu Gly Pro Phe Val Thr Leu Asp Met Glu Asp Cys Gly Tyr Asn
 370 375 380
 Ile Pro Gln Thr Asp Glu Ser Thr Leu Met Thr Ile Ala Tyr Val Met
 385 390 395 400
 Ala Ala Ile Cys Ala Leu Phe Met Leu Pro Leu Cys Leu Met Val Cys
 405 410 415
 Gln Trp Arg Cys Leu Arg Cys Leu Arg Gln Gln His Asp Asp Phe Ala
 420 425 430
 Asp Asp Ile Ser Leu Leu Lys
 435

<210> 70

<211> 390

<212> PRT

<213> Homo sapiens

<400> 70

Met Val Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu
 1 5 10 15
 Met Thr Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp Thr
 20 25 30
 Gly Ser Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu His
 35 40 45

Arg Tyr Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys
 50 55 60
 Gly Val Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu Gly
 65 70 75 80
 Thr Asp Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg Ala
 85 90 95
 Asn Ile Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser
 100 105 110
 Asn Trp Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg Pro
 115 120 125
 Asp Asp Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr His
 130 135 140
 Val Pro Asn Leu Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro Leu
 145 150 155 160
 Asn Gln Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile Gly
 165 170 175
 Gly Ile Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro Ile
 180 185 190
 Arg Arg Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile Asn
 195 200 205
 Gly Gln Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys Ser
 210 215 220
 Ile Val Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val Phe
 225 230 235 240
 Glu Ala Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys Phe
 245 250 255
 Pro Asp Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala Gly
 260 265 270
 Thr Thr Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met Gly
 275 280 285
 Glu Val Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln Tyr
 290 295 300
 Leu Arg Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr Lys
 305 310 315 320
 Phe Ala Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val Ile
 325 330 335
 Met Glu Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile Gly
 340 345 350
 Phe Ala Val Ser Ala Cys His Val His Asp Glu Phe Arg Thr Ala Ala
 355 360 365
 Val Glu Gly Pro Phe Val Thr Leu Asp Met Glu Asp Cys Gly Tyr Asn
 370 375 380
 Ile Pro Gln Thr Asp Glu
 385 390

<210> 71
 <211> 374
 <212> PRT
 <213> Homo sapiens

<400> 71
 Glu Thr Asp Glu Glu Pro Glu Glu Pro Gly Arg Arg Gly Ser Phe Val
 1 5 10 15
 Glu Met Val Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val
 20 25 30
 Glu Met Thr Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp
 35 40 45
 Thr Gly Ser Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu
 50 55 60
 His Arg Tyr Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg
 65 70 75 80

Lys Gly Val Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu
 85 90 95
 Gly Thr Asp Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg
 100 105 110
 Ala Asn Ile Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly
 115 120 125
 Ser Asn Trp Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg
 130 135 140
 Pro Asp Asp Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr
 145 150 155 160
 His Val Pro Asn Leu Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro
 165 170 175
 Leu Asn Gln Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile
 180 185 190
 Gly Gly Ile Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro
 195 200 205
 Ile Arg Arg Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile
 210 215 220
 Asn Gly Gln Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys
 225 230 235 240
 Ser Ile Val Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val
 245 250 255
 Phe Glu Ala Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys
 260 265 270
 Phe Pro Asp Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala
 275 280 285
 Gly Thr Thr Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met
 290 295 300
 Gly Glu Val Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln
 305 310 315 320
 Tyr Leu Arg Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr
 325 330 335
 Lys Phe Ala Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val
 340 345 350
 Ile Met Glu Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile
 355 360 365
 Gly Phe Ala Val Ser Ala
 370

<210> 72
 <211> 14
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> P10-P4'staD-V peptide inhibitor

<221> MOD_RES
 <222> 10
 <223> Xaa is statine moiety

<400> 72
 Lys Thr Glu Glu Ile Ser Glu Val Asn Xaa Val Ala Glu Phe
 1 5 10

<210> 73
 <211> 9
 <212> PRT
 <213> Artificial Sequence

<220>

<223> P4-P4 staD-V peptide inhibitor

<221> MOD_RES

<222> 5

<223> Xaa is statine moiety

<400> 73

Ser Glu Val Asn Xaa Val Ala Glu Phe

1

5

<210> 74

<211> 431

<212> PRT

<213> Homo sapiens

<400> 74

Thr Gln His Gly Ile Arg Leu Pro Leu Arg Ser Gly Leu Gly Gly Ala
1 5 10 15
Pro Leu Gly Leu Arg Leu Pro Arg Glu Thr Asp Glu Glu Pro Glu Glu
20 25 30
Pro Gly Arg Arg Gly Ser Phe Val Glu Met Val Asp Asn Leu Arg Gly
35 40 45
Lys Ser Gly Gln Gly Tyr Tyr Val Glu Met Thr Val Gly Ser Pro Pro
50 55 60
Gln Thr Leu Asn Ile Leu Val Asp Thr Gly Ser Ser Asn Phe Ala Val
65 70 75 80
Gly Ala Ala Pro His Pro Phe Leu His Arg Tyr Tyr Gln Arg Gln Leu
85 90 95
Ser Ser Thr Tyr Arg Asp Leu Arg Lys Gly Val Tyr Val Pro Tyr Thr
100 105 110
Gln Gly Lys Trp Glu Gly Glu Leu Gly Thr Asp Leu Val Ser Ile Pro
115 120 125
His Gly Pro Asn Val Thr Val Arg Ala Asn Ile Ala Ala Ile Thr Glu
130 135 140
Ser Asp Lys Phe Phe Ile Asn Gly Ser Asn Trp Glu Gly Ile Leu Gly
145 150 155 160
Leu Ala Tyr Ala Glu Ile Ala Arg Pro Asp Asp Ser Leu Glu Pro Phe
165 170 175
Phe Asp Ser Leu Val Lys Gln Thr His Val Pro Asn Leu Phe Ser Leu
180 185 190
Gln Leu Cys Gly Ala Gly Phe Pro Leu Asn Gln Ser Glu Val Leu Ala
195 200 205
Ser Val Gly Gly Ser Met Ile Ile Gly Gly Ile Asp His Ser Leu Tyr
210 215 220
Thr Gly Ser Leu Trp Tyr Thr Pro Ile Arg Arg Glu Trp Tyr Tyr Glu
225 230 235 240
Val Ile Ile Val Arg Val Glu Ile Asn Gly Gln Asp Leu Lys Met Asp
245 250 255
Cys Lys Glu Tyr Asn Tyr Asp Lys Ser Ile Val Asp Ser Gly Thr Thr
260 265 270
Asn Leu Arg Leu Pro Lys Lys Val Phe Glu Ala Ala Val Lys Ser Ile
275 280 285
Lys Ala Ala Ser Ser Thr Glu Lys Phe Pro Asp Gly Phe Trp Leu Gly
290 295 300
Glu Gln Leu Val Cys Trp Gln Ala Gly Thr Thr Pro Trp Asn Ile Phe
305 310 315 320
Pro Val Ile Ser Leu Tyr Leu Met Gly Glu Val Thr Asn Gln Ser Phe
325 330 335
Arg Ile Thr Ile Leu Pro Gln Gln Tyr Leu Arg Pro Val Glu Asp Val
340 345 350

Ala Thr Ser Gln Asp Asp Cys Tyr Lys Phe Ala Ile Ser Gln Ser Ser
 355 360 365
 Thr Gly Thr Val Met Gly Ala Val Ile Met Glu Gly Phe Tyr Val Val
 370 375 380
 Phe Asp Arg Ala Arg Lys Arg Ile Gly Phe Ala Val Ser Ala Cys His
 385 390 395 400
 Val His Asp Glu Phe Arg Thr Ala Ala Val Glu Gly Pro Phe Val Thr
 405 410 415
 Leu Asp Met Glu Asp Cys Gly Tyr Asn Ile Pro Gln Thr Asp Glu
 420 425 430

<210> 75

<211> 361

<212> PRT

<213> Homo sapiens

<400> 75

Met Val Asp Asn Leu Arg Gly Lys Ser Gly Gln Gly Tyr Tyr Val Glu
 1 5 10 15
 Met Thr Val Gly Ser Pro Pro Gln Thr Leu Asn Ile Leu Val Asp Thr
 20 25 30
 Gly Ser Ser Asn Phe Ala Val Gly Ala Ala Pro His Pro Phe Leu His
 35 40 45
 Arg Tyr Tyr Gln Arg Gln Leu Ser Ser Thr Tyr Arg Asp Leu Arg Lys
 50 55 60
 Gly Val Tyr Val Pro Tyr Thr Gln Gly Lys Trp Glu Gly Glu Leu Gly
 65 70 75 80
 Thr Asp Leu Val Ser Ile Pro His Gly Pro Asn Val Thr Val Arg Ala
 85 90 95
 Asn Ile Ala Ala Ile Thr Glu Ser Asp Lys Phe Phe Ile Asn Gly Ser
 100 105 110
 Asn Trp Glu Gly Ile Leu Gly Leu Ala Tyr Ala Glu Ile Ala Arg Pro
 115 120 125
 Asp Asp Ser Leu Glu Pro Phe Phe Asp Ser Leu Val Lys Gln Thr His
 130 135 140
 Val Pro Asn Leu Phe Ser Leu Gln Leu Cys Gly Ala Gly Phe Pro Leu
 145 150 155 160
 Asn Gln Ser Glu Val Leu Ala Ser Val Gly Gly Ser Met Ile Ile Gly
 165 170 175
 Gly Ile Asp His Ser Leu Tyr Thr Gly Ser Leu Trp Tyr Thr Pro Ile
 180 185 190
 Arg Arg Glu Trp Tyr Tyr Glu Val Ile Ile Val Arg Val Glu Ile Asn
 195 200 205
 Gly Gln Asp Leu Lys Met Asp Cys Lys Glu Tyr Asn Tyr Asp Lys Ser
 210 215 220
 Ile Val Asp Ser Gly Thr Thr Asn Leu Arg Leu Pro Lys Lys Val Phe
 225 230 235 240
 Glu Ala Ala Val Lys Ser Ile Lys Ala Ala Ser Ser Thr Glu Lys Phe
 245 250 255
 Pro Asp Gly Phe Trp Leu Gly Glu Gln Leu Val Cys Trp Gln Ala Gly
 260 265 270
 Thr Thr Pro Trp Asn Ile Phe Pro Val Ile Ser Leu Tyr Leu Met Gly
 275 280 285
 Glu Val Thr Asn Gln Ser Phe Arg Ile Thr Ile Leu Pro Gln Gln Tyr
 290 295 300
 Leu Arg Pro Val Glu Asp Val Ala Thr Ser Gln Asp Asp Cys Tyr Lys
 305 310 315 320
 Phe Ala Ile Ser Gln Ser Ser Thr Gly Thr Val Met Gly Ala Val Ile
 325 330 335
 Met Glu Gly Phe Tyr Val Val Phe Asp Arg Ala Arg Lys Arg Ile Gly
 340 345 350

Phe Ala Val Ser Ala Cys His Val His
355 360

<210> 76
<211> 63
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<222> (1)...(63)
<223> n = A,T,C or G

<400> 76
garacngayg argarccnga rgarccnggn mgnmgnggnw snttygtnga ratggtnGay 60
aay 63

<210> 77
<211> 21
<212> PRT
<213> Homo sapiens

<400> 77
Glu Thr Asp Glu Glu Pro Glu Glu Pro Gly Arg Arg Gly Ser Phe Val
1 5 10 15
Glu Met Val Asp Asn
20

<210> 78
<211> 7
<212> PRT
<213> Artificial Sequence

<220>
<223> Peptide inhibitor P3-P4' XD-V

<221> MOD_RES
<222> 3
<223> Xaa is hydroxyethylene or statine

<400> 78
Val Met Xaa Val Ala Glu Phe
1 5

<210> 79
<211> 11
<212> PRT
<213> Homo sapiens

<400> 79
Pro Glu Glu Pro Gly Arg Arg Gly Ser Phe Val
1 5 10

<210> 80
<211> 419
<212> DNA
<213> Artificial Sequence

<220>
<223> nucleotide insert in vector pCF

```

<400> 80
ctgttgggct cgcggttgag gacaaactct tcgcggtctt tccagtactc ttggatcgga      60
aaccgcgcgg cctccgaacg gtactccgcc accgaggagc ctgagcgagt ccgcatcgac      120
cggatcggaa aacctctcga ctgttggggt gagtactccc tctcaaaagc gggcatgact      180
tctgcgctaa gattgtcagt ttccaaaaac gaggaggatt tgatattcac ctggcccgcg      240
gtgatgcctt tgagggtggc cgcgtccatc tggtcagaaa agacaatctt tttgtgtgca      300
agcttgaggt gtggcaggct tgagatctgg ccatacactt gagtgacaat gacatccact      360
ttgcctttct ctccacaggc gtccactccc aggtccaact gcaggtcgac tctagaccc      419

```

```

<210> 81
<211> 8
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Peptide inhibitor P4-P4' XD-V

```

```

<221> MOD_RES
<222> 4
<223> Xaa is hydroxyethylene or statine

```

```

<400> 81
Glu Val Met Xaa Val Ala Glu Phe
1                5

```

```

<210> 82
<211> 9
<212> PRT
<213> Homo sapiens

```

```

<220>
<223> APP fragment P5-P4' wt

```

```

<400> 82
Ser Glu Val Lys Met Asp Ala Glu Phe
1                5

```

```

<210> 83
<211> 9
<212> PRT
<213> Homo sapiens

```

```

<220>
<223> APP fragment P5-P4'wt

```

```

<400> 83
Ser Glu Val Asn Leu Asp Ala Glu Phe
1                5

```

```

<210> 84
<211> 9
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> APP fragment

```

```

<400> 84
Ser Glu Val Lys Leu Asp Ala Glu Phe
1                5

```

<210> 85
 <211> 9
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> APP fragment
 <400> 85
 Ser Glu Val Lys Phe Asp Ala Glu Phe
 1 5
 <210> 86
 <211> 9
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> APP fragment
 <400> 86
 Ser Glu Val Asn Phe Asp Ala Glu Phe
 1 5
 <210> 87
 <211> 9
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> APP fragment
 <400> 87
 Ser Glu Val Lys Met Ala Ala Glu Phe
 1 5
 <210> 88
 <211> 9
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> APP fragment
 <400> 88
 Ser Glu Val Asn Leu Ala Ala Glu Phe
 1 5
 <210> 89
 <211> 9
 <212> PRT
 <213> Artificial Sequence
 <220>
 <223> APP fragment
 <400> 89
 Ser Glu Val Lys Leu Ala Ala Glu Phe
 1 5
 <210> 90

<211> 9
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> APP fragment

 <400> 90
 Ser Glu Val Lys Met Leu Ala Glu Phe
 1 5

 <210> 91
 <211> 9
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> APP fragment

 <400> 91
 Ser Glu Val Asn Leu Leu Ala Glu Phe
 1 5

 <210> 92
 <211> 9
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> APP fragment

 <400> 92
 Ser Glu Val Lys Leu Leu Ala Glu Phe
 1 5

 <210> 93
 <211> 9
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> APP fragment

 <400> 93
 Ser Glu Val Lys Phe Ala Ala Glu Phe
 1 5

 <210> 94
 <211> 9
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> APP fragment

 <400> 94
 Ser Glu Val Asn Phe Ala Ala Glu Phe
 1 5

 <210> 95
 <211> 9

<212> PRT
<213> Artificial Sequence

<220>
<223> APP fragment

<400> 95
Ser Glu Val Lys Phe Leu Ala Glu Phe
1 5

<210> 96
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> APP fragment

<400> 96
Ser Glu Val Asn Phe Leu Ala Glu Phe
1 5

<210> 97
<211> 14
<212> PRT
<213> Artificial Sequence

<220>
<223> APP-derived fragment P10-P4' (D-V)

<400> 97
Lys Thr Glu Glu Ile Ser Glu Val Asn Leu Val Ala Glu Phe
1 5 10

<210> 98
<211> 35
<212> DNA
<213> Homo sapiens

<400> 98
cccgaagagc ccggccggag gggcagcttt gtcga

35

<210> 99
<211> 11
<212> PRT
<213> Homo sapiens

<400> 99
Glu Thr Asp Glu Glu Pro Glu Glu Pro Gly Arg
1 5 10

<210> 100
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Recombinant 293T cells

<400> 100

Thr Gln His Gly Ile Arg Leu Pro Leu Arg
1 5 10

<210> 101
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> Recombinant 293T cells

<400> 101
Met Val Asp Asn Leu Arg Gly Lys Ser
1 5

<210> 102
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Recombinant CosA2 cells

<400> 102
Gly Ser Phe Val Glu Met Val Asp Asn Leu
1 5 10

<210> 103
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> APP substrate fragment:WT Sequence

<400> 103
Val Lys Met Asp
1

<210> 104
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> APP substrate fragment:Swedish Sequence

<400> 104
Val Asn Leu Asp
1

1. The first part of the report is a summary of the work done during the year.

2. The second part is a detailed account of the work done during the year.

3. The third part is a summary of the work done during the year.

4. The fourth part is a summary of the work done during the year.

5. The fifth part is a summary of the work done during the year.

6. The sixth part is a summary of the work done during the year.

7. The seventh part is a summary of the work done during the year.

8. The eighth part is a summary of the work done during the year.

9. The ninth part is a summary of the work done during the year.

10. The tenth part is a summary of the work done during the year.

11. The eleventh part is a summary of the work done during the year.

12. The twelfth part is a summary of the work done during the year.

13. The thirteenth part is a summary of the work done during the year.

14. The fourteenth part is a summary of the work done during the year.

15. The fifteenth part is a summary of the work done during the year.

16. The sixteenth part is a summary of the work done during the year.

17. The seventeenth part is a summary of the work done during the year.

18. The eighteenth part is a summary of the work done during the year.

19. The nineteenth part is a summary of the work done during the year.

20. The twentieth part is a summary of the work done during the year.